



# DCC User's Guide

## DCCs for Radionuclides

[PDF of User's Guide.](#)

Welcome to the EPA's "Dose Compliance Concentrations for Radionuclides at Superfund Sites" (DCC) user's guide. Here you will find descriptions, equations and default exposure parameters used to calculate the dose-based DCCs. Additional guidance is also provided on sources of parameters and proper DCC use. It is suggested that users read the [DCC FAQ](#) page before proceeding. The user guide is extensive so please use the "Open All Sections" and "Close All Sections" links below as needed. Individual sections can be opened and closed by clicking on the section titles. Before proceeding through the user's guide please read the [Disclaimer](#).

- [Home Page](#)
- [User's Guide](#)
- [What's New](#)
- [Frequent Questions](#)
- [Equations](#)
- [Calculator](#)
- [Generic Tables](#)

[Open All Sections](#) | [Close All Sections](#)

## Disclaimer

This guidance document sets forth recommended approaches based on EPA's best thinking to date with respect to dose assessment for response actions at CERCLA sites. This document does not establish binding rules. Alternative dose assessment approaches may be more appropriate at specific sites (e.g., where site circumstances do not match the underlying assumptions, conditions, and models of the guidance). The decision to use an alternative approach and a description of any such approach should be placed in the Administrative Record for the site. Accordingly, if comments are received at individual sites questioning the use of the approaches recommended in this guidance, the comments should be considered and an explanation provided for the selected approach.

The policies set out in the Radionuclide ARAR DCC User Guide provide guidance to EPA staff. It also educates the public and regulated community on how EPA intends the National Oil and Hazardous Substances Pollution Contingency Plan (NCP) be implemented. EPA may change this guidance in the future, as appropriate. This calculator is intended for use by risk assessors, health physicists, and other qualified environmental protection specialists.

It should also be noted that DCCs do not address human cancer risk, noncancer toxicity, or potential ecological risk. Of the radionuclides generally found at CERCLA sites, only uranium has potentially significant noncancer toxicity. When assessing sites with uranium as a contaminant, it may also be necessary to consider the noncancer toxicity of uranium using other tools, such as EPA's Regional Screening Levels ([RSLs](#)) for Chemical Contaminants at Superfund Sites electronic calculator for uranium in soil, water, or air, and the [WTC](#) for uranium inside buildings. EPA's [SDCC](#) Calculator should be used to assess radionuclide dose for hard outside surfaces, and the [BDCC](#) Calculator for radionuclide dose inside buildings. EPA's [PRG](#) Calculator should be used to assess radionuclide cancer risk for soil, water, and air, [BPRG](#) Calculator for radionuclide cancer risk inside buildings, and the [SPRG](#) Calculator for radionuclide cancer risk for hard outside surfaces. Similarly, some sites with radiological contaminants in sensitive ecological settings may also need to be evaluated for potential ecological risk. EPA's guidance "[Ecological Risk Assessment Guidance for Superfund: Process for Designing and Conducting Ecological Risk Assessment](#)" contains an eight step process for using benchmarks for ecological effects in the remedy selection process.

This web calculator may be used to develop generic DCCs for radionuclides for several different exposure scenarios. The calculator is flexible and may be used to derive site-specific DCCs as more site characterization is obtained (EPA 2000a). Models reviewed by EPA in the [Soil Screening Guidance for Radionuclides: Technical Background Document](#) are presented in Section 3-2. This report provides a detailed technical analysis of five unsaturated zone fate and transport models for radionuclides. This report supports the information provided in [Part 3 - Unsaturated Zone Models for Radionuclide Fate and Transport](#) [PDF 383KB, 25 pages] of the [Soil Guidance for Radionuclides: Technical Background Document](#) on determining the general applicability of the models to subsurface conditions and assessing each model's potential applicability to the soil screening process.

## 1. Introduction

Radionuclide ARAR Dose Compliance Concentrations (DCCs) are dose concentrations derived from standardized equations that combine exposure information, assumptions, and dose conversion factors (DCFs).

The Dose Assessment Guidance is a tool that the U.S. Environmental Protection Agency developed to help standardize the evaluation and cleanup of radioactively contaminated sites when doses are being assessed. This guidance provides a methodology for radiation professionals to calculate dose-based, site-specific, dose compliance concentrations (DCCs) for radionuclides complying with a dose-based standard as an ARAR. This guidance supersedes the dose assessment methodology contained in the "Risk Assessment Guidance for Superfund Volume I Human Health Evaluation Manual (Part A) (EPA/540/1-89/002).

A number of different radiation standards may be used as Applicable or Relevant and Appropriate Requirements (ARARs) to establish cleanup levels at a site. Cleanup levels may be based on a number of Federal or State ARARs. Federal standards expressed in terms of dose that are potential ARARs at CERCLA sites include 40 CFR Part 190, "Environmental Radiation Protection Standards for Nuclear Power Operations," 40 CFR Part 191, "Environmental Radiation Protection Standards for Management and Disposal of Spent Nuclear Fuel, High-Level and Transuranic Radioactive Wastes," or 10 CFR Part 61, "Licensing Requirements for Land Disposal of Radioactive Waste," among others.

One set of radiation standards consists of a combination of whole body and critical organ dose annual limits, generally either (1) 25 mrem to the whole body, 75 mrem to the thyroid, and 25 mrem to any other critical organ besides the thyroid or (2) 25 mrem/year to the whole body and 75 mrem/year to any critical organ (including the thyroid). Another set of standards consists of a single limit (e.g., 10 mrem/year). The type of dose limit used in the standard would be the same type of dose methodology used for dose assessments to demonstrate ARAR compliance.

The approach to dose limitation and the methods used to calculate doses have evolved over time. The first two radiation protection standards listed above (the 25/75/25 and 25/75 mrem annual dose limits) are based on the older, critical organ concept of dose limitation. This approach limits dose

and long-term effects to a specific target tissue or organ (e.g., the thyroid), the most radiosensitive tissue or organ, or the tissue or organ receiving the highest dose. Under this approach, introduced in 1959 by the International Commission on Radiological Protection (ICRP) in its Publication 2, "Report of Committee II on Permissible Dose for Internal Radiation" (ICRP, 1959), the dose to an organ from internally-deposited radionuclides is calculated separately from the dose due to external exposure, and the whole body is essentially treated as one of the critical organs.

Later, standards were based on the effective dose equivalent concept of dose limitation, introduced in 1977 by the ICRP in its Publication 26, "Recommendations of the International Commission on Radiological Protection" (ICRP, 1977). The effective dose equivalent approach accounts for the differences in the cancer induction rates in organs and tissues subjected to equal doses of radiation and normalizes these doses and effects on a whole body basis. Under this approach, the effective dose equivalent dose is calculated as the weighted sum of the committed dose equivalents (from ingested and inhaled radionuclides) and the dose equivalent (for external exposure from photon-emitting radionuclides) to all organs and tissues. The weighting factors used in these calculations are organ-specific and correspond to the fractional contribution of each organ or tissue to the total risk of fatal cancers when the body is uniformly irradiated. Thus, the summation of all organ and tissue factors is equal to one.

ICRP has since updated the effective dose equivalent concept with the introduction of effective dose quantity in its Publication 60, "1990 recommendations of the International Commission on Radiological Protection" (ICRP, 1991). While similar to the effective dose equivalent approach, the effective dose quantity incorporates updated scientific information in the dose conversion factors. Effective dose quantity incorporates a greater number of organs, updated information on organ-specific risk, and age-specific dose coefficients for internal exposure that incorporate new physiologically-based biokinetic models.

ICRP Publication 107 (ICRP 2008) provides an electronic database of the physical data for calculations of radionuclide-specific protection and operational quantities. This database supersedes the data of ICRP 38 and will be used in future ICRP publications of dose coefficients for the intake of or exposure to radionuclides in the workplace and the environment.

The purpose of this document is to guide EPA personnel in calculating release criteria based on regulations promulgated under various methods of dose calculation. This guidance will relate these dose limits to a single measure, cleanup concentration. This guidance will assist RPMs in making decisions at these sites.

Note: use of this calculator to develop dose compliance concentrations for some dose-based ARARs does not affect the CERCLA requirement to comply with all other Federal and State ARARs at a site (e.g., 40 CFR 141.66, 40 CFR 192.12). ARARs are determined site-specifically. For a list of "Likely Federal Radiation Applicable or Relevant and Appropriate (ARARs)", see Attachment A of EPA's guidance "[Establishment of Cleanup Levels for CERCLA sites with Radioactive Contamination](#)." For additional guidance documents on compliance with ARARs at radioactively contaminated sites, go to the following webpage: <https://www.epa.gov/superfund/radiation-superfund-sites>

This website combines current EPA DCFs with "standard" exposure factors to estimate contaminant concentrations in environmental media (soil and water) that are protective of humans (including sensitive groups) over a lifetime. Exceeding a DCC usually suggests that further evaluation of the potential doses is appropriate. The DCC concentrations presented on this website can be used to screen pollutants in environmental media, trigger further investigation, and provide initial cleanup goals, if applicable. DCCs should be applied in accordance with guidance from EPA Regions.

In addition to this guidance, for relevant training, see the internet-based course "Radiation Risk Assessment: Updates and Tools. [https://clu-in.org/conf/itrc/rads\\_051507/](https://clu-in.org/conf/itrc/rads_051507/)

## 2. Understanding the DCC Website

### 2.1 General Considerations

DCCs are isotope concentrations that correspond to certain levels of dose in air, soil, water, and biota. Dose Coefficients (DCFs), for a given radionuclide, represent the dose equivalent per unit intake (i.e., ingestion or inhalation) or external exposure of that radionuclide. In dose assessments, these DCFs are used in calculations with radionuclide concentrations and exposure assumptions to estimate dose from exposure to radioactive contamination. The calculations may be rearranged to generate DCCs for a specified level of dose. DCFs may be specified for specific body organs or tissues of interest or as a weighted sum of individual organ dose, termed the effective dose equivalent. These DCFs may be multiplied by the total activity of each radionuclide inhaled or ingested per year or the external exposure concentration to which a receptor may be exposed to estimate the dose to the receptor. Dose Coefficients used are provided by the [Center for Radiation Protection Knowledge](#). The main report is [Calculations of Slope Factors and Dose Coefficients](#) and the tables of DCFs are in a separate [appendix](#).

Inhalation dose coefficients are tabulated separately for each of the three lung absorption types considered in the lung model currently recommended by the International Commission on Radiological Protection (ICRP) and, where appropriate, for inhalation of radionuclides in vapor or gaseous forms.

The designations "F", "M", and "S" presented in the Radionuclide Table under the heading "ICRP Lung Type" refer to the lung absorption type for inhaled particulate radionuclides, expressed as fast (F), medium (M), or slow (S), used in the current ICRP model of the respiratory tract. The inhalation dose coefficient values tabulated in the Radionuclide Table for each radionuclide have been selected based on the following guidelines: (1) For those elements where Table 4.1 of Federal Guidance Report No. 13 (and Table 2 of ICRP Publication 72) specifies a recommended default lung absorption type for particulates, the inhalation dose coefficient for that type is tabulated in the Radionuclide Table for each radioisotope of that element. (2) For those elements where no specific lung absorption type is recommended and multiple types are indicated as plausible choices, the inhalation dose coefficient reported in the Radionuclide Table for each radioisotope of that element is the maximum of the values for each of the plausible lung absorption types. (3) Where Federal Guidance Report No. 13 specifies risk coefficients for multiple chemical forms of certain elements (tritium, carbon, sulfur, iodine, and mercury), the inhalation dose coefficient value for the form estimated to pose the maximum dose is reported in the Radionuclide Table in most cases.

Inhaled particulates are assumed to have an activity median aerodynamic diameter (AMAD) of 1  $\mu$ m, as recommended by the ICRP for consideration of environmental exposures in the absence of specific information about physical characteristics of the aerosol. Where appropriate, radionuclides may be present in gas or vapor form, are designated by "G" and "V", respectively; such radionuclides include tritium, carbon, sulfur, nickel, ruthenium, iodine, tellurium, and mercury.

The most common land uses and exposure assumptions are included in the equations on this website: [Resident Soil](#), [Composite Worker Soil](#), [Outdoor Worker Soil](#), [Indoor Worker Soil](#), [Construction worker Soil](#), [Farmer Soil](#), [Recreator Soil](#), [resident Air](#), [Composite Worker Air](#), [Outdoor Worker Air](#), [Indoor Worker Air](#), [Construction worker Air](#), [Farmer Air](#), [Recreator Air](#), [Tapwater](#), [Soil to Groundwater](#) and [Ingestion of Fish](#).

The DCCs are generated with [standard exposure route equations](#) using EPA DCFs and exposure [parameters](#). For the calculation of oral dose coefficients, area correction factors, and gamma shielding factors, a standard soil density of 1.6 g/cm<sup>3</sup> has been used.

## 2.2 DCC Output Options

The calculator offers three options for calculating DCCs. Previous versions of this calculator employed slope factors that included progeny ingrowth for 100 years, designated "+D." The +D slope factors are no longer included in the pick list. This section describes the potential applications of the three choices and recommends a default DCC calculation.

### 2.2.1 DCC Output Option #1: Assumes Secular Equilibrium Throughout the Chain (no decay)

This is the preferred DCC calculation option and is marked as the default selection in the calculator. When a single isotope is selected, the calculator identifies all the daughters in the chain. The DCCs for each daughter are combined with the parent on a fractional basis. The fractional basis is determined by branching fractions, where a progeny may decay into more than one isotope. The resulting DCC is now based on secular equilibrium of the full chain. For straight chain decay, all the progeny would be at the same activity of the parent, and the DCC provided in the output would be the inverse sum of the reciprocal DCCs of the parent and all the progeny. Currently, all the soil DCC equation images are presented with a radioactive decay term to account for half-lives shorter than the exposure duration. Decay is not included in this DCC option, as the assumption of secular equilibrium is that the parent is continually being renewed.

When the secular equilibrium DCC output option is selected, the DCC Calculator now gives the option to show the individual progeny contributions for the DCC (and dose) output. When the option to display progeny contribution is selected, the DCC Calculator output gives the secular equilibrium DCC and the individual progeny DCCs in separate tables.

- A total DCC is calculated using the following formula.

Total secular equilibrium DCC for parent isotope;

$$DCC_{SE-tot} = \frac{1}{\sum_{i=1}^n \frac{1}{DCC_{SE-route} i}}$$

where:

n = total number of exposure routes;

Route secular equilibrium DCC for parent isotope:

$$DCC_{SE-route} = \frac{1}{\sum_{i=1}^n \left( \frac{DCC}{FC} \right)_i}$$

where:

n = total number of isotopes in decay chain;

FC = fractional contribution of isotope in decay chain;

DCC = DCC for isotope in decay chain without decay.

### 2.2.2 DCC Output Option #2: Does Not Assume Secular Equilibrium, Provides Results for Progeny Throughout Chain (with decay)

This option displays the DCCs calculated with half-life decay as identified in the DCC equation images. In addition to the selected isotope, all the individual progeny DCCs are displayed. Each DCC is determined with each isotope's respective half-life and not that of its parent isotope. This option does not assume secular equilibrium and presents all the individual progeny DCCs, so that the risk assessor can identify any isotopes that will be present and those that have no data. Users can alter progeny half-life to match the parent isotope or other progeny or to account for ingrowth and decay over a chain.

### 2.2.3 DCC Output Option #3: Does Not Assume Secular Equilibrium, No Progeny Included (with decay)

This option displays DCCs for only the selected isotopes with half-life decay as identified in the DCC equation images. In this output, secular equilibrium is not assumed, progeny DCCs are not displayed, and progeny contribution is not combined into the DCC for the selected isotope. This option is useful when contamination is from one radionuclide with a very long half-life, where secular equilibrium would be too conservative.

## 2.3 Dose Conversion Factors (DCFs)

Users should choose the DCFs [International Commission on Radiological Protection (ICRP) 30, 60 or 107] required by the ARAR. If DCFs are not specified within the regulation (for example, the Code of Federal Regulations for a federal standard that is being complied with as an ARAR), then users should generally choose ICRP 107 DCFs. This recommendation is consistent with the guidance contained in "Use of IRIS Values in Superfund Risk Assessment" (OSWER 9285.7-16) for EPA to evaluate dose based upon its best scientific judgment. For further discussion of the scientific differences between ICRP 30 and 60 methodologies, see ["Dosimetric Significance of the ICRP's Updated Guidance and Models, 1989-2003, and Implications for U.S. Federal Guidance"](#) (August 2003, ORNL/TM-2003/207). For a discussion of the impacts of the ICRP 107 nuclear decay data, see [Impact of the New Nuclear Decay Data of ICRP Publication 107 on Inhalation Dose Coefficients for Workers](#).

EPA classifies all radionuclides as Group A carcinogens ("carcinogenic to humans"). Group A classification is used only when there is sufficient evidence from epidemiologic studies to support a causal association between exposure to the agents and cancer. The [appendix radionuclide table](#), from the [Center for Radiation Protection Knowledge](#), lists ingestion, inhalation, and external exposure dose coefficients for radionuclides in conventional units of picocuries (pCi). Ingestion and inhalation dose coefficients are central estimates in a linear model

of the age-averaged, lifetime attributable radiation cancer incidence (fatal and nonfatal cancer) dose per unit of activity inhaled or ingested, expressed as *mrem/pCi*. External exposure dose coefficients are central estimates of lifetime attributable radiation dose for each year of exposure to external radiation from photon-emitting radionuclides distributed uniformly in a thick layer of soil and are expressed as *mrem/year per pCi/gram soil*. External exposure dose coefficients can also be used that have units of *mrem/year per pCi/cm<sup>2</sup> soil*. When combined with site-specific media concentration data and appropriate exposure assumptions, dose coefficients can be used to estimate annual dose to members of the general population due to radionuclide exposures. EPA currently provides guidance on inhalation risk assessment in [RAGS Part F](#) (Risk Assessment Guidance for Superfund Volume I: Human Health Evaluation Manual, Part F, Supplemental Guidance for Inhalation Risk Assessment). This guidance only addresses chemicals. The development of inhalation dose coefficients for radionuclides differs from the guidance presented in RAGS Part F for development of inhalation unit risk (IUR) values for chemicals.

The DCFs from the [Center for Radiation Protection Knowledge](#) differ from the values presented in [FGR 12 CD supplement](#). The DCFs were calculated using ORNL's DCAL software in the manner of Federal Guidance Report 12 and 13. For the calculation of oral dose coefficients, a standard soil density of 1.6 g/cm<sup>3</sup> has been used. The radionuclides presented are those provided in the International Commission on Radiological Protection (ICRP) [Publication 107](#). This document contains a revised database of nuclear decay data (energies and intensities of emitted radiations, physical half-lives, and decay modes) for 1,252 naturally occurring and man-made radionuclides. ICRP Publication 107 supersedes the previous database, ICRP Publication 38, published in 1983.

### 2.3.1 ICRP 30

Unlike ICRP 2, which did not calculate DCF per se, ICRP 30 does present DCFs that may be used to calculate either organ dose equivalent or effective dose equivalent for ingestion and inhalation. For each radionuclide, ICRP 30 provides values for the organ dose equivalent conversion factors, hT,50, and the effective dose equivalent conversion factor, hE,50 (calculated using the organ weighting factors w<sub>T</sub>). These values are also presented in [Federal Guidance Report No. 11](#). Organ DCFs are provided for those organs that have specific weighting factors, namely the gonads, breast, red marrow, lungs, thyroid, and bone surfaces. Organ DCFs are also given for the remainder, which include the five remaining tissues that receive the next highest doses. These include the liver, kidneys, spleen, brain, small intestine, upper large intestine, lower large intestine, etc.

Organ dose equivalent conversion factors and effective dose equivalent conversion factors for all radionuclides selected for this analysis are provided in Attachment A, Table A.2 (inhalation) and Table A.3 (ingestion). These values, in units of mrem/pCi, have been taken from Tables 2.1 and 2.2 respectively of [Federal Guidance Report No. 11](#).

### 2.3.2 ICRP 60

ICRP 60 also presents DCFs. Most of the world's radiation standards are based on this document. ICRP 60 is similar to ICRP 30, except it is based on more recent findings. ICRP 60 risk estimates increased due to cancers in Japanese populations exposed to radiation from World War II bombings. There were also reevaluations of the radiation dose calculations. These values are also presented in [Federal Guidance Report No. 13](#).

Isotopes that decay by spontaneous fission at greater than 0.1% (Cf-252, Cf-254, Cm-248, Cm-250 and Pu-244) are not in FGR-13. They are released in ICRP 72, which is analogous to the FGR 13 CD that contains most of the same values.

The use of dose conversion factors of ICRP 60/72 is mandated in the European Union by [European Council Directive 96/29](#) of May 13, 1996. If requested, NRC can grant a licensee an exemption to use the new dosimetric data of the ICRP (e.g. ICRP 68 for occupational exposures). In accordance with a June 8, 2007 [Federal Register](#) notice, DOE no longer requires a facility to get an exception to use ICRP 68 dosimetric data for occupational exposure. Non-regulatory studies (e.g., risk assessments) use the technically best available dose coefficients, which are those of the recent ICRP Publications. In addition, the IAEA in its Safety Series has adopted the ICRP Publication 60 Recommendations and the subsequent dose coefficients. For example, the dose coefficients of ICRP Publication 68 are contained in the IAEA Safety Guide entitled "Assessment of Occupational Exposure Due to Intakes of Radionuclides", RS-G-1.2, issued in 1999.

### 2.3.3 ICRP 107

ICRP Publication 107 (ICRP 2008) provides an electronic database of the physical data for calculations of radionuclide-specific protection and operational quantities. This database supersedes the data of ICRP 38 and will be used in future ICRP publications of dose coefficients for the intake of or exposure to radionuclides in the workplace and the environment.

The database contains information on the half-lives, decay chains, and yields and energies of radiations emitted in nuclear transformations of 1252 radionuclides of 97 elements. The CD accompanying the publication provides electronic access to complete tables of the emitted radiations as well as the beta and neutron spectra. The database has been constructed such that user-developed software can extract the data for further calculations of a radionuclide of interest. A Windows-based application is provided to display summary information on a user-specified radionuclide as well as the general characterization of the nuclides contained in the database. In addition, the application allows users to export the emissions of a specified radionuclide for use in subsequent calculations.

### 2.3.4 Federal Guidance Report 12

ICRP Publications 30 and 60 provide dose coefficients for the ingestion and inhalation intake of radionuclides. Dose coefficients for exposure to the radiations emitted by radionuclides present outside the body are given in Federal Guidance Report 12. That report addresses radionuclides uniformly distributed in air, in water, on the surface of the soil, and within the volume of the soil. The published report is consistent with ICRP Publication 26; however, the CD Supplement to Federal Guidance Report 13 provides values for the effective dose as defined in ICRP Publication 60.

### 2.3.5 Metastable Isotopes

Most dose and risk coefficients are presented for radionuclides in their ground state. In the decay process, the newly formed nucleus may be in an excited state and emit radiation (e.g., gamma rays) to lose the energy of the state. The excited nucleus is said to be in a metastable state, which is denoted by the chemical symbol and atomic number appended by "m" (e.g., Ba-137m). If additional higher energy metastable states are present, then "n", "p", ... is appended. Metastable states have different physical half-lives and emit different radiations and thus unique dose and risk coefficients. In decay data tabulations of [ICRP 107](#), if the half-life of a metastable state was less than 1 minute, then the radiations emitted in de-excitation are included with those of the parent radionuclide. Click to see a graphical representation of the decay of [Cs-137 to Ba-137](#).

Eu-152, in addition to its ground state, has two metastable states: Eu-152m and Eu-152n. The half-lives of Eu-152, Eu-152m and Eu-152n are: 13.5 y, 9.31 min and 96 min, respectively, and the energy emitted per decay is 1.30 MeV, 0.080 MeV, and 0.14 MeV,

respectively.

## 2.4 Radionuclide-Specific Parameters

Several radionuclide-specific parameters are needed for development of the DCCs. The parameters are selected from a hierarchy of sources.

### 2.4.1 Sources

Many sources are used to populate the database of radionuclide-specific parameters. They are briefly described below.

1. [IAEA TRS 472 \(IAEA\)](#). Handbook of Parameter Values for the Prediction of Radionuclide Transfer in Terrestrial and Freshwater Environments. Technical Reports Series No. 472. International Atomic Energy Agency, Vienna. 2010. (IAEA TRS 364. Handbook of Parameter Values for the Prediction of Radionuclide Transfer in Temperate Environments (Technical Reports Series No. 364), 1994 was used to supplement egg, poultry and swine transfer factors.) [Spreadsheet of values](#).
2. [NCRP 123 \(NCRP\)](#). NCRP Report No. 123, Screening Models for Releases of Radionuclides to the Atmosphere, Surface Water, and Ground. National Council on Radiation Protection and Measurements. January 22, 1996. [Spreadsheet of values](#).
3. [EPA Radionuclide Soil Screening Level \(SSL\)](#). Soil Screening Guidance for Radionuclides: User's Guide. Office of Solid Waste and Emergency Response (OSWER) Directive 9355.4-16A. October 2000. [Spreadsheet of values](#).
4. [RESRAD](#). User's Manual for RESRAD Version 6. Environmental Assessment Division, Argonne National Laboratory. ANL/EAD-4. July 2001. [Spreadsheet of values](#).
5. [BAES](#). A Review and Analysis of Parameters for Assessing Transport of Environmentally Released Radionuclides through Agriculture. C. F. Baes III, R. D. Sharp, A. L. Sjoreen, R.W. Shor. Oak Ridge National Laboratory 1984. [Spreadsheet of values](#).
6. [ICRP 2008](#). Nuclear Decay Data for Dosimetric Calculations. International Commission on Radiological Protection Publication 107. Ann. ICRP 38 (3), 2008.
7. [EPAKD](#). Understanding Variation in Partition Coefficient, K<sub>d</sub>, Values. Volume II: Review of Geochemistry and Available K<sub>d</sub> Values for Cadmium, Cesium, Chromium, Lead, Plutonium, Radon, Strontium, Thorium, Tritium (3H), and Uranium. Office of Air and Radiation. EPA 402-R-99-004B. August 1999. and Volume III: Review of Geochemistry and Available K<sub>d</sub> Values for Americium, Arsenic, Curium, Iodine, Neptunium, Radium, and Technetium. Office of Air and Radiation. EPA 402-R-04-002C. July 2004. [Spreadsheet of values](#).

### 2.4.2 Hierarchy by Parameter

Generally the hierarchies below are followed.

- Half-life (yr), Decay mode, Atomic weight, Atomic number and Decay energy. [ICRP 107](#).
- Milk transfer factor (TF<sub>dairy</sub>) (day/L). IAEA, EA, NCRP, RESRAD. TF<sub>dairy</sub> is the volumetric activity density in milk (pCi/L) divided by the daily intake of radionuclide (pCi/day).
- Beef transfer factor (TF<sub>beef</sub>) (day/kg). IAEA, EA, NCRP, RESRAD. TF<sub>beef</sub> is the mass activity density in beef (pCi/kg fresh weight) divided by the daily intake of radionuclide (in pCi/d).
- Fish bioconcentration factor (BCF) (L/kg). IAEA, EA, RESRAD. BCF is the ratio of the radionuclide concentration in the fish tissue (pCi/kg fresh weight) from all exposure pathways relative to that in water (pCi/L).
- Poultry transfer factor (TF<sub>poultry</sub>) (day/kg). IAEA. TF<sub>poultry</sub> is the mass activity density in poultry (pCi/kg fresh weight) divided by the daily intake of radionuclide (in pCi/d).
- Egg transfer factor (TF<sub>egg</sub>) (day/kg). IAEA. TF<sub>egg</sub> is the mass activity density in egg (pCi/kg fresh weight) divided by the daily intake of radionuclide (in pCi/d).
- Swine transfer factor (TF<sub>swine</sub>) (day/kg). IAEA. TF<sub>swine</sub> is the mass activity density in swine (pCi/kg fresh weight) divided by the daily intake of radionuclide (in pCi/d).
- Sheep Milk transfer factor (TF<sub>sheep-milk</sub>) (day/L). IAEA. TF<sub>sheep-milk</sub> is the volumetric activity density in sheep milk (pCi/L) divided by the daily intake of radionuclide (pCi/day).
- Sheep transfer factor (TF<sub>sheep</sub>) (day/kg). IAEA, EA. TF<sub>sheep</sub> is the mass activity density in sheep (pCi/kg fresh weight) divided by the daily intake of radionuclide (in pCi/d).
- Goat Milk transfer factor (TF<sub>goat-milk</sub>) (day/L). IAEA. TF<sub>goat-milk</sub> is the volumetric activity density in goat milk (pCi/L) divided by the daily intake of radionuclide (pCi/day).
- Goat transfer factor (TF<sub>goat</sub>) (day/kg). IAEA. TF<sub>goat</sub> is the mass activity density in goat (pCi/kg fresh weight) divided by the daily intake of radionuclide (in pCi/d).
- Soil to water partition coefficient (K<sub>d</sub>) (mg/kg-soil per mg/L water or simplified = L/kg). EPAKD, IAEA, SSL, RESRAD, BAES. K<sub>d</sub> is the ratio of the mass activity density (pCi/kg) of the specified solid phase (usually on a dry mass basis) to the volumetric activity density (Bq/L) of the specified liquid phase.
- Soil to plant transfer factor-wet (Bv<sub>wet</sub>) (pCi/g plant per pCi/g soil). IAEA, EA, NCRP, SSL, RESRAD, BAES. The values for cereal grain are used from IAEA. Bv<sub>wet</sub> is the ratio of the activity concentration of radionuclide in the plant (pCi/kg wet mass) to that in the soil (pCi/kg dry mass). Note: Some Bv<sub>wet</sub> values were derived from Bv<sub>dry</sub> sources, assuming the ratio of dry mass to fresh mass was presented in the source documents.

- For carbon, the only value in the hierarchy is found in RESRAD. This value is excluded, as it overestimates root uptake. See section 2.5.4 for a detailed discussion of the carbon transfer factor derivation.

- Soil to plant transfer factor-dry (Bv<sub>dry</sub>) (pCi/g plant per pCi/g soil). IAEA, EA, NCRP, SSL, RESRAD, BAES. The values for cereal grain are used. Bv<sub>dry</sub> is the ratio of the activity concentration of radionuclide in the plant (pCi/kg dry mass) to that in the soil (pCi/kg dry mass). Note: Some Bv<sub>dry</sub> values were derived from Bv<sub>wet</sub> sources, assuming the ratio of dry mass to fresh mass was presented in the source documents.

- For carbon, the only value in the hierarchy is found in RESRAD. This value is excluded, as it overestimates root uptake. See section 2.5.4 for a detailed discussion of the carbon transfer factor derivation.

Bv<sub>wet</sub> and Bv<sub>dry</sub> can be determined using the following equations.

$$Bv_{wet} = Bv_{dry} \times \left( \frac{100 - MC}{100} \right)$$

where:  
MC = percent moisture content (%)

and:

$$Bv_{dry} = Bv_{wet} \times \left( \frac{100}{100 - MC} \right)$$

where:  
MC = percent moisture content (%)

2.5 Biota Modeling

2.5.1 Produce Modeling

There are 22 individually calculated DCCs that make up the default produce DCC. Each individual DCC is determine based on produce specific data such as intake rate, soil to plant transfer factors ( $Bv_{wet}$ ), and soil mass loading factor (MLF). These 22 individual DCCs are then summed by inverse reciprocal to determine a total produce DCC. The MLFs and intake rates used in the determination of the default biota DCCs are based on fresh weight. Intake rates, MLFs, and tranfer factors were updated in July 2016. A pdf of the Technical Memorandum, released for this update, can be found [here](#).

2.5.1.1 Intake Rates (g/day)

Table 2.5.1-A provides all of the default produce intake rates that are used to determine the total produce DCC. The delineation of (FW) in the column header indicates that the intake rates are for fresh weight; these are the intake rates used when the tool is run in default mode. In site-specific mode, the user may choose between Fresh Weight (FW) or Cooked Weight (CPW), which takes cooking and preparation loss into account. In addition, the user may also add rice and cereal grain to the produce output. These intake rates can be found in Table 2.5.1-B below and are only given in dry weight (DW). In user-provided mode, the user may change produce specific and element specific parameters to model produce that is not provided in this tool, such as soil to plant transfer factor, mass loading factor, contaminated fraction, and intake rates. Another source for intake rates is the [Food Commodity Intake Database](#) (FCID). If the FCID is used, the user must convert the data to g/day, as it is required for use in this tool.

Table 2.5.1-A	Intake Rate for Farmer Child (g/day) (FW)	Intake Rate for Farmer Adult (g/day) (FW)	Intake Rate for Resident Child (g/day) (FW)	Intake Rate for Resident Adult (g/day) (FW)	Intake Rate for Farmer Child (g/day) (CPW)	Intake Rate for Farmer Adult (g/day) (CPW)	Intake Rate for Resident Child (g/day) (CPW)	Intake Rate for Resident Adult (g/day) (CPW)
Apples	82.9	84.7	72.2	73.7	43.0	43.9	37.4	38.2
Citrus	194.4	309.4	194.1	309.4	100.6	160.4	100.6	160.4
Berries other than Strawberries	23.9	35.4	23.9	35.4	12.4	18.3	12.4	18.3
Peaches	99.3	103.1	111.4	115.7	51.5	53.5	57.7	60.0
Pears	76.9	59.9	66.7	51.9	39.9	31.1	34.6	26.9
Strawberries	25.3	40.5	25.3	40.5	13.1	21.0	13.1	21.0
Asparagas	12.0	39.3	12.0	39.3	8.2	26.8	8.2	26.8
Beets	3.9	33.9	3.9	33.9	2.7	23.2	2.7	23.2
Broccoli	14.4	35.3	13.1	32.0	9.9	24.1	8.9	21.9
Cabbage	11.5	85.7	12.3	92.1	7.8	58.6	8.4	62.9
Carrots	13.3	24.4	14.9	27.3	9.1	16.6	10.2	18.7
Corn	32.7	82.0	23.8	59.8	22.3	56.0	16.3	40.9
Cucumbers	16.9	54.9	25.4	82.4	11.5	37.5	17.3	56.3
Lettuce	4.2	37.5	4.2	37.5	2.9	25.6	2.9	25.6
Lima Beans	6.5	33.8	6.5	33.8	4.5	23.1	4.5	23.1
Okra	5.3	30.2	5.3	30.2	3.6	20.7	3.6	20.7

Onions	7.2	27.2	5.8	21.8	4.9	18.6	4.0	14.9
Peas	28.7	31.7	32.1	35.4	19.6	21.7	21.9	24.2
Pumpkins	45.2	64.8	45.2	64.8	30.9	44.2	30.9	44.2
Snap Beans	27.5	54.2	27.3	53.9	18.8	37.0	18.7	36.8
Tomatoes	34.9	94.2	29.7	80.3	23.8	64.4	20.3	54.8
White Potatoes	57.3	141.8	51.7	127.8	39.1	96.9	35.3	87.3

Table 2.5.1-B	Intake Rate for Farmer Child (g/day) (DW)	Intake Rate for Farmer Adult (g/day) (DW)	Intake Rate for Resident Child (g/day) (DW)	Intake Rate for Resident Adult (g/day) (DW)
Rice	34.8	88.5	28.8	73.2
Cereal Grain	46.0	91.9	38.0	76.0

To determine which produce are commonly cultivated in the area around the site, users should contact their county extension office. The [National Pesticide Information Center](#) has an interactive map that allows users to choose their state and county and then connects them to their county extension office.


2.5.1.2 Soil to Plant Transfer Factors (B<sub>v<sub>wet</sub></sub>)

The new soil to plant transfer factors (B<sub>v<sub>wet</sub></sub>) from IAEA ([TRS-472](#)) are unique to climate zone, soil type, and produce type. There are three climate zones (Temperate, Tropical, and Subtropical), seven soil types (Default, Sand, Loam, Clay, Organic, Coral Sand, and Other), and 24 produce types implemented in the DCC calculator. When the tool is run in default mode, the climate zone is temperate, the soil type is Default, which applies to all soil types, and 22 produce are used. Corn and rice are not used in default mode because the parameters used for these are based on dry weight, whereas the other 22 produce are based on fresh weight. For rice, IAEA did not specify a particular climate zone; therefore, the rice transfer factors have been applied to all three climate zones.

Climate Zones

The following map shows how the climate zones are distributed across the United States.





\*[The Holdridge life zones of the conterminous United States in relation to ecosystem mapping](#). Journal of Biogeography, 26, 1032.

Soil Types

Table 2 below describes the soil classification used in [TRS-472](#). The Coral Sand and Other soil types are not listed in Table 2. The 'Other' soil type classification, in a temperate climate, was created for soils without characterization data and for mineral soils with unknown sand and clay contents (TRS-472 pg. 9). For tropical climates, the 'Coral Sand' soil type classification was changed from 'Other' (given in TRS-472, because it refers to soils that are outside the classification scheme used here, such as Marshall Island soils, classified by the authors as coral sand soil) (TRS-472, pg. 73).


TABLE 2. TYPICAL RANGES OF VALUES OF SELECTED SOIL PARAMETERS FOR THE FOUR SOIL GROUPS

Soil group	pH	Organic matter content (%)	Cation exchange capacity (cmol/kg)	Sand content in the mineral matter fraction (%)	Clay content in the mineral matter fraction (%)
Sand	3.5–6.5	0.5–3.0	3.0–15.0	≥65	<18
Loam	4.0–6.0	2.0–6.5	5.0–25.0	65–82	18–35
Clay	5.0–8.0	3.5–10.0	20.0–70.0	—	≥35
Organic	3.0–5.0	≥20	20.0–200.0	—	—

\*[Technical Report Series no. 472](#).

p>





\*Land Data Assimilation Systems (LDAS). [NASA](#)


**Produce Types**

The following table illustrates which soil to plant transfer factor categories from IAEA are used for each produce type in the DCC calculator. The individual produce output only lists the category name from IAEA; however, a value from a secondary source may be being utilized.

Produce Type	Primary Transfer Factor Category	Primary Transfer Factor Source	Secondary Transfer Factor Category	Secondary Transfer Factor Source	Tertiary Transfer Factor Category	Tertiary Transfer Factor Source
Apples	Woody Tree	IAEA	Fruit	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Citrus	Woody Tree	IAEA	Fruit	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Berries other than Strawberries	Shrub	IAEA	Fruit	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Peaches	Woody Tree	IAEA	Fruit	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Pears	Woody Tree	IAEA	Fruit	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Strawberries	Herbaceous	IAEA	Fruit	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Asparagus	Leafy Vegetable	IAEA	Green Vegetable	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Beets	Root	IAEA	Root Vegetable	EA	None	NCRP-123, RADSSL, RESRAD, and Baes

Broccoli	Non-Leafy Vegetable	IAEA	Green Vegetable	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Cabbage	Leafy Vegetable	IAEA	Green Vegetable	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Carrots	Root	IAEA	Root Vegetable	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Corn	Maize Grain	IAEA	Green Vegetable	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Cucumbers	Non-Leafy Vegetable	IAEA	Green Vegetable	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Lettuce	Leafy Vegetable	IAEA	Green Vegetable	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Lima Beans	Legume Seed	IAEA	Green Vegetable	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Okra	Non-Leafy Vegetable	IAEA	Green Vegetable	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Onions	Non-Leafy Vegetable	IAEA	Root Vegetable	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Peas	Legume Seed	IAEA	Green Vegetable	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Pumpkins	Non-Leafy Vegetable	IAEA	Green Vegetable	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Snap Beans	Legume Seed	IAEA	Green Vegetable	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Tomatoes	Non-Leafy Vegetable	IAEA	Green Vegetable	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
White Potatoes	Tuber	IAEA	Root Vegetable	EA	None	NCRP-123, RADSSL, RESRAD, and Baes
Rice	Rice	IAEA	None	NCRP-123	None	RADSSL, RESRAD, and Baes
Cereal Grain	Cereal Grain	IAEA	None	NCRP-123	None	RADSSL, RESRAD, and Baes

- While included in the initial hierarchy analysis, RESRAD and BAES sources do not contribute to the current output. They are retained in the user guide for informational purposes.



- Please click on flow chart to view a larger image. Carbon is missing from this chart. Please see section 2.5.4 of this user guide for information about how a soil to plant transfer factor was derived for Carbon in the DCC calculator.

## 2.5.2 Animal Product Modeling

### 2.5.2.1 Intake Rates (g/day)

In default mode, the (FW) intake rates from Table 2.5.2-A are used. Similar to produce, there is an option to select Cooked Weight in site-specific mode. The intake rates for poultry include chicken, turkey, and duck. In default mode, the parameters used for poultry and eggs are for chicken specifically (i.e.,  $Q_p$ , etc.). If eggs and poultry are selected in site specific mode, the user will have the option to switch between chicken, turkey, duck, and goose, which will change the previously mentioned respective parameters; however, the transfer factor used will still be meant for chicken or duck, per TRS-472. If the user has a specific transfer factor for turkey or goose, then user-provided mode should be used.


Table 2.5.2-A	Intake Rate for Farmer Child (g/day) (FW)	Intake Rate for Farmer Adult (g/day) (FW)	Intake Rate for Farmer Child (g/day) (CPW)	Intake Rate for Farmer Adult (g/day) (CPW)
Dairy - Cow	994.7	676.4	994.7	676.4
Beef	62.8	165.3	31.1	81.7
Swine	33.7	92.5	16.6	45.7
Poultry	46.9	107.4	23.2	53.1
Egg	31.7	59.6	31.7	59.6
Fish	57.4	831.8	35.2	509.9

Site-specific mode will also offer the user the option to add animal products from Table 2.4.2-B to the output. The tool has transfer factors for these products; however, the user will need to enter their own intake rate data, as the tool does not provide any. Again, the [Food Commodity Intake Database](#) (FCID) may be used to find intake rate data but the user must convert the data to g/day, as it is required for use in this tool.

Table 2.5.2-B	Intake Rate for Farmer Child (g/day) (FW)	Intake Rate for Farmer Adult (g/day) (FW)	Intake Rate for Farmer Child (g/day) (CPW)	Intake Rate for Farmer Adult (g/day) (CPW)
Sheep	n/a	n/a	n/a	n/a
Sheep Milk	n/a	n/a	n/a	n/a
Goat	n/a	n/a	n/a	n/a
Goat Milk	n/a	n/a	n/a	n/a

### 2.5.2.2 Animal Transfer Factors (TF)

- While included in the initial hierarchy analysis, RESRAD and BAES sources do not contribute to the current output. They are retained in the user guide for informational purposes.



- Please click on flow chart to view a larger image. Carbon is missing from this chart. Please see section 2.5.4 of this user guide for information about how a soil to plant transfer factor was derived for Carbon in the DCC calculator.

2.5.3 Mass Loading Factor

A mass loading factor (MLF) is the amount, or mass, of soil that adheres to the plant surface. The following table lists the MLFs used in this tool according to plant type. For more information on how these were derived, please see the Technical Memorandum: [Biota Modeling in EPA's Preliminary Remediation Goal and Dose Compliance Concentration Calculators for Use in EPA Superfund Risk Assessment: Explanation of Intake Rate Derivation, Transfer Factor Compilation, and Mass Loading Factor Sources](#). This file is engineered for 2 sided printing.

Table 2.5.3-A	Mass Loading Facotr (MLF)	Units	Reference
Apples	1.60E-04	g dry soil / g fresh plant	<a href="#">EA (2009)</a>
Citrus	1.57E-04	g dry soil / g fresh plant	<a href="#">EA (2009)</a>
Berries other than Strawberries	1.66E-04	g dry soil / g fresh plant	<a href="#">EA (2009)</a>
Peaches	1.50E-04	g dry soil / g fresh plant	<a href="#">EA (2009)</a>
Pears	1.60E-04	g dry soil / g fresh plant	<a href="#">EA (2009)</a>
Strawberries	8.00E-05	g dry soil / g fresh plant	<a href="#">EA (2009)</a>
Asparagus	7.90E-05	g dry soil / g fresh plant	<a href="#">EA (2009)</a>
Beets	1.38E-04	g dry soil / g fresh plant	<a href="#">EA (2009)</a>
Broccoli	1.01E-03	g dry soil / g fresh plant	Hinton (1992), <a href="#">SSG-Appendix G</a>
Cabbage	1.05E-04	g dry soil / g fresh plant	<a href="#">EA (2009)</a>
Carrots	9.70E-05	g dry soil / g fresh plant	<a href="#">EA (2009)</a>
Corn	1.45E-04	g dry soil / g fresh plant	Pinder & McLeod (1989), <a href="#">SSG-Appendix G</a>

Cucumbers	4.00E-05	g dry soil / g fresh plant	<a href="#">EA (2009)</a>
Lettuce	1.35E-02	g dry soil / g fresh plant	Hinton (1992), <a href="#">SSG-Appendix G</a>
Lima Beans	3.83E-03	g dry soil / g fresh plant	Hinton (1992), <a href="#">SSG-Appendix G</a>
Okra	8.00E-05	g dry soil / g fresh plant	<a href="#">EA (2009)</a>
Onions	9.70E-05	g dry soil / g fresh plant	<a href="#">EA (2009)</a>
Peas	1.78E-04	g dry soil / g fresh plant	<a href="#">EA (2009)</a>
Pumpkins	5.80E-05	g dry soil / g fresh plant	<a href="#">EA (2009)</a>
Snap Beans	5.00E-03	g dry soil / g fresh plant	Hinton (1992), <a href="#">SSG-Appendix G</a>
Tomatoes	1.59E-03	g dry soil / g fresh plant	Hinton (1992), <a href="#">SSG-Appendix G</a>
White Potatoes	2.10E-04	g dry soil / g fresh plant	<a href="#">EA (2009)</a>
Cereal Grains	2.50E-01	g dry soil / g dry plant	Hinton (1992)
Rice	2.50E-01	g dry soil / g dry plant	Hinton (1992)
Pasture	2.50E-01	g dry soil / g dry plant	Hinton (1992)

2.5.4 Soil to Plant Transfer Factor Derivation for Carbon

The value of 5.5 given in Table D.3 of the RESRAD User's Manual for carbon root uptake was derived from data in Ng et al. 1968. Table 4 of Ng et al. 1968 presents a carbon composition in typical agricultural soil of 2.00E+04 ppm and a carbon composition in terrestrial plants of 1.10E+05 ppm in Table 10A. 1.10E+05 divided by 2.00E+04 gives the value of 5.5 reported in RESRAD. This value assumes that all the carbon in the plant is taken up by the roots; however, this is not the case. Photosynthesis is the primary source of carbon in plants. Carbon may be present in gas form in soils and volatilize into the plant canopy, where it may be taken up by the plant in some fraction depending on atmospheric conditions. It is typically estimated that 2% of plant carbon comes from soil (either directly or by uptake from the sub-canopy atmosphere). The other 98% of plant carbon comes from the above-canopy atmosphere, which is assumed not to contain carbon from the contaminated site. Consider that a plant is about 90% water and of the 10% dry matter about 40% is carbon. That plant would comprise about 4% carbon on a fresh weight basis. A mineral soil is typically about 2% to 5% organic matter, which corresponds to 0.8% to 2% carbon on a dry mass basis. Thus, taking the ratio of carbon contents results in a transfer factor of 4%/(0.8% to 2%) = 5.0 to 2.0 g fresh plant/g dry soil. The next step is to apply the 2% fraction of plant carbon derived from soil. The resulting range of transfer factors is 0.1 to 0.04; (2%\*(5.0 to 2.0)). The value of 0.1 is chosen for the calculation of DCCs and is used for all the BV<sub>wet</sub> values. BV<sub>dry</sub> values are derived for each plant type based on individual moisture content. For comparison purposes, the 5.5 value from RESRAD gives a transfer factor of 0.11 if the 2% assumption is made.

The above derivation assumes that all the carbon taken up by the plant is radioactive. In situations where radioactive carbon is mixing with stable carbon, a site-specific transfer factor can be derived using a model called "specific activity". Essentially, specific activity is the concentration ratio of the radioactive form to the stable form of carbon. Specific activity assumes that, within a compartment (i.e., soil), the radioactive contaminant mixes with the stable form both chemically and physically. Plants uptake the element in the same ratio as it exists in the soil compartment, resulting in the same ratio in the plant as in the soil compartment.

To determine a site-specific soil to plant transfer factor, actual site data must be available. Further, the flux rate of the element must be in a steady-state condition. The environmental compartments must be well defined and the fluxes between compartments well understood. For further information, refer to the following: [AMEC/004041/007](#) section 5, [ANL/EAD-4](#) Appendix L, and [IAEA TECDOC 1616](#) page 550.

2.6 DCCs in Context of Superfund Modeling Framework

This DCC calculator focuses on the application of generic and simple site-specific approaches that are part of a larger framework for calculating concentration levels to comply with dose-based ARARs. Generic DCCs for a 1 mrem standard are provided by viewing either the tables in the [Download Area](#) section of this tool or by running the [calculator](#) with "Defaults" selected for Site Info Type. [Part 3](#) of the [Soil Screening Guidance for Radionuclides: Technical Background Document](#) provides more information about 5 more detailed soil to groundwater models that are part of the same framework.

Generic DCCs are calculated from the same equations presented in the site-specific portion of the calculator, but they are based on a number of default assumptions chosen to be protective of human health for most site conditions. Generic DCCs, which should be scaled to



the same dose level as the standard being complied (e.g., multiplied by a factor of ten for a 10 mrem/year standard), can be used in place of site-specific DCC levels; however, in general, they are expected to be more protective than site-specific levels. The site manager should weigh the cost of collecting the data necessary to develop site-specific DCCs with the potential for deriving a higher DCC that provides an appropriate level of protection.

The framework presented in [Part 3](#) of the [Soil Screening Guidance for Radionuclides: Technical Background Document](#) includes more detailed modeling approaches that take into account more complex site conditions than the generic or simple site-specific methodology used in the Soil to Groundwater scenario in this calculator. More detailed approaches may be appropriate when site conditions (e.g., very deep water table, very thick uncontaminated unsaturated zone, soils underlain by karst or fractured rock aquifers) are different than those assumed in the generic or simple-site methodology. Further information on using more detailed approaches may be found in "[Simulating Radionuclide Fate and Transport in the Unsaturated Zone: Evaluation and Sensitivity Analyses of Select Computer Models](#)". This report provides a detailed technical analysis of five unsaturated zone fate and transport models for radionuclides.

To avoid unnecessary inconsistency between radiological and chemical risk assessment and radiological dose assessment at the same site, users should generally use the same model for chemical and radionuclide risk assessment and radionuclide dose assessment. If there is a reason for using another model, then justification for doing so should be developed. The justification should include specific supporting data and information in the administrative record. The justification would normally include the model runs, using both the recommended EPA DCC model and the alternative model. Users are cautioned that they should have a thorough understanding of both the DCC recommended model and any alternative model when evaluating whether a different approach is appropriate. When alternative models are used, the user should adjust the default input parameters to be as close as possible to the DCC inputs, which may be difficult, since models tend to use different definitions for parameters. Numerous computerized mathematical models have been developed by EPA and other organizations to predict the fate and transport of radionuclides in the environment; these models include single-media unsaturated zone models (for example, groundwater transport) as well as multi-media models. These models have been designed for a variety of goals, objectives, and applications; as such, no single model may be appropriate for all site-specific conditions. Generally, even when a different model is used to predict fate and transport of radionuclides through different media, EPA recommends using the DCC calculators for the remedial program to establish the dose-based concentrations to ensure consistency with CERCLA, the NCP, and EPA's Superfund guidance for remedial sites. Prior to using another model for dose assessment at a CERCLA remedial site, EPA regional staff should consult with the Superfund remedial program's National Radiation Expert (Stuart Walker, at (703) 603-8748 or [walker.stuart@epa.gov](mailto:walker.stuart@epa.gov)). For more information on this issue, please see questions 10 and 16 on pages 12, 17, and 18 of [Radiation Risk Assessment At CERCLA Sites: Q & A](#) (EPA 540-R-012-13, May 2014).

## 2.7 Understanding Dose Output on the DCC Website

The DCC [calculator](#) provides an option to select dose output. Selecting dose output requires the calculator to be run in "Site Specific" mode. The "Soil to Groundwater" medium does not have dose output and will become disabled when dose output is selected. The dose values presented on this site are radionuclide-specific values for individual contaminants in air, water, soil, and biota that may warrant further investigation or site cleanup.

### 2.7.1 General Considerations for the Dose Output

The first step in the dose assessment process is hazard identification where site data is screened against DCCs to identify radionuclides of potential concern (ROPCs). The "Dose Characterization" step, in the dose assessment process, incorporates the outcome of the exposure and toxicity assessment steps to calculate the annual dose identified in the data screening part of the hazard identification step. Dose is calculated for each land use determined appropriate for the site. The annual dose sum is presented for each exposure route for all ROPCs and for each ROPC across all exposure routes. The process used to calculate dose in this calculator follows the traditional method of first calculating a CDI (Chronic Daily Intake).


The basic equation for calculating total Dose is:

$$\begin{aligned} \text{Dose}_{\text{media-tot}} &= \sum_{i,k} \left( \text{Dose}_{i,k} \right) \\ \text{where:} \\ i &= \text{th route;} \\ k &= k^{\text{th}} \text{ isotope;} \\ \left( \text{Dose}_{\text{route-tot}} \right)_k &= \sum_i \left( \text{Dose}_{i,k} \right) \\ \text{and:} \\ \left( \text{Dose}_{\text{ROPC-tot}} \right)_i &= \sum_k \left( \text{Dose}_{i,k} \right) \\ \text{where:} \\ \text{Dose}_{i,k} &= \left( \text{CDI}_{\text{route}} \times \text{SF}_{\text{route}} \right) \\ \text{where:} \\ \text{CDI} &= \text{Chronic Daily Intake expressed in } \left[ \frac{\text{pCi}}{\text{year}}, \frac{\text{pCi}}{\text{g}}, \frac{\text{pCi}}{\text{cm}^2}, \frac{\text{pCi}}{\text{m}^3}, \frac{\text{pCi}}{\text{L}} \right] \\ \text{DCF} &= \text{Dose Coefficient expressed in } \left[ \frac{\text{mrem}}{\text{pCi}}, \frac{\text{mrem}}{\text{pCi/g}}, \frac{\text{mrem}}{\text{pCi/cm}^2}, \frac{\text{mrem}}{\text{pCi/m}^3}, \frac{\text{mrem}}{\text{pCi/L}} \right] \end{aligned}$$

The secular equilibrium equation for calculating total Dose is:

As presented in the previous section, the dose output is dependent on the DCC calculated. Section 2.2 discusses the DCC output options. To summarize section 2.2, the DCC options are either secular equilibrium or not. If the data is collected from a site where secular equilibrium is assumed to be present, the user need only enter the activity of the parent in the calculator, and a representative dose of the parent and all progeny will be presented in the calculator output. In the case of non-secular equilibrium, the current "state of the chain" may not be known or easily calculated. For relatively fast decaying isotopes, significant decay or ingrowth of progeny may have occurred since the sample date. Further, determining future activity of the contaminants may be useful in planning for future release of a property.

A [Decay Chain Activity Projection Tool](#) has been developed where the user can select an isotope, enter a length of time to allow decay and ingrowth, and enter the beginning activity of the parent. The results of this tool, pictured below, are the activities of the parent and progeny at the end of the decay and ingrowth of progeny time. These activities can be entered into the DCC calculator to calculate dose using the second and third DCC Output options.




The tables in the [DCC Download Area](#) provide generic concentrations in the absence of site-specific dose assessments. Screening concentrations can be used for:

- Prioritizing multiple sites within a facility or exposure units,
- Setting dose-based detection limits for contaminants of potential concern (COPCs), and
- Focusing future dose assessment efforts.

When using DCCs, the exposure pathways of concern and site conditions must match those taken into account by the screening levels. Thus, it is necessary to develop a conceptual site model (CSM) to identify likely contaminant source areas, exposure pathways, and potential receptors. This information can be used to determine the applicability of screening levels at the site and the need for additional information. The final CSM diagram represents linkages among contaminant sources, release mechanisms, exposure pathways, and routes and receptors based on historical information. It summarizes the understanding of the contamination problem. A separate CSM for ecological receptors can be useful. Part 2 and Attachment A of the Soil Screening Guidance for Radionuclides: Users Guide (EPA 2000a) contain the steps for developing a CSM.

A conceptual site model for the land uses presented in this calculator is presented below.



As a final check, the CSM should answer the following questions:

- Are there potential ecological concerns?
- Is there potential for land use other than those covered by the DCC levels (i.e., resident and worker)?
- Are there other likely human exposure pathways that were not considered in development of the DCC levels?
- Are there unusual site conditions (e.g., large areas of contamination, high fugitive dust levels, potential for indoor air contamination)?

The DCCs may need to be adjusted to reflect the answers to these questions.

### 3.2 Background Radiation

Natural background radiation should be considered prior to applying DCCs as cleanup levels. Background and site-related levels of radiation will be addressed as they are for other contaminants at CERCLA sites; for further information, see EPA's guidance "[Role of Background in the CERCLA Cleanup Program](#)", April 2002, (OSWER 9285.6-07P). Note that certain ARARs specifically address how to factor background into cleanup levels. For example, some radiation ARAR levels are established as increments above background concentrations. In these circumstances, background should be addressed in the manner prescribed by the ARAR.

### 3.3 Potential Problems and Limitations

As with any dose-based tool, the potential exists for misapplication. In most cases, this results from not understanding the intended use of the DCCs. In order to prevent misuse of the DCCs, the following should be avoided:

- Applying DCC levels to a site without adequately developing a conceptual site model that identifies relevant exposure pathways and exposure scenarios,
- Use of DCC levels as cleanup levels without the consideration of other relevant criteria such as ARARs,
- Using DCC levels as cleanup levels without verifying numbers with a health physicist/risk assessor,
- Using outdated DCC tables that have been superseded by more recent publications,
- Not considering the effects from the presence of multiple isotopes, and
- Not considering the individual model limitations as described in section 4 (e.g., inhalation of tapwater only considers C-14, H-3, Ra-224, Ra-226, Rn-220, and Rn-222).

## 4. Land Use Descriptions, Equations, and Technical Documentation

The DCCs consider human exposure from contact with contaminated soils and water. The equations and technical discussion are aimed at developing compliance levels for dose-based ARARs. The following text presents the land use equations and their exposure routes. [Table 1](#) (at the end of the User's Guide) presents the definitions of the variables and their default values. The default values and exposure models are consistent with the Regional Screening Levels for Chemical Contaminants at Superfund Sites ([RSL](#)) calculator where the same pathways are addressed (e.g., ingestion and inhalation) and are analogous when pathways are similar (e.g., dermal and external exposure). This calculator and the RSL both follow the recommendations in the [OSWER Directive](#) concerning use of exposure parameters from the [2011 Exposure Factors Handbook](#). Any alternative values or assumptions used in remedy evaluation or selection at a CERCLA site should be presented with supporting rationale in Administrative Records.

The DCC equations have evolved over time and are a combination of the following guidance documents:

- [Risk Assessment Guidance for Superfund: Volume I, Human Health Evaluation Manual \(Part B, Development of Risk-based Preliminary Remediation Goals\)](#) (RAGS Part B).
- [U.S. EPA, 2005. Human Health Risk Assessment Protocol for Hazardous Waste Combustion Facilities. Washington, DC. OSWER No. 5305W, EPA530-R-05-006](#)
- [U.S. EPA, 2000a. Soil Screening Guidance for Radionuclides: User's Guide. Office of Emergency and Remedial Response and Office of Radiation and Indoor Air. Washington, DC. OSWER No. 9355.4-16A EPA/540-R-00-007](#)
- [U.S. EPA, 2000b. Soil Screening Guidance for Radionuclides: Technical Background Document. Office of Emergency and Remedial Response and Office of Radiation and Indoor Air. Washington, DC. OSWER No. 9355.4-16](#)
- [U.S. EPA 2002. Supplemental Guidance for Developing Soil Screening Levels for Superfund Sites. OSWER 9355.4-24. December 2002.](#)
- U.S. EPA 1994b. Radiation Site Cleanup Regulations: Technical Support Documents for the Development of Radiation Cleanup Levels for Soil - Review Draft. Office of Radiation and Indoor Air, Washington, DC. EPA 402-R-96-011A. [PDF document](#) View Appendix C [here](#).
- [U.S. EPA, 1994c. Revised Draft Guidance for Performing Screening Level Risk Analyses at Combustion Facilities Burning Hazardous Wastes. Attachment C. Office of Emergency and Remedial Response. Office of Solid Waste. December 14.](#)
- [U.S. EPA, 1996a. Soil Screening Guidance: User's Guide. Office of Emergency and Remedial Response. Washington, DC. OSWER No. 9355.4-23](#)
- [U.S. EPA, 1996b. Soil Screening Guidance: Technical Background Document. Office of Emergency and Remedial Response. Washington, DC. OSWER No. 9355.4-17A](#)
- U.S. EPA, 1998. Human Health Risk Assessment Protocol for Hazardous Waste Combustion Facilities. Office of Solid Waste, Washington, DC. EPA530-D-98-001A. A secure PDF can be downloaded [here](#).
- [NCRP 1996. Screening Models for Releases of Radionuclides to Atmosphere, Surface Water, and Ground, Vols. 1 and 2. NCRP Report No. 123. National Council on Radiation Protection and Measurements.](#)


Note that if a route of exposure (e.g., ingesting fish from the pond in the farmer soil exposure scenario) is considered to be unreasonable at a site, both currently and in the future, eliminate the route in the site-specific option by entering zero for the ingestion rate of that route (e.g., replacing default fish ingestion rates in farmer soil scenario of 156.6 and 32.8 g/day with 0.0).

## 4.1 Resident

### 4.1.1 Resident Soil

This receptor spends most, if not all, of the day at home except for the hours spent at work. The activities for this receptor involve typical home making chores (cooking, cleaning, and laundering) as well as gardening. The resident is assumed to be exposed to contaminants via the following pathways: incidental ingestion of soil, external radiation from contaminants in soil, inhalation of fugitive dust, and consumption of home grown produce (100% of fruit and vegetables). Adults and children exhibit different ingestion rates for soil and produce. For example, the child resident is assumed to ingest 200 mg per day while the adult ingests 100 mg per day. Therefore, age-adjusted intake equations were developed to account for changes in intake as the receptor ages.

#### Graphical Representation



#### DCC Equations

The resident soil land use equation, presented here, contains the following exposure routes:

- incidental ingestion of soil

$$DCC_{res-soil-ing} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_o \left( \frac{mrem}{pCi} \right) \times IFS_{res-adj} \left( \frac{43,050 \text{ mg}}{year} \right) \times \left( \frac{g}{1000 \text{ mg}} \right)}$$

where:

$$IFS_{res-adj} \left( \frac{43,050 \text{ mg}}{year} \right) = \left( \left( EF_{res-c} \left( \frac{350 \text{ days}}{year} \right) \times IRS_{res-c} \left( \frac{200 \text{ mg}}{day} \right) \times AAF_{res-c} (0.23) \right) + \left( EF_{res-a} \left( \frac{350 \text{ days}}{year} \right) \times IRS_{res-a} \left( \frac{100 \text{ mg}}{day} \right) \times AAF_{res-a} (0.77) \right) \right)$$

where:

$$AAF_{res-c} (0.23) = \left( \frac{ED_{res-c} (6 \text{ years})}{ED_{res} (26 \text{ years})} \right) \text{ and: } AAF_{res-a} (0.77) = \left( \frac{ED_{res-a} (20 \text{ years})}{ED_{res} (26 \text{ years})} \right)$$

- inhalation of particulates emitted from soil

$$DCC_{res-soil-inh} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_i \left( \frac{mrem}{pCi} \right) \times IFA_{res-adj} \left( \frac{6,195 \text{ m}^3}{year} \right) \times \frac{1}{PEF \left( \frac{\text{m}^3}{kg} \right)} \times \left( \frac{1000 \text{ g}}{kg} \right)}$$

where:

$$IFA_{res-adj} \left( \frac{6,195 \text{ m}^3}{year} \right) = \left( \left( EF_{res-c} \left( \frac{350 \text{ days}}{year} \right) \times ET_{res-c} \left( \frac{24 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{res-c} \left( \frac{10 \text{ m}^3}{day} \right) \times AAF_{res-c} (0.23) \right) + \left( EF_{res-a} \left( \frac{350 \text{ days}}{year} \right) \times ET_{res-a} \left( \frac{24 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{res-a} \left( \frac{20 \text{ m}^3}{day} \right) \times AAF_{res-a} (0.77) \right) \right)$$

where:

$$AAF_{res-c} (0.23) = \left( \frac{ED_{res-c} (6 \text{ years})}{ED_{res} (26 \text{ years})} \right) \text{ and: } AAF_{res-a} (0.77) = \left( \frac{ED_{res-a} (20 \text{ years})}{ED_{res} (26 \text{ years})} \right)$$

- external exposure to ionizing radiation

$$DCC_{res-soil-ext} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-sv} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{res} \left( \frac{350 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ACF_{ext-sv} \times \left[ \left( ET_{res-o} \left( \frac{1,752 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-sv} (1.0) \right) + \left( ET_{res-i} \left( \frac{16,416 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{i-total} \right) \right]}$$

- consumption of produce (fruits and vegetables) - back-calculated to soil. Sections 9 and 13 of the 2011 Exposure Factors Handbook were used to derive the intakes for home-grown produce.

$$DCC_{res-soil-produce-ing-tot} (pCi/g) = \frac{1}{\sum_{i=1}^n DCC_{res-soil-produce-ing} (pCi/g)_i}$$

where:  
n = total number of produce items included

and:

$$DCC_{res-soil-produce-ing} (pCi/g) = \frac{DCC_{res-produce-ing} (pCi/g)}{(R_{upv} + R_{es})}$$

where:

$$R_{upv} = Bv_{wet} \left( \frac{pCi / g-fresh \text{ plant}}{pCi / g-dry \text{ soil}} \right); R_{es} = MLF_{produce} \left( \frac{g-dry \text{ soil}}{g-fresh \text{ plant}} \right)$$

where:

$$DCC_{res-produce-ing} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_o \left( \frac{mrem}{pCi} \right) \times IF_{res-adj} \left( \frac{g}{year} \right) \times CF_{res-produce} (1)}$$

where:

$$IF_{res-adj} \left( \frac{g}{year} \right) = \left( \left( EF_{res-c} \left( \frac{350 \text{ days}}{year} \right) \times IR_{res-c} \left( \frac{g}{day} \right) \times AAF_{res-c} (0.23) \right) + \left( EF_{res-a} \left( \frac{350 \text{ days}}{year} \right) \times IR_{res-a} \left( \frac{g}{day} \right) \times AAF_{res-a} (0.77) \right) \right)$$

and:

$$AAF_{res-c} (0.23) = \left( \frac{ED_{res-c} (6 \text{ years})}{ED_{res} (26 \text{ years})} \right) \text{ and: } AAF_{res-a} (0.77) = \left( \frac{ED_{res-a} (20 \text{ years})}{ED_{res} (26 \text{ years})} \right)$$

The consumption of produce exposure route drives the DCCs lower than all the other routes. It is recommended that produce-specific transfer factors ( $Bv_{wet}$ ) be used when available for a site. Further, the default transfer factors ( $Bv_{wet}$ ) from IAEA, used in these DCC calculations, are based on a composite of all soil groups. Transfer factors ( $Bv_{wet}$ ) for sand, loam, clay, organic, coral sand, and other soil types that may be more suited to a particular site are also provided. The site-specific option of the calculator can be used to focus on ingestion of individual produce types. When "Site-specific" is selected, if the user changes the "Select Isotope Info Type" to "User-provided", then a specific transfer factor may be changed.

- total



$$DCC_{res-soil-tot} (pCi/g) = \frac{1}{\frac{1}{DCC_{res-soil-ing}} + \frac{1}{DCC_{res-soil-inh}} + \frac{1}{DCC_{res-soil-ext}} + \frac{1}{DCC_{res-soil-produce-ing-tot}}}$$

A number of studies have shown that inadvertent ingestion of soil is common among children 6 years old and younger (Calabrese et al. 1989, Davis et al. 1990, Van Wijnen et al. 1990). Therefore, the dose method uses an age-adjusted soil ingestion factor that takes into account the difference in daily soil ingestion rates, body weights, and exposure duration for children from 1 to 6 years old and others from 7 to 26 years old. The equation is presented below. This health-protective approach is chosen to take into account the higher daily rates of soil ingestion in children as well as the longer duration of exposure that is anticipated for a long-term resident. For more on this method, see [RAGS Part B](#).

Age-adjusted intake factors are also used for inhalation of particulates emitted from soil and consumption of produce (presented above).

Definitions of the input variables are in [Table 1](#).

#### CDI Equations

- incidental ingestion of soil

$$CDI_{res-soil-ing} \left( \frac{pCi}{year} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times IFS_{res-adj} \left( \frac{43,050 \text{ mg}}{year} \right) \times \left( \frac{g}{1000 \text{ mg}} \right)$$

where:

$$IFS_{res-adj} \left( \frac{43,050 \text{ mg}}{year} \right) = \left( \left( EF_{res-c} \left( \frac{350 \text{ days}}{year} \right) \times IRS_{res-c} \left( \frac{200 \text{ mg}}{day} \right) \times AAF_{res-c}(0.23) \right) + \left( EF_{res-a} \left( \frac{350 \text{ days}}{year} \right) \times IRS_{res-a} \left( \frac{100 \text{ mg}}{day} \right) \times AAF_{res-a}(0.77) \right) \right)$$

where:

$$AAF_{res-c}(0.23) = \left( \frac{ED_{res-c}(6 \text{ years})}{ED_{res}(26 \text{ years})} \right) \text{ and: } AAF_{res-a}(0.77) = \left( \frac{ED_{res-a}(20 \text{ years})}{ED_{res}(26 \text{ years})} \right)$$

- inhalation of particulates emitted from soil

$$CDI_{res-soil-inh} \left( \frac{pCi}{year} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times IFA_{res-adj} \left( \frac{6,195 \text{ m}^3}{year} \right) \times \frac{1}{PEF \left( \frac{\text{m}^3}{kg} \right)} \times \left( \frac{1000 \text{ g}}{kg} \right)$$

where:

$$IFA_{res-adj} \left( \frac{6,195 \text{ m}^3}{year} \right) = \left( \left( EF_{res-c} \left( \frac{350 \text{ days}}{year} \right) \times ET_{res-c} \left( \frac{24 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{res-c} \left( \frac{10 \text{ m}^3}{day} \right) \times AAF_{res-c}(0.23) \right) + \left( EF_{res-a} \left( \frac{350 \text{ days}}{year} \right) \times ET_{res-a} \left( \frac{24 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{res-a} \left( \frac{20 \text{ m}^3}{day} \right) \times AAF_{res-a}(0.77) \right) \right)$$

where:

$$AAF_{res-c}(0.23) = \left( \frac{ED_{res-c}(6 \text{ years})}{ED_{res}(26 \text{ years})} \right) \text{ and: } AAF_{res-a}(0.77) = \left( \frac{ED_{res-a}(20 \text{ years})}{ED_{res}(26 \text{ years})} \right)$$

- external exposure to ionizing radiation

$$CDI_{res-soil-ext} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{res} \left( \frac{350 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ACF_{ext-sv} \times \left[ \left( \frac{ET_{res-o} \left( \frac{1,752 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-sv}(1.0) \right) + \left( \frac{ET_{res-i} \left( \frac{16,416 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{i-total} \right) \right]$$

- consumption of fruits and vegetables back-calculated to soil

$$CDI_{res-soil-produce-ing-tot} \left( \frac{pCi}{year} \right) = \sum_{i=1}^n CDI_{res-soil-produce-ing} \left( \frac{pCi}{year} \right)_i$$

where:

n = total number of produce items included

and:

$$CDI_{res-produce-ing} \left( \frac{pCi}{year} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times (R_{upv} + R_{es}) \times IF_{res-adj} \left( \frac{g}{year} \right) \times CF_{res-produce} \quad (1)$$

where:

$$R_{upv} = Bv_{wet} \left( \frac{pCi / g-fresh \text{ plant}}{pCi / g-dry \text{ soil}} \right); R_{es} = MLF_{produce} \left( \frac{g-dry \text{ soil}}{g-fresh \text{ plant}} \right)$$

and:

$$IF_{res-adj} \left( \frac{g}{year} \right) = \left( \left( EF_{res-c} \left( \frac{350 \text{ days}}{year} \right) \times IR_{res-c} \left( \frac{g}{day} \right) \times AAF_{res-c}(0.23) \right) + \left( EF_{res-a} \left( \frac{350 \text{ days}}{year} \right) \times R_{res-a} \left( \frac{g}{day} \right) \times AAF_{res-a}(0.77) \right) \right)$$

where:


$$AAF_{res-c}(0.23) = \left( \frac{ED_{res-c}(6 \text{ years})}{ED_{res}(26 \text{ years})} \right) \text{ and: } AAF_{res-a}(0.77) = \left( \frac{ED_{res-a}(20 \text{ years})}{ED_{res}(26 \text{ years})} \right)$$

#### 4.1.2 Resident Soil 2-D External Exposure

This receptor spends most, if not all, of the day at home except for the hours spent at work. The activities for this receptor involve typical home making chores (cooking, cleaning, and laundering) as well as gardening.

This analysis is designed to look at external exposure from contamination of different area sizes. Areas considered are 1 to 1,000,000 square meters. Isotope-specific area correction factors (ACF) were developed for this analysis.

### Graphical Representation



### DCC Equations

- Direct External Exposure to contamination at infinite depth

$$DCC_{res-soil-sv} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-sv} \left( \frac{mrem/yr}{pCi/g} \right) \times EF_{res} \left( \frac{350 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ACF_{ext-sv} \times \left[ \left( ET_{res-o} \left( \frac{1.752 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-sv} (1.0) \right) + \left( ET_{res-i} \left( \frac{16.416 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hour}} \right) \times GSF_{i-total} \right) \right]}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the soil volume for external exposure.

- Direct External Exposure to contamination 1 cm thick

$$DCC_{res-soil-1cm} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-1cm} \left( \frac{mrem/yr}{pCi/g} \right) \times EF_{res} \left( \frac{350 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ACF_{ext-1cm} \times \left[ \left( ET_{res-o} \left( \frac{1.752 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-1cm} (1.0) \right) + \left( ET_{res-i} \left( \frac{16.416 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hour}} \right) \times GSF_{i-total} \right) \right]}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 1cm soil volume for external exposure.

- Direct External Exposure to contamination 5 cm thick

$$DCC_{res-soil-5cm} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-5cm} \left( \frac{mrem/yr}{pCi/g} \right) \times EF_{res} \left( \frac{350 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ACF_{ext-5cm} \times \left[ \left( ET_{res-o} \left( \frac{1.752 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-5cm} (1.0) \right) + \left( ET_{res-i} \left( \frac{16.416 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hour}} \right) \times GSF_{i-total} \right) \right]}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 5cm soil volume for external exposure.

- Direct External Exposure to contamination 15 cm thick

$$DCC_{res-soil-15cm} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-15cm} \left( \frac{mrem/yr}{pCi/g} \right) \times EF_{res} \left( \frac{350 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ACF_{ext-15cm} \times \left[ \left( ET_{res-o} \left( \frac{1.752 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-15cm} (1.0) \right) + \left( ET_{res-i} \left( \frac{16.416 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hour}} \right) \times GSF_{i-total} \right) \right]}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 15cm soil volume for external exposure.

- Direct External Exposure to surface contamination

$$DCC_{\text{res-soil-gp}} \left( \frac{\text{pCi}}{\text{cm}^2} \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_{\text{ext-gp}} \left( \frac{\text{mrem/yr}}{\text{pCi/cm}^2} \right) \times EF_{\text{res}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ACF_{\text{ext-gp}} \times \left[ \left( ET_{\text{res-o}} \left( \frac{1,752 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{ext-gp}} (1.0) \right) + \left( ET_{\text{res-i}} \left( \frac{16,416 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{i-total}} \right) \right]}$$

The resulting units for this recommended DCC are in pCi/cm<sup>2</sup>. The units are based on area, because the DCF used is the ground plane for external exposure.

Definitions of the input variables are in [Table 1](#).

**CDI Equations**

- Direct External Exposure to contamination at infinite depth

$$CDI_{\text{res-soil-sv}} \left( \frac{\text{pCi}}{\text{g}} \right) = C_{\text{soil}} \left( \frac{\text{pCi}}{\text{g}} \right) \times EF_{\text{res}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ACF_{\text{ext-sv}} \times \left[ \left( ET_{\text{res-o}} \left( \frac{1,752 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{ext-sv}} (1.0) \right) + \left( ET_{\text{res-i}} \left( \frac{16,416 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{i-total}} \right) \right]$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the soil volume for external exposure.

- Direct External Exposure to contamination 1 cm thick

$$CDI_{\text{res-soil-1cm}} \left( \frac{\text{pCi}}{\text{g}} \right) = C_{\text{soil}} \left( \frac{\text{pCi}}{\text{g}} \right) \times EF_{\text{res}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ACF_{\text{ext-1cm}} \times \left[ \left( ET_{\text{res-o}} \left( \frac{1,752 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{ext-1cm}} (1.0) \right) + \left( ET_{\text{res-i}} \left( \frac{16,416 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{i-total}} \right) \right]$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 1cm soil volume for external exposure.

- Direct External Exposure to contamination 5 cm thick

$$CDI_{\text{res-soil-5cm}} \left( \frac{\text{pCi}}{\text{g}} \right) = C_{\text{soil}} \left( \frac{\text{pCi}}{\text{g}} \right) \times EF_{\text{res}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ACF_{\text{ext-5cm}} \times \left[ \left( ET_{\text{res-o}} \left( \frac{1,752 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{ext-5cm}} (1.0) \right) + \left( ET_{\text{res-i}} \left( \frac{16,416 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{i-total}} \right) \right]$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 5cm soil volume for external exposure.

- Direct External Exposure to contamination 15 cm thick

$$CDI_{\text{res-soil-15cm}} \left( \frac{\text{pCi}}{\text{g}} \right) = C_{\text{soil}} \left( \frac{\text{pCi}}{\text{g}} \right) \times EF_{\text{res}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ACF_{\text{ext-15cm}} \times \left[ \left( ET_{\text{res-o}} \left( \frac{1,752 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{ext-15cm}} (1.0) \right) + \left( ET_{\text{res-i}} \left( \frac{16,416 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{i-total}} \right) \right]$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 15cm soil volume for external exposure.

- Direct External Exposure to surface contamination

$$CDI_{\text{res-soil-gp}} \left( \frac{\text{pCi}}{\text{cm}^2} \right) = C_{\text{soil}} \left( \frac{\text{pCi}}{\text{cm}^2} \right) \times EF_{\text{res}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ACF_{\text{ext-gp}} \times \left[ \left( ET_{\text{res-o}} \left( \frac{1,752 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{ext-gp}} (1.0) \right) + \left( ET_{\text{res-i}} \left( \frac{16,416 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{i-total}} \right) \right]$$

The resulting units for this recommended DCC are in pCi/cm<sup>2</sup>. The units are based on area, because the DCF used is the ground plane for external exposure.


Definitions of the input variables are in [Table 1](#).

**4.1.3 Resident Air**

This receptor spends most, if not all, of the day at home except for the hours spent at work. The activities for this receptor involve typical home making chores (cooking, cleaning, and laundering) as well as gardening. The resident is assumed to be exposed to contaminants via the following pathways: inhalation of ambient air and external radiation from contaminants in ambient air. To take into account the different inhalation rates for children and adults, age-adjusted intake equations were developed to account for changes in intake as the receptor ages.

Two ambient air exposure conditions are offered for this scenario. The first scenario includes a half-life decay function, and the second scenario does not. In situations where the contaminant in the air is not being replenished (e.g., an accidental one-time air release from a factory), equations for the first scenario should be used. In situations where the contaminant in the air has a continual source (e.g., indoor radon from radium in the soil or an operating factory or landfill cap), equations for the second scenario should be used.

**Graphical Representation**



### DCC Equations

The resident ambient air land use equation, presented here, contains the following exposure routes:

- inhalation

$$DCC_{res-air-inh-nodecay} \left( \frac{pCi}{m^3} \right) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_i \left( \frac{mrem}{pCi} \right) \times IFA_{res-adj} \left( \frac{6,195 m^3}{year} \right)}$$

where:

$$IFA_{res-adj} \left( \frac{6,195 m^3}{year} \right) = \left( \left( EF_{res-c} \left( \frac{350 days}{year} \right) \times ET_{res-c} \left( \frac{24 hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times IRA_{res-c} \left( \frac{10 m^3}{day} \right) \times AAF_{res-c} (0.23) \right) + \left( EF_{res-a} \left( \frac{350 days}{year} \right) \times ET_{res-a} \left( \frac{24 hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times IRA_{res-a} \left( \frac{20 m^3}{day} \right) \times AAF_{res-a} (0.77) \right) \right)$$

where:

$$AAF_{res-c} (0.23) = \left( \frac{ED_{res-c} (6 years)}{ED_{res} (26 years)} \right) \text{ and: } AAF_{res-a} (0.77) = \left( \frac{ED_{res-a} (20 years)}{ED_{res} (26 years)} \right)$$

- external exposure to ionizing radiation

$$DCC_{res-air-sub-nodecay} \left( \frac{pCi}{m^3} \right) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{sub} \left( \frac{mrem/year}{pCi/m^3} \right) \times EF_{res} \left( \frac{350 days}{year} \right) \times \left( \frac{1 year}{365 days} \right) \times ET_{res} \left( \frac{24 hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times GSF_a (1.0)}$$

- total

$$DCC_{res-air-tot-nodecay} \left( \frac{pCi}{m^3} \right) = \frac{1}{\frac{1}{DCC_{res-air-inh-nodecay}} + \frac{1}{DCC_{res-air-sub-nodecay}}}$$

Definitions of the input variables are in [Table 1](#).

### CDI Equations

The resident ambient air land use equation, presented here, contains the following exposure routes:

- inhalation

$$CDI_{res-air-inh-nodecay} \left( \frac{pCi}{year} \right) = C_{air} \left( \frac{pCi}{m^3} \right) \times IFA_{res-adj} \left( \frac{6,195 m^3}{year} \right)$$

where:

$$IFA_{res-adj} \left( \frac{6,195 m^3}{year} \right) = \left( \left( EF_{res-c} \left( \frac{350 days}{year} \right) \times ET_{res-c} \left( \frac{24 hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times IRA_{res-c} \left( \frac{10 m^3}{day} \right) \times AAF_{res-c} (0.23) \right) + \left( EF_{res-a} \left( \frac{350 days}{year} \right) \times ET_{res-a} \left( \frac{24 hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times IRA_{res-a} \left( \frac{20 m^3}{day} \right) \times AAF_{res-a} (0.77) \right) \right)$$

where:

$$AAF_{res-c} (0.23) = \left( \frac{ED_{res-c} (6 years)}{ED_{res} (26 years)} \right) \text{ and: } AAF_{res-a} (0.77) = \left( \frac{ED_{res-a} (20 years)}{ED_{res} (26 years)} \right)$$

- external exposure to ionizing radiation

$$CDI_{res-air-sub-nodecay} \left( \frac{pCi}{m^3} \right) = C_{air} \left( \frac{pCi}{m^3} \right) \times EF_{res} \left( \frac{350 days}{year} \right) \times \left( \frac{1 year}{365 days} \right) \times ET_{res} \left( \frac{24 hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times GSF_a (1.0)$$


Definitions of the input variables are in [Table 1](#).

#### 4.1.4 Resident Tapwater

This receptor is exposed to radionuclides that are delivered into a residence. Ingestion of drinking water is an appropriate pathway for all radionuclides. Activities such as showering, laundering, and dish washing also contribute to inhalation. The inhalation exposure route is only calculated for C-14, H-3, Ra-224, and Ra-226, which all volatilize. External exposure to immersion in

tapwater and exposure to produce irrigated with contaminated tapwater are also considered.

### Graphical Representation



### DCC Equations

The tapwater land use equation, presented here, contains the following exposure routes:

- ingestion of tapwater

$$DCC_{\text{water-ing}} (\text{pCi/L}) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_o \left( \frac{\text{mrem}}{\text{pCi}} \right) \times IFW_{\text{res-adj}} \left( \frac{737 \text{ L}}{\text{year}} \right)}$$

where:

$$IFW_{\text{res-adj}} \left( \frac{737 \text{ L}}{\text{year}} \right) = \left( \left( EF_{\text{res-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRW_{\text{res-c}} \left( \frac{0.78 \text{ L}}{\text{day}} \right) \times AAF_{\text{res-c}} (0.23) \right) + \left( EF_{\text{res-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRW_{\text{res-a}} \left( \frac{2.5 \text{ L}}{\text{day}} \right) \times AAF_{\text{res-a}} (0.77) \right) \right)$$

where:

$$AAF_{\text{res-c}} (0.23) = \left( \frac{ED_{\text{res-c}} (6 \text{ years})}{ED_{\text{res}} (26 \text{ years})} \right) \text{ and } AAF_{\text{res-a}} (0.77) = \left( \frac{ED_{\text{res-a}} (20 \text{ years})}{ED_{\text{res}} (26 \text{ years})} \right)$$

- inhalation (The inhalation exposure route is only calculated for C-14, H-3, Ra-224, and Ra-226). Also, volatilization in the equation comes from household uses of water (e.g., showering, laundering, dish washing)

$$DCC_{\text{water-inh}} (\text{pCi/L}) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_i \left( \frac{\text{mrem}}{\text{pCi}} \right) \times IFA_{\text{res-adj}} \left( \frac{6,195 \text{ m}^3}{\text{year}} \right) \times K \left( \frac{0.5 \text{ L}}{\text{m}^3} \right)}$$

where:

$$IFA_{\text{res-adj}} \left( \frac{6,195 \text{ m}^3}{\text{year}} \right) = \left( \left( EF_{\text{res-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times ET_{\text{res-c}} \left( \frac{24 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{res-c}} \left( \frac{10 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{res-c}} (0.23) \right) + \left( EF_{\text{res-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times ET_{\text{res-a}} \left( \frac{24 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{res-a}} \left( \frac{20 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{res-a}} (0.77) \right) \right)$$

where:

$$AAF_{\text{res-c}} (0.23) = \left( \frac{ED_{\text{res-c}} (6 \text{ years})}{ED_{\text{res}} (26 \text{ years})} \right) \text{ and } AAF_{\text{res-a}} (0.77) = \left( \frac{ED_{\text{res-a}} (20 \text{ years})}{ED_{\text{res}} (26 \text{ years})} \right)$$

- immersion

$$DCC_{\text{water-imm}} (\text{pCi/L}) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_{\text{imm}} \left( \frac{\text{mrem/year}}{\text{pCi/L}} \right) \times \left( \frac{1 \text{ year}}{8760 \text{ hours}} \right) \times DFA_{\text{res-adj}} \left( \frac{235 \text{ hours}}{\text{year}} \right)}$$

where:

$$DFA_{\text{res-adj}} \left( \frac{235 \text{ hours}}{\text{year}} \right) = \left( \left( EF_{\text{res-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times EV_{\text{res-c}} \left( \frac{1 \text{ event}}{\text{day}} \right) \times ET_{\text{event-res-c}} \left( \frac{0.54 \text{ hours}}{\text{event}} \right) \times AAF_{\text{res-c}} (0.23) \right) + \left( EF_{\text{res-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times EV_{\text{res-a}} \left( \frac{1 \text{ event}}{\text{day}} \right) \times ET_{\text{event-res-a}} \left( \frac{0.71 \text{ hours}}{\text{event}} \right) \times AAF_{\text{res-a}} (0.77) \right) \right)$$

where:

$$AAF_{\text{res-c}} (0.23) = \left( \frac{ED_{\text{res-c}} (6 \text{ years})}{ED_{\text{res}} (26 \text{ years})} \right) \text{ and } AAF_{\text{res-a}} (0.77) = \left( \frac{ED_{\text{res-a}} (20 \text{ years})}{ED_{\text{res}} (26 \text{ years})} \right)$$

- consumption of produce (fruits and vegetables) - back-calculated to tapwater. Sections 9 and 13 of the 2011 Exposure Factors Handbook were used to derive the intakes for home-grown produce.



$$DCC_{\text{water-res-produce-ing-tot}} \left( \frac{\text{pCi}}{\text{L}} \right) = \frac{1}{\sum_{i=1}^n \frac{1}{DCC_{\text{water-res-produce-ing}} \left( \frac{\text{pCi}}{\text{L}} \right)_i}}$$

where:

n = total number of produce items included

and:

$$DCC_{\text{water-res-produce-ing}} \left( \frac{\text{pCi}}{\text{L}} \right) = \frac{DCC_{\text{res-produce-ing}} \left( \frac{\text{pCi}}{\text{g}} \right)}{\left( \frac{1 \text{ kg}}{1000 \text{ g}} \right) \times \left( Irr_{\text{rup}} \left( \frac{\text{L}}{\text{kg}} \right) + Irr_{\text{res}} \left( \frac{\text{L}}{\text{kg}} \right) + Irr_{\text{dep}} \left( \frac{\text{L}}{\text{kg}} \right) \right)}$$

where:

$$Irr_{\text{res}} \left( \frac{\text{L}}{\text{kg}} \right) = \frac{Ir \left( \frac{\text{L}}{\text{m}^2 \cdot \text{day}} \right) \times F \times MLF_{\text{produce}} \times \left[ 1 - \exp \left( - \left( \frac{\lambda_B}{\text{day}} \right) \times t_b \text{ (days)} \right) \right]}{P \left( \frac{\text{kg}}{\text{m}^2} \right) \times \left( \frac{\lambda_B}{\text{day}} \right)};$$

$$Irr_{\text{rup}} \left( \frac{\text{L}}{\text{kg}} \right) = \frac{Ir \left( \frac{\text{L}}{\text{m}^2 \cdot \text{day}} \right) \times F \times B_{\text{wet}} \times \left[ 1 - \exp \left( - \left( \frac{\lambda_B}{\text{day}} \right) \times t_b \text{ (days)} \right) \right]}{P \left( \frac{\text{kg}}{\text{m}^2} \right) \times \left( \frac{\lambda_B}{\text{day}} \right)};$$

and:

$$Irr_{\text{dep}} \left( \frac{\text{L}}{\text{kg}} \right) = \frac{Ir \left( \frac{\text{L}}{\text{m}^2 \cdot \text{day}} \right) \times F \times I_f \times T \times \left[ 1 - \exp \left( - \left( \frac{\lambda_E}{\text{day}} \right) \times t_b \text{ (days)} \right) \right]}{Y_v \left( \frac{\text{kg}}{\text{m}^2} \right) \times \left( \frac{\lambda_E}{\text{day}} \right)}$$

where:

$$DCC_{\text{res-produce-ing}} \left( \frac{\text{pCi}}{\text{g}} \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_o \left( \frac{\text{mrem}}{\text{pCi}} \right) \times IF_{\text{res-adj}} \left( \frac{\text{g}}{\text{year}} \right) \times CF_{\text{res-produce}} (1)}$$

where:

$$IF_{\text{res-adj}} \left( \frac{\text{g}}{\text{year}} \right) = \left( \left( EF_{\text{res-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IR_{\text{res-c}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{res-c}} (0.23) \right) + \left( EF_{\text{res-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IR_{\text{res-a}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{res-a}} (0.77) \right) \right)$$

and:

$$AAF_{\text{res-c}} (0.23) = \left( \frac{ED_{\text{res-c}} (6 \text{ years})}{ED_{\text{res}} (26 \text{ years})} \right) \text{ and: } AAF_{\text{res-a}} (0.77) = \left( \frac{ED_{\text{res-a}} (20 \text{ years})}{ED_{\text{res}} (26 \text{ years})} \right)$$

- total

$$DCC_{\text{water-tot}} \left( \frac{\text{pCi}}{\text{L}} \right) = \frac{1}{\frac{1}{DCC_{\text{water-ing}}} + \frac{1}{DCC_{\text{water-inh}}} + \frac{1}{DCC_{\text{water-imm}}} + \frac{1}{DCC_{\text{water-produce-tot}}}}$$

Definitions of the input variables are in [Table 1](#).

### CDI Equations

The tapwater land use equation, presented here, contains the following exposure routes:

- ingestion of tapwater

$$CDI_{\text{water-res-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{water}} \left( \frac{\text{pCi}}{\text{L}} \right) \times IFW_{\text{res-adj}} \left( \frac{737 \text{ L}}{\text{year}} \right)$$

where:

$$IFW_{\text{res-adj}} \left( \frac{737 \text{ L}}{\text{year}} \right) = \left( \left( EF_{\text{res-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRW_{\text{res-c}} \left( \frac{0.78 \text{ L}}{\text{day}} \right) \times AAF_{\text{res-c}} (0.23) \right) + \left( EF_{\text{res-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRW_{\text{res-a}} \left( \frac{2.5 \text{ L}}{\text{day}} \right) \times AAF_{\text{res-a}} (0.77) \right) \right)$$

where:

$$AAF_{\text{res-c}} (0.23) = \left( \frac{ED_{\text{res-c}} (6 \text{ years})}{ED_{\text{res}} (26 \text{ years})} \right) \text{ and: } AAF_{\text{res-a}} (0.77) = \left( \frac{ED_{\text{res-a}} (20 \text{ years})}{ED_{\text{res}} (26 \text{ years})} \right)$$

- inhalation (The inhalation exposure route is only calculated for C-14, H-3, Ra-224, and Ra-226). Also, volatilization in the equation comes from household uses of water (e.g., showering, laundering, dish washing)

$$CDI_{\text{water-res-inh}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{water}} \left( \frac{\text{pCi}}{\text{L}} \right) \times IFA_{\text{res-adj}} \left( \frac{6,195 \text{ m}^3}{\text{year}} \right) \times K \left( \frac{0.5 \text{ L}}{\text{m}^3} \right)$$

where:

$$IFA_{\text{res-adj}} \left( \frac{6,195 \text{ m}^3}{\text{year}} \right) = \left( \left( EF_{\text{res-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times ET_{\text{res-c}} \left( \frac{24 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{res-c}} \left( \frac{10 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{res-c}} (0.23) \right) + \left( EF_{\text{res-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times ET_{\text{res-a}} \left( \frac{24 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{res-a}} \left( \frac{20 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{res-a}} (0.77) \right) \right)$$

where:

$$AAF_{\text{res-c}} (0.23) = \left( \frac{ED_{\text{res-c}} (6 \text{ years})}{ED_{\text{res}} (26 \text{ years})} \right) \text{ and: } AAF_{\text{res-a}} (0.77) = \left( \frac{ED_{\text{res-a}} (20 \text{ years})}{ED_{\text{res}} (26 \text{ years})} \right)$$

- immersion

$$CDI_{\text{water-res-imm}} \left( \frac{\text{pCi}}{\text{L}} \right) = C_{\text{water}} \left( \frac{\text{pCi}}{\text{L}} \right) \times \left( \frac{1 \text{ year}}{8760 \text{ hours}} \right) \times DFA_{\text{res-adj}} \left( \frac{235 \text{ hours}}{\text{year}} \right)$$

where:

$$DFA_{\text{res-adj}} \left( \frac{235 \text{ hours}}{\text{year}} \right) = \left( \left( EF_{\text{res-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times EV_{\text{res-c}} \left( \frac{1 \text{ event}}{\text{day}} \right) \times ET_{\text{event-res-c}} \left( \frac{0.54 \text{ hours}}{\text{event}} \right) \times AAF_{\text{res-c}} (0.23) \right) + \left( EF_{\text{res-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times EV_{\text{res-a}} \left( \frac{1 \text{ event}}{\text{day}} \right) \times ET_{\text{event-res-a}} \left( \frac{0.71 \text{ hours}}{\text{event}} \right) \times AAF_{\text{res-a}} (0.77) \right) \right)$$

where:

$$AAF_{\text{res-c}} (0.23) = \left( \frac{ED_{\text{res-c}} (6 \text{ years})}{ED_{\text{res}} (26 \text{ years})} \right) \text{ and: } AAF_{\text{res-a}} (0.77) = \left( \frac{ED_{\text{res-a}} (20 \text{ years})}{ED_{\text{res}} (26 \text{ years})} \right)$$

- consumption of produce (fruits and vegetables) - back-calculated to tapwater. Sections 9 and 13 of the 2011 Exposure Factors Handbook were used to derive the intakes for home-grown produce.

$$CDI_{\text{water-res-produce-ing-tot}} \left( \frac{\text{pCi}}{\text{year}} \right) = \sum_{i=1}^n CDI_{\text{water-res-produce-ing}} \left( \frac{\text{pCi}}{\text{year}} \right)_i$$

where:  
n = total number of produce items included

and:

$$CDI_{\text{water-res-produce-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = \left( C_{\text{water}} \left( \frac{\text{pCi}}{\text{L}} \right) \times IF_{\text{res-adj}} \left( \frac{\text{g}}{\text{year}} \right) \times CF_{\text{res-produce}} (1) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right) \times \left( Irr_{\text{rup}} \left( \frac{\text{L}}{\text{kg}} \right) + Irr_{\text{res}} \left( \frac{\text{L}}{\text{kg}} \right) + Irr_{\text{dep}} \left( \frac{\text{L}}{\text{kg}} \right) \right) \right)$$

where:

$$Irr_{\text{res}} \left( \frac{\text{L}}{\text{kg}} \right) = \frac{Ir \left( \frac{\text{L}}{\text{m}^2 \cdot \text{day}} \right) \times F \times MLF_{\text{produce}} \times \left[ 1 - \exp \left( - \left( \frac{\lambda_B}{\text{day}} \right) \times t_b \text{ (days)} \right) \right]}{P \left( \frac{\text{kg}}{\text{m}^2} \right) \times \left( \frac{\lambda_B}{\text{day}} \right)}$$

$$Irr_{\text{rup}} \left( \frac{\text{L}}{\text{kg}} \right) = \frac{Ir \left( \frac{\text{L}}{\text{m}^2 \cdot \text{day}} \right) \times F \times Bv_{\text{wet}} \times \left[ 1 - \exp \left( - \left( \frac{\lambda_B}{\text{day}} \right) \times t_b \text{ (days)} \right) \right]}{P \left( \frac{\text{kg}}{\text{m}^2} \right) \times \left( \frac{\lambda_B}{\text{day}} \right)}$$

$$Irr_{\text{dep}} \left( \frac{\text{L}}{\text{kg}} \right) = \frac{Ir \left( \frac{\text{L}}{\text{m}^2 \cdot \text{day}} \right) \times F \times I_f \times T \times \left[ 1 - \exp \left( - \left( \frac{\lambda_E}{\text{day}} \right) \times t_b \text{ (days)} \right) \right]}{Y_v \left( \frac{\text{kg}}{\text{m}^2} \right) \times \left( \frac{\lambda_E}{\text{day}} \right)}$$

and:

$$IF_{\text{res-adj}} \left( \frac{\text{g}}{\text{year}} \right) = \left( \left( EF_{\text{res-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IR_{\text{res-c}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{res-c}} (0.23) \right) + \left( EF_{\text{res-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IR_{\text{res-a}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{res-a}} (0.77) \right) \right)$$

where:

$$AAF_{\text{res-c}} (0.23) = \left( \frac{ED_{\text{res-c}} (6 \text{ years})}{ED_{\text{res}} (26 \text{ years})} \right) \text{ and: } AAF_{\text{res-a}} (0.77) = \left( \frac{ED_{\text{res-a}} (20 \text{ years})}{ED_{\text{res}} (26 \text{ years})} \right)$$

Definitions of the input variables are in [Table 1](#).


## 4.2 Composite Worker

### 4.2.1 Composite Worker Soil

This long-term receptor is a full time employee working on-site who spends most of the workday conducting maintenance activities outdoors. The activities for this receptor (e.g., moderate digging, landscaping) typically involve on-site exposures to surface soils. The composite worker is expected to have an elevated soil ingestion rate (100 mg per day) and is assumed to be exposed to contaminants via the following pathways: incidental ingestion of soil, external radiation from contaminants in soil, and inhalation of fugitive dust. The composite worker combines the most protective exposure assumptions of the outdoor and indoor workers. The only difference between the outdoor worker and the composite worker is that the composite worker uses the more protective exposure frequency of 250 days/year from the indoor worker scenario.

This land use is for developing industrial default screening levels that are presented in the [Download Area](#).

### Graphical Representation



DCC Equations

The composite worker soil land use equations, presented here, contain the following exposure routes:

- incidental ingestion of soil

$$DCC_{w-soil-ing} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{oa} \left( \frac{mrem}{pCi} \right) \times EF_w \left( \frac{250 \text{ days}}{year} \right) \times IRS_w \left( \frac{100 \text{ mg}}{day} \right) \times \left( \frac{g}{1000 \text{ mg}} \right)}$$

- inhalation of particulates emitted from soil

$$DCC_{w-soil-inh} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_i \left( \frac{mrem}{pCi} \right) \times EF_w \left( \frac{250 \text{ days}}{year} \right) \times ET_w \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_w \left( \frac{60 \text{ m}^3}{day} \right) \times \frac{1}{PEF \left( \frac{m^3}{kg} \right)} \times \left( \frac{1000 \text{ g}}{kg} \right)}$$

- external exposure to ionizing radiation

$$DCC_{w-soil-ext} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-sv} \left( \frac{mrem/year}{pCi/g} \right) \times EF_w \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_w \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-sv} (1.0) \times ACF_{ext-sv}}$$

- total

$$DCC_{w-soil-tot} (pCi/g) = \frac{1}{\frac{1}{DCC_{w-soil-ing}} + \frac{1}{DCC_{w-soil-inh}} + \frac{1}{DCC_{w-soil-ext}}}$$

Definitions of the input variables are in [Table 1](#).

CDI Equations

The composite worker soil land use equations, presented here, contain the following exposure routes:

- incidental ingestion of soil

$$CDI_{w-soil-ing} \left( \frac{pCi}{year} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_w \left( \frac{250 \text{ days}}{year} \right) \times IRS_w \left( \frac{100 \text{ mg}}{day} \right) \times \left( \frac{g}{1000 \text{ mg}} \right)$$

- inhalation of particulates emitted from soil

$$CDI_{w-soil-inh} \left( \frac{pCi}{year} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_w \left( \frac{250 \text{ days}}{year} \right) \times ET_w \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_w \left( \frac{60 \text{ m}^3}{day} \right) \times \frac{1}{PEF \left( \frac{m^3}{kg} \right)} \times \left( \frac{1000 \text{ g}}{kg} \right)$$

- external exposure to ionizing radiation

$$CDI_{w-soil-ext} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_w \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_w \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-sv} (1.0) \times ACF_{ext-sv}$$

Definitions of the input variables are in [Table 1](#).

4.2.2 Composite Worker Soil 2-D External Exposure

This long-term receptor is a full time employee working on-site who spends most of the workday conducting maintenance activities outdoors. The activities for this receptor (e.g., moderate digging, landscaping) typically involve on-site exposures to surface soils.

This analysis is designed to look at external exposure from contamination of different area sizes. Areas considered are 1 to 1,000,000 square meters. Isotope-specific area correction factors (ACF) were developed for this analysis.

Graphical Representation



### DCC Equations

- Direct External Exposure to contamination at infinite depth

$$DCC_{w-soil-sv} \left( \frac{pCi}{g} \right) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-sv} \left( \frac{mrem/year}{pCi/g} \right) \times EF_w \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_w \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-sv} (1.0) \times ACF_{ext-sv}}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the soil volume for external exposure.

- Direct External Exposure to contamination 1 cm thick

$$DCC_{w-soil-1cm} \left( \frac{pCi}{g} \right) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-1cm} \left( \frac{mrem/year}{pCi/g} \right) \times EF_w \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_w \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-1cm} (1.0) \times ACF_{ext-1cm}}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 1cm soil volume for external exposure.

- Direct External Exposure to contamination 5 cm thick

$$DCC_{w-soil-5cm} \left( \frac{pCi}{g} \right) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-5cm} \left( \frac{mrem/year}{pCi/g} \right) \times EF_w \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_w \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-5cm} (1.0) \times ACF_{ext-5cm}}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 5cm soil volume for external exposure.

- Direct External Exposure to contamination 15 cm thick

$$DCC_{w-soil-15cm} \left( \frac{pCi}{g} \right) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-15cm} \left( \frac{mrem/year}{pCi/g} \right) \times EF_w \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_w \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-15cm} (1.0) \times ACF_{ext-15cm}}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 15cm soil volume for external exposure.

- Direct External Exposure to surface contamination

$$DCC_{w-soil-gp} \left( \frac{pCi}{cm^2} \right) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-gp} \left( \frac{mrem/year}{pCi/cm^2} \right) \times EF_w \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_w \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-gp} (1.0) \times ACF_{ext-gp}}$$

The resulting units for this recommended DCC are in pCi/cm<sup>2</sup>. The units are based on area, because the DCF used is the ground plane for external exposure.

Definitions of the input variables are in [Table 1](#).

### CDI Equations

- Direct External Exposure to contamination at infinite depth

$$CDI_{w-soil-sv} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_w \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_w \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-sv} (1.0) \times ACF_{ext-sv}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the soil volume for external exposure.

- Direct External Exposure to contamination 1 cm thick

$$CDI_{w-soil-1cm} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_w \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_w \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-1cm} (1.0) \times ACF_{ext-1cm}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 1cm soil volume for external exposure.

- Direct External Exposure to contamination 5 cm thick

$$CDI_{w-soil-5cm} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_w \left( \frac{250 \text{ days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_w \left( \frac{8 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-5cm} (1.0) \times ACF_{ext-5cm}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 5cm soil volume for external exposure.

- Direct External Exposure to contamination 15 cm thick

$$CDI_{w-soil-15cm} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_w \left( \frac{250 \text{ days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_w \left( \frac{8 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-15cm} (1.0) \times ACF_{ext-15cm}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 15cm soil volume for external exposure.

- Direct External Exposure to surface contamination

$$CDI_{w-soil-gp} \left( \frac{pCi}{cm^2} \right) = C_{soil} \left( \frac{pCi}{cm^2} \right) \times EF_w \left( \frac{250 \text{ days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_w \left( \frac{8 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-gp} (1.0) \times ACF_{ext-gp}$$

The resulting units for this recommended DCC are in pCi/cm<sup>2</sup>. The units are based on area, because the DCF used is the ground plane for external exposure.


Definitions of the input variables are in [Table 1](#).

4.2.3 Composite Worker Air

This long-term receptor is a full time employee working on-site who spends most of the workday conducting maintenance activities outdoors. The activities for this receptor (e.g., moderate digging, landscaping) typically involve on-site exposures to surface soils. The composite worker is assumed to be exposed to contaminants via the following pathways: inhalation of ambient air and external radiation from contaminants in ambient air. The composite worker combines the most protective exposure assumptions of the outdoor and indoor workers. The only difference between the outdoor worker and the composite worker is that the composite worker uses the more protective exposure frequency of 250 days/year from the indoor worker scenario.

Two ambient air exposure conditions are offered for this scenario. The first scenario includes a half-life decay function and the second scenario does not. In situations where the contaminant in the air is not being replenished (e.g., an accidental one-time air release from a factory), equations for the first scenario should be used. In situations where the contaminant in the air has a continual source (e.g., indoor radon from radium in the soil or an operating factory or landfill cap), equations for the second scenario should be used.

Graphical Representation



DCC Equations

The composite worker ambient air land use equations, presented here, contain the following exposure routes:

- inhalation

$$DCC_{w-air-inh-nodecay} (pCi/m^3) = \frac{DL \left( \frac{mrem}{\text{year}} \right)}{DCF_i \left( \frac{mrem}{pCi} \right) \times EF_w \left( \frac{250 \text{ days}}{\text{year}} \right) \times ET_w \left( \frac{8 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_w \left( \frac{60 \text{ m}^3}{\text{day}} \right)}$$

- external exposure to ionizing radiation

$$DCC_{w-air-sub-nodecay} (pCi/m^3) = \frac{DL \left( \frac{mrem}{\text{year}} \right)}{DCF_{sub} \left( \frac{mrem/year}{pCi/m^3} \right) \times EF_w \left( \frac{250 \text{ days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_w \left( \frac{8 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_a (1.0)}$$

- total

$$DCC_{w-air-tot-nodecay} (pCi/m^3) = \frac{1}{\frac{1}{DCC_{w-air-inh-nodecay}} + \frac{1}{DCC_{w-air-sub-nodecay}}}$$

Definitions of the input variables are in [Table 1](#).



### CDI Equations

The composite worker ambient air land use equation, presented here, contains the following exposure routes:

- inhalation

$$CDI_{w-air-inh-no decay} \left( \frac{pCi}{year} \right) = C_{air} \left( \frac{pCi}{m^3} \right) \times EF_w \left( \frac{250 \text{ days}}{year} \right) \times ET_w \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_w \left( \frac{60 \text{ m}^3}{day} \right)$$

- external exposure to ionizing radiation

$$CDI_{w-air-sub-no decay} \left( \frac{pCi}{m^3} \right) = C_{air} \left( \frac{pCi}{m^3} \right) \times EF_w \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_w \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_a (1.0)$$

Definitions of the input variables are in [Table 1](#).


## 4.3 Outdoor Worker

### 4.3.1 Outdoor Worker Soil

This long-term receptor is a full time employee working on-site who spends most of the workday conducting maintenance activities outdoors. The activities for this receptor (e.g., moderate digging, landscaping) typically involve on-site exposures to surface soils. The outdoor worker is expected to have an elevated soil ingestion rate (100 mg per day) and is assumed to be exposed to contaminants via the following pathways: incidental ingestion of soil, external radiation from contaminants in soil, and inhalation of fugitive dust. The outdoor worker receives more exposure than the indoor worker under commercial/industrial conditions.

The outdoor worker soil land use is not provided in the [DCC Download Area](#), but DCCs can be created by using the Calculator to modify the exposure parameters for the composite worker to match the equations that follow.

### Graphical Representation



### DCC Equations

The outdoor worker soil land use equations, presented here, contain the following exposure routes:

- incidental ingestion of soil

$$DCC_{ow-soil-ing} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{oa} \left( \frac{mrem}{pCi} \right) \times EF_{ow} \left( \frac{225 \text{ days}}{year} \right) \times IRS_{ow} \left( \frac{100 \text{ mg}}{day} \right) \times \left( \frac{g}{1000 \text{ mg}} \right)}$$

- inhalation of particulates emitted from soil

$$DCC_{ow-soil-inh} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_i \left( \frac{risk}{pCi} \right) \times EF_{ow} \left( \frac{225 \text{ days}}{year} \right) \times ET_{ow} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{ow} \left( \frac{60 \text{ m}^3}{day} \right) \times \frac{1}{PEF} \left( \frac{m^3}{kg} \right) \times \left( \frac{1000 \text{ g}}{kg} \right)}$$

- external exposure to ionizing radiation

$$DCC_{ow-soil-ext} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-sv} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{ow} \left( \frac{225 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{ow} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-sv} (1.0) \times ACF_{ext-sv}}$$

- total

$$DCC_{ow-soil-tot} (pCi/g) = \frac{1}{\frac{1}{DCC_{ow-soil-ing}} + \frac{1}{DCC_{ow-soil-inh}} + \frac{1}{DCC_{ow-soil-ext}}}$$

Definitions of the input variables are in [Table 1](#).

### CDI Equations

The outdoor worker soil land use equations, presented here, contain the following exposure routes:

- incidental ingestion of soil

$$CDI_{ow-soil-ing} \left( \frac{pCi}{year} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{ow} \left( \frac{225 \text{ days}}{year} \right) \times IRS_{ow} \left( \frac{100 \text{ mg}}{day} \right) \times \left( \frac{g}{1000 \text{ mg}} \right)$$

- inhalation of particulates emitted from soil/li>

$$CDI_{ow-soil-inh} \left( \frac{pCi}{year} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{ow} \left( \frac{225 \text{ days}}{year} \right) \times ET_{ow} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{ow} \left( \frac{60 \text{ m}^3}{day} \right) \times \frac{1}{PEF \left( \frac{\text{m}^3}{kg} \right)} \times \left( \frac{1000 \text{ g}}{kg} \right)$$

- external exposure to ionizing radiation

$$CDI_{ow-soil-ext} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{ow} \left( \frac{225 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{ow} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-sv} (1.0) \times ACF_{ext-sv}$$

Definitions of the input variables are in [Table 1](#).

### 4.3.2 Outdoor Worker Soil 2-D External Exposure

This long-term receptor is a full time employee working on-site who spends most of the workday conducting maintenance activities outdoors. The activities for this receptor (e.g., moderate digging, landscaping) typically involve on-site exposures to surface soils.

This analysis is designed to look at external exposure from contamination of different area sizes. Areas considered are 1 to 1,000,000 square meters. Isotope-specific area correction factors (ACF) were developed for this analysis.

#### Graphical Representation



#### DCC Equations

- Direct External Exposure to contamination at infinite depth

$$DCC_{ow-soil-sv} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-sv} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{ow} \left( \frac{225 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{ow} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-sv} (1.0) \times ACF_{ext-sv}}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the soil volume for external exposure.

- Direct External Exposure to contamination 1 cm thick

$$DCC_{ow-soil-1cm} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-1cm} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{ow} \left( \frac{225 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{ow} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-1cm} (1.0) \times ACF_{ext-1cm}}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 1cm soil volume for external exposure.

- Direct External Exposure to contamination 5 cm thick

$$DCC_{ow-soil-5cm} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-5cm} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{ow} \left( \frac{225 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{ow} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-5cm} (1.0) \times ACF_{ext-5cm}}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 5cm soil volume for external exposure.

- Direct External Exposure to contamination 15 cm thick

$$DCC_{ow-soil-15cm} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-15cm} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{ow} \left( \frac{225 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{ow} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-15cm} (1.0) \times ACF_{ext-15cm}}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 15cm soil volume for external exposure.

- Direct External Exposure to surface contamination

$$DCC_{ow-soil-gp} \left( \frac{pCi}{cm^2} \right) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-gp} \left( \frac{mrem/year}{pCi/cm^2} \right) \times EF_{ow} \left( \frac{225 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{ow} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-gp} (1.0) \times ACF_{ext-gp}}$$

The resulting units for this recommended DCC are in pCi/cm<sup>2</sup>. The units are based on area, because the DCF used is the ground plane for external exposure.

Definitions of the input variables are in [Table 1](#).

CDI Equations

- Direct External Exposure to contamination at infinite depth

$$CDI_{ow-soil-sv} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{ow} \left( \frac{225 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{ow} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-sv} (1.0) \times ACF_{ext-sv}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the soil volume for external exposure.

- Direct External Exposure to contamination 1 cm thick

$$CDI_{ow-soil-1cm} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{ow} \left( \frac{225 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{ow} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-1cm} (1.0) \times ACF_{ext-1cm}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 1cm soil volume for external exposure.

- Direct External Exposure to contamination 5 cm thick

$$CDI_{ow-soil-5cm} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{ow} \left( \frac{225 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{ow} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-5cm} (1.0) \times ACF_{ext-5cm}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 5cm soil volume for external exposure.

- Direct External Exposure to contamination 15 cm thick

$$CDI_{ow-soil-15cm} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{ow} \left( \frac{225 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{ow} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-15cm} (1.0) \times ACF_{ext-15cm}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 15cm soil volume for external exposure.

- Direct External Exposure to surface contamination

$$CDI_{ow-soil-gp} \left( \frac{pCi}{cm^2} \right) = C_{soil} \left( \frac{pCi}{cm^2} \right) \times EF_{ow} \left( \frac{225 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{ow} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-gp} (1.0) \times ACF_{ext-gp}$$

The resulting units for this recommended DCC are in pCi/cm<sup>2</sup>. The units are based on area, because the DCF used is the ground plane for external exposure.


Definitions of the input variables are in [Table 1](#).

4.3.3 Outdoor Worker Air

This long-term receptor is a full time employee working on-site who spends most of the workday conducting maintenance activities outdoors. The activities for this receptor (e.g., moderate digging, landscaping) typically involve on-site exposures to surface soils. The outdoor worker is assumed to be exposed to contaminants via the following pathways: inhalation of ambient air and external radiation from contaminants in ambient air.

Two ambient air exposure conditions are offered for this scenario. The first scenario includes a half-life decay function and the second scenario does not. In situations where the contaminant in the air is not being replenished (e.g., an accidental one-time air release from a factory), equations for the first scenario should be used. In situations where the contaminant in the air has a continual source (e.g., indoor radon from radium in the soil, or an operating factory or landfill cap), equations for the second scenario should be used.

Graphical Representation



DCC Equations

The outdoor worker ambient air land use equation, presented here, contains the following exposure routes:

- inhalation

$$DCC_{ow-air-inh-no decay} \left( \frac{pCi}{m^3} \right) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_i \left( \frac{mrem}{pCi} \right) \times EF_{ow} \left( \frac{225 \text{ days}}{year} \right) \times ET_{ow} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{ow} \left( \frac{60 \text{ m}^3}{day} \right)}$$

- external exposure to ionizing radiation

$$DCC_{ow-air-sub-no decay} \left( \frac{pCi}{m^3} \right) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{sub} \left( \frac{mrem/year}{pCi/m^3} \right) \times EF_{ow} \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ day}} \right) \times ET_{ow} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_a (1.0)}$$

- total

$$DCC_{ow-air-tot-no decay} \left( \frac{pCi}{m^3} \right) = \frac{1}{\frac{1}{DCC_{ow-air-inh-no decay}} + \frac{1}{DCC_{ow-air-sub-no decay}}}$$

Definitions of the input variables are in [Table 1](#).

CDI Equations

The outdoor worker ambient air land use equation, presented here, contains the following exposure routes:

- inhalation

$$CDI_{ow-air-inh-no decay} \left( \frac{pCi}{year} \right) = C_{air} \left( \frac{pCi}{m^3} \right) \times EF_{ow} \left( \frac{225 \text{ days}}{year} \right) \times ET_{ow} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{ow} \left( \frac{60 \text{ m}^3}{day} \right)$$

- external exposure to ionizing radiation

$$CDI_{ow-air-sub-no decay} \left( \frac{pCi}{m^3} \right) = C_{air} \left( \frac{pCi}{m^3} \right) \times EF_{ow} \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ day}} \right) \times ET_{ow} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_a (1.0)$$

Definitions of the input variables are in [Table 1](#).


4.4 Indoor Worker

4.4.1 Indoor Worker Soil

This receptor spends most, if not all, of the workday indoors. Thus, an indoor worker has no direct contact with outdoor soils. This worker may, however, be exposed to contaminants through ingestion of contaminated soils that have been incorporated into indoor dust, external radiation from contaminants in soil, and the inhalation of contaminants present in indoor air. DCCs calculated for this receptor are expected to be protective of both workers engaged in low intensity activities, such as office work, and those engaged in more strenuous activity (e.g., factory or warehouse workers).

The indoor worker soil land use is not provided in the [DCC Download Area](#), but DCCs can be created by using the Calculator to modify the exposure parameters for the composite worker to match the equations that follow.

Graphical Representation



### DCC Equations

The indoor worker soil land use equation, presented here, contains the following exposure routes:

- incidental ingestion of soil

$$DCC_{iw-soil-ing} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{oa} \left( \frac{mrem}{pCi} \right) \times EF_{iw} \left( \frac{250 \text{ days}}{year} \right) \times IRS_{iw} \left( \frac{50 \text{ mg}}{day} \right) \times \left( \frac{g}{1000 \text{ mg}} \right)}$$

- inhalation of particulates emitted from soil

$$DCC_{iw-soil-inh} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{SF_i \left( \frac{mrem}{pCi} \right) \times EF_{iw} \left( \frac{250 \text{ days}}{year} \right) \times ET_{iw} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{iw} \left( \frac{60 \text{ m}^3}{day} \right) \times \frac{1}{PEF \left( \frac{m^3}{kg} \right)} \times \left( \frac{1000 \text{ g}}{kg} \right)}$$

- external exposure to ionizing radiation

$$DCC_{iw-soil-ext} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-sv} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{iw} \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ day}} \right) \times ET_{iw} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{i-total} \times ACF_{ext-sv}}$$

- total

$$DCC_{iw-soil-tot} (pCi/g) = \frac{1}{\frac{1}{DCC_{iw-soil-ing}} + \frac{1}{DCC_{iw-soil-inh}} + \frac{1}{DCC_{iw-soil-ext}}}$$

Definitions of the input variables are in [Table 1](#).

### CDI Equations

The indoor worker soil land use equation, presented here, contains the following exposure routes:

- incidental ingestion of soil

$$CDI_{iw-soil-ing} \left( \frac{pCi}{year} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{iw} \left( \frac{250 \text{ days}}{year} \right) \times IRS_{iw} \left( \frac{50 \text{ mg}}{day} \right) \times \left( \frac{g}{1000 \text{ mg}} \right)$$

- inhalation of particulates emitted from soil

$$CDI_{iw-soil-inh} \left( \frac{pCi}{year} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{iw} \left( \frac{250 \text{ days}}{year} \right) \times ET_{iw} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{iw} \left( \frac{60 \text{ m}^3}{day} \right) \times \frac{1}{PEF \left( \frac{m^3}{kg} \right)} \times \left( \frac{1000 \text{ g}}{kg} \right)$$

- external exposure to ionizing radiation

$$CDI_{ow-soil-ext} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{iw} \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ day}} \right) \times ET_{iw} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{i-total} \times ACF_{ext-sv}$$


Definitions of the input variables are in [Table 1](#).

### 4.4.2 Indoor Worker Soil 2-D External Exposure

This receptor spends most, if not all, of the workday indoors. Thus, an indoor worker has no direct contact with outdoor soils. A gamma shielding factor is applied for this scenario to account for shielding provided by floors and foundation slabs.

This analysis is designed to look at external exposure from contamination of different area sizes. Areas considered are 1 to 1,000,000 square meters. Isotope-specific area correction factors (ACF) were developed for this analysis.

### Graphical Representation



### DCC Equations

- Direct External Exposure to contamination at infinite depth

$$DCC_{IW-soil-sv} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-sv} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{IW} \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{IW} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{i-total} \times ACF_{ext-sv}}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the soil volume for external exposure.

- Direct External Exposure to contamination 1 cm thick

$$DCC_{IW-soil-1cm} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-1cm} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{IW} \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{IW} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{i-total} \times ACF_{ext-1cm}}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 1cm soil volume for external exposure.

- Direct External Exposure to contamination 5 cm thick

$$DCC_{IW-soil-5cm} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-5cm} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{IW} \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{IW} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{i-total} \times ACF_{ext-5cm}}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 5cm soil volume for external exposure.

- Direct External Exposure to contamination 15 cm thick

$$DCC_{IW-soil-15cm} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-15cm} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{IW} \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{IW} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{i-total} \times ACF_{ext-15cm}}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 15cm soil volume for external exposure.

- Direct External Exposure to surface contamination

$$DCC_{IW-soil-gp} (pCi/cm^2) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-gp} \left( \frac{mrem/year}{pCi/cm^2} \right) \times EF_{IW} \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{IW} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{i-total} \times ACF_{ext-gp}}$$

The resulting units for this recommended DCC are in pCi/cm<sup>2</sup>. The units are based on area, because the DCF used is the ground plane for external exposure.

Definitions of the input variables are in [Table 1](#).

### CDI Equations

- Direct External Exposure to contamination at infinite depth

$$CDI_{ow-soil-sv} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{IW} \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ day}} \right) \times ET_{IW} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{i-total} \times ACF_{ext-sv}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the soil volume for external exposure.

- Direct External Exposure to contamination 1 cm thick

$$CDI_{ow-soil-1cm} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{IW} \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ day}} \right) \times ET_{IW} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{i-total} \times ACF_{ext-1cm}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 1cm soil volume for external exposure.

- Direct External Exposure to contamination 5 cm thick



$$CDI_{ow-soil-5cm} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{iw} \left( \frac{250 \text{ days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ day}} \right) \times ET_{iw} \left( \frac{8 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{i-total} \times ACF_{ext-5cm}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 5cm soil volume for external exposure.

- Direct External Exposure to contamination 15 cm thick

$$CDI_{ow-soil-15cm} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{iw} \left( \frac{250 \text{ days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ day}} \right) \times ET_{iw} \left( \frac{8 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{i-total} \times ACF_{ext-15cm}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 15cm soil volume for external exposure.

- Direct External Exposure to surface contamination

$$CDI_{ow-soil-gp} \left( \frac{pCi}{cm^2} \right) = C_{soil} \left( \frac{pCi}{cm^2} \right) \times EF_{iw} \left( \frac{250 \text{ days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ day}} \right) \times ET_{iw} \left( \frac{8 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{i-total} \times ACF_{ext-gp}$$

The resulting units for this recommended DCC are in pCi/cm<sup>2</sup>. The units are based on area, because the DCF used is the ground plane for external exposure.


Definitions of the input variables are in [Table 1](#).

### 4.4.3 Indoor Worker Air

This long-term receptor is a full time employee working on-site who spends most, if not all, of the workday indoors. The indoor worker is assumed to be exposed to contaminants via the following pathways: inhalation of ambient air and external radiation from contaminants in ambient air.

Two ambient air exposure conditions are offered for this scenario. The first scenario includes a half-life decay function and the second scenario does not. In situations where the contaminant in the air is not being replenished (e.g., an accidental one-time air release from a factory), equations for the first scenario should be used. In situations where the contaminant in the air has a continual source (e.g., indoor radon from radium in the soil or an operating factory or landfill cap), equations for the second scenario should be used.

#### Graphical Representation



#### DCC Equations

The indoor worker ambient air land use equation, presented here, contains the following exposure routes:

- inhalation

$$DCC_{iw-air-inh-nodecay} \left( pCi/m^3 \right) = \frac{DL \left( \frac{mrem}{\text{year}} \right)}{DCF_i \left( \frac{mrem}{pCi} \right) \times EF_{iw} \left( \frac{250 \text{ days}}{\text{year}} \right) \times ET_{iw} \left( \frac{8 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{iw} \left( \frac{60 \text{ m}^3}{\text{day}} \right)}$$

- external exposure to ionizing radiation

$$DCC_{iw-air-sub-nodecay} \left( pCi/m^3 \right) = \frac{DL \left( \frac{mrem}{\text{year}} \right)}{DCF_{sub} \left( \frac{mrem/year}{pCi/m^3} \right) \times EF_{iw} \left( \frac{250 \text{ days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ day}} \right) \times ET_{iw} \left( \frac{8 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_a (1.0)}$$

- total

$$DCC_{iw-air-tot-nodecay} \left( pCi/m^3 \right) = \frac{1}{\frac{1}{DCC_{iw-air-inh-nodecay}} + \frac{1}{DCC_{iw-air-sub-nodecay}}}$$

Definitions of the input variables are in [Table 1](#).

#### CDI Equations

The indoor worker ambient air land use equation, presented here, contains the following exposure routes:

- inhalation

$$CDI_{lw-air-inh-nodecay} \left( \frac{pCi}{year} \right) = C_{air} \left( \frac{pCi}{m^3} \right) \times EF_{lw} \left( \frac{250 \text{ days}}{year} \right) \times ET_{lw} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{lw} \left( \frac{60 \text{ m}^3}{day} \right)$$

- external exposure to ionizing radiation

$$CDI_{lw-air-sub-nodecay} \left( \frac{pCi}{m^3} \right) = C_{air} \left( \frac{pCi}{m^3} \right) \times EF_{lw} \left( \frac{250 \text{ days}}{year} \right) \times \left( \frac{1 \text{ year}}{365 \text{ day}} \right) \times ET_{lw} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_a (1.0)$$

Definitions of the input variables are in [Table 1](#).

## 4.5 Construction Worker

### 4.5.1 Construction Worker Soil Exposure to Unpaved Road Traffic

This short-term receptor is exposed during the work day to vehicles suspending dust in the air. The activities for this receptor (e.g., trenching, excavating) typically involve on-site exposures to surface soils. The construction worker is expected to have an elevated soil ingestion rate (330 mg per day) and is assumed to be exposed to contaminants via the following pathways: incidental ingestion of soil, external radiation from contaminants in soil, and inhalation of fugitive dust. The only difference between this construction worker and the one described in section 4.5.3 is that this construction worker uses a different PEF.

The construction worker soil land use is not provided in the Generic Tables, but DCCs can be created by using the Calculator. The construction land use is described in the [supplemental soil screening guidance](#). This land use is limited to an exposure duration of 1 year and is thus subchronic. Other unique aspects of this scenario are that the PEF is based on mechanical disturbance of the soil. Two types of mechanical soil disturbance are addressed: standard vehicle traffic and other than standard vehicle traffic (e.g. wind, grading, dozing, tilling and excavating). In general, the intakes and contact rates are all greater than the outdoor worker. Exhibit 5-1 in the [supplemental soil screening guidance](#) presents the exposure parameters.

#### Graphical Representation



#### DCC Equations

The construction worker soil land use equations, presented here, contain the following exposure routes:

- incidental ingestion of soil

$$DCC_{cw-soil-ing} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{oa} \left( \frac{mrem}{pCi} \right) \times EF_{cw} \left( \frac{EW_{cw} \text{ 50 weeks}}{year} \times DW_{cw} \frac{5 \text{ days}}{week} \right) \times IRS_{cw} \left( \frac{330 \text{ mg}}{day} \right) \times \left( \frac{g}{1000 \text{ mg}} \right)}$$

- inhalation of particulates emitted from soil

$$DCC_{cw-soil-inh} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_i \left( \frac{mrem}{pCi} \right) \times EF_{cw} \left( \frac{EW_{cw} \text{ 50 weeks}}{year} \times DW_{cw} \frac{5 \text{ days}}{week} \right) \times ET_{cw} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{cw} \left( \frac{60 \text{ m}^3}{day} \right) \times \frac{1}{PEF_{sc} \left( \frac{m^3}{kg} \right) \times \left( \frac{1000 \text{ g}}{kg} \right)}$$

- external exposure to ionizing radiation

$$DCC_{cw-soil-ext} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-sv} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{cw} \left( \frac{EW_{cw} \text{ 50 weeks}}{year} \times DW_{cw} \frac{5 \text{ days}}{week} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{cw} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-sv} (1.0) \times ACF_{ext-sv}}$$

- total

$$DCC_{cw-soil-tot} (pCi/g) = \frac{1}{\frac{1}{DCC_{cw-soil-ing}} + \frac{1}{DCC_{cw-soil-inh}} + \frac{1}{DCC_{cw-soil-ext}}}$$

Definitions of the input variables are in [Table 1](#).

CDI Equations

The construction worker soil land use equations, presented here, contain the following exposure routes:

- incidental ingestion of soil

$$DCC_{cw-soil-ing} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{oa} \left( \frac{mrem}{pCi} \right) \times EF_{cw} \left( \frac{EW_{cw} \ 50 \ weeks}{year} \times \frac{DW_{cw} \ 5 \ days}{week} \right) \times IRS_{cw} \left( \frac{330 \ mg}{day} \right) \times \left( \frac{g}{1000 \ mg} \right)}$$

- inhalation of particulates emitted from soil

$$DCC_{cw-soil-inh} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_i \left( \frac{mrem}{pCi} \right) \times EF_{cw} \left( \frac{EW_{cw} \ 50 \ weeks}{year} \times \frac{DW_{cw} \ 5 \ days}{week} \right) \times ET_{cw} \left( \frac{8 \ hours}{day} \right) \times \left( \frac{1 \ day}{24 \ hours} \right) \times IRA_{cw} \left( \frac{60 \ m^3}{day} \right) \times \frac{1}{PEF_{sc} \left( \frac{m^3}{kg} \right)} \times \left( \frac{1000 \ g}{kg} \right)}$$

- external exposure to ionizing radiation

$$DCC_{cw-soil-ext} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-sv} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{cw} \left( \frac{EW_{cw} \ 50 \ weeks}{year} \times \frac{DW_{cw} \ 5 \ days}{week} \right) \times \left( \frac{1 \ year}{365 \ days} \right) \times ET_{cw} \left( \frac{8 \ hours}{day} \right) \times \left( \frac{1 \ day}{24 \ hours} \right) \times GSF_{ext-sv} (1.0) \times ACF_{ext-sv}}$$

Definitions of the input variables are in [Table 1](#).

4.5.2 Construction Worker Soil Exposure to Unpaved Road Traffic 2-D External Exposure

This short-term receptor is a full time employee working on-site and who spends most of the workday conducting maintenance activities outdoors. The activities for this receptor (e.g., trenching, excavating, wind, grading, dozing, and tilling) typically involve on-site exposures to surface soils.

This analysis is designed to look at external exposure from contamination of different area sizes. Areas considered are 1 to 1,000,000 square meters. Isotope-specific area correction factors (ACF) were developed for this analysis.

Graphical Representation



Equations

- Direct External Exposure to contamination at infinite depth

$$DCC_{cw-soil-sv} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-sv} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{cw} \left( \frac{EW_{cw} \ 50 \ weeks}{year} \times \frac{DW_{cw} \ 5 \ days}{week} \right) \times \left( \frac{1 \ year}{365 \ days} \right) \times ET_{cw} \left( \frac{8 \ hours}{day} \right) \times \left( \frac{1 \ day}{24 \ hours} \right) \times GSF_{ext-sv} (1.0) \times ACF_{ext-sv}}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the soil volume for external exposure.

- Direct External Exposure to contamination 1 cm thick

$$DCC_{cw-soil-1cm} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-1cm} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{cw} \left( \frac{EW_{cw} \ 50 \ weeks}{year} \times \frac{DW_{cw} \ 5 \ days}{week} \right) \times \left( \frac{1 \ year}{365 \ days} \right) \times ET_{cw} \left( \frac{8 \ hours}{day} \right) \times \left( \frac{1 \ day}{24 \ hours} \right) \times GSF_{ext-1cm} (1.0) \times ACF_{ext-1cm}}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 1cm soil volume for external exposure.

- Direct External Exposure to contamination 5 cm thick

$$DCC_{cw-soil-5cm} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-5cm} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{cw} \left( \frac{EW_{cw} \frac{50 weeks}{year} \times DW_{cw} \frac{5 days}{week} \right) \times \left( \frac{1 year}{365 days} \right) \times ET_{cw} \left( \frac{8 hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times GSF_{ext-5cm} (1.0) \times ACF_{ext-5cm}}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 5cm soil volume for external exposure.

- Direct External Exposure to contamination 15 cm thick

$$DCC_{cw-soil-15cm} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-15cm} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{cw} \left( \frac{EW_{cw} \frac{50 weeks}{year} \times DW_{cw} \frac{5 days}{week} \right) \times \left( \frac{1 year}{365 days} \right) \times ET_{cw} \left( \frac{8 hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times GSF_{ext-15cm} (1.0) \times ACF_{ext-15cm}}$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 15cm soil volume for external exposure.

- Direct External Exposure to surface contamination

$$DCC_{cw-soil-gp} (pCi/cm^2) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-gp} \left( \frac{mrem/year}{pCi/cm^2} \right) \times EF_{cw} \left( \frac{EW_{cw} \frac{50 weeks}{year} \times DW_{cw} \frac{5 days}{week} \right) \times \left( \frac{1 year}{365 days} \right) \times ET_{cw} \left( \frac{8 hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times GSF_{ext-gp} (1.0) \times ACF_{ext-gp}}$$

The resulting units for this recommended DCC are in pCi/cm<sup>2</sup>. The units are based on area, because the DCF used is the ground plane for external exposure.

Definitions of the input variables are in [Table 1](#).

#### Equations

- Direct External Exposure to contamination at infinite depth

$$CDI_{cw-soil-sv} \left( \frac{pCi}{g} \right) = \left( \frac{C_{soil} \left( \frac{pCi}{g} \right) \times EF_{cw} \left( \frac{EW_{cw} \frac{50 weeks}{year} \times DW_{cw} \frac{5 days}{week} \right) \times \left( \frac{1 year}{365 day} \right) \times ET_{cw} \left( \frac{8 hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times GSF_{ext-sv} (1.0) \times ACF_{ext-sv}} \right)$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the soil volume for external exposure.

- Direct External Exposure to contamination 1 cm thick

$$CDI_{cw-soil-1cm} \left( \frac{pCi}{g} \right) = \left( \frac{C_{soil} \left( \frac{pCi}{g} \right) \times EF_{cw} \left( \frac{EW_{cw} \frac{50 weeks}{year} \times DW_{cw} \frac{5 days}{week} \right) \times \left( \frac{1 year}{365 day} \right) \times ET_{cw} \left( \frac{8 hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times GSF_{ext-1cm} (1.0) \times ACF_{ext-1cm}} \right)$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 1cm soil volume for external exposure.

- Direct External Exposure to contamination 5 cm thick

$$CDI_{cw-soil-5cm} \left( \frac{pCi}{g} \right) = \left( \frac{C_{soil} \left( \frac{pCi}{g} \right) \times EF_{cw} \left( \frac{EW_{cw} \frac{50 weeks}{year} \times DW_{cw} \frac{5 days}{week} \right) \times \left( \frac{1 year}{365 day} \right) \times ET_{cw} \left( \frac{8 hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times GSF_{ext-5cm} (1.0) \times ACF_{ext-5cm}} \right)$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 5cm soil volume for external exposure.

- Direct External Exposure to contamination 15 cm thick

$$CDI_{cw-soil-15cm} \left( \frac{pCi}{g} \right) = \left( \frac{C_{soil} \left( \frac{pCi}{g} \right) \times EF_{cw} \left( \frac{EW_{cw} \frac{50 weeks}{year} \times DW_{cw} \frac{5 days}{week} \right) \times \left( \frac{1 year}{365 day} \right) \times ET_{cw} \left( \frac{8 hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times GSF_{ext-15cm} (1.0) \times ACF_{ext-15cm}} \right)$$

The resulting units for this recommended DCC are in pCi/g. The units are based on mass, because the DCF used is the 15cm soil volume for external exposure.

- Direct External Exposure to surface contamination

$$CDI_{cw-soil-gp} \left( \frac{pCi}{cm^2} \right) = \left( \frac{C_{soil} \left( \frac{pCi}{cm^2} \right) \times EF_{cw} \left( \frac{EW_{cw} \frac{50 weeks}{year} \times DW_{cw} \frac{5 days}{week} \right) \times \left( \frac{1 year}{365 day} \right) \times ET_{cw} \left( \frac{8 hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times GSF_{ext-gp} (1.0) \times ACF_{ext-gp}} \right)$$

The resulting units for this recommended DCC are in pCi/cm<sup>2</sup>. The units are based on area, because the DCF used is the ground plane for external exposure.

Definitions of the input variables are in [Table 1](#).

### 4-5.3 Construction Worker Air

This short-term receptor is exposed during the work day to heavy construction activities outdoors. The activities for this receptor

(e.g., trenching, excavating, wind, grading, dozing, and tilling) typically involve on-site exposures to surface soils. The construction worker is assumed to be exposed to contaminants via the following pathways: inhalation of ambient air and external radiation from contaminants in ambient air.

Two ambient air exposure conditions are offered for this scenario. The first scenario includes a half-life decay function and the second scenario does not. In situations where the contaminant in the air is not being replenished (e.g., an accidental one-time air release from a factory), equations for the first scenario should be used. In situations where the contaminant in the air has a continual source (e.g., indoor radon from radium in the soil or an operating factory or landfill cap), equations for the second scenario should be used.

Graphical Representation



DCC Equations

The composite worker ambient air land use equation, presented here, contains the following exposure routes without half-life decay:

- inhalation

$$DCC_{cw-air-inh-nodecay} \left( \frac{pCi}{m^3} \right) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_i \left( \frac{mrem}{pCi} \right) \times EF_{cw} \left( \frac{50 \text{ weeks}}{year} \right) \times DW_{cw} \left( \frac{5 \text{ days}}{week} \right) \times ET_{cw} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{cw} \left( \frac{60 \text{ m}^3}{day} \right)}$$

- external exposure to ionizing radiation

$$DCC_{cw-air-sub-nodecay} \left( \frac{pCi}{m^3} \right) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{sub} \left( \frac{mrem/year}{pCi/m^3} \right) \times EF_{cw} \left( \frac{50 \text{ weeks}}{year} \right) \times DW_{cw} \left( \frac{5 \text{ days}}{week} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{cw} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_a (1.0)}$$

- total

$$DCC_{cw-air-tot-nodecay} \left( \frac{pCi}{m^3} \right) = \frac{1}{\frac{1}{DCC_{cw-air-inh-nodecay}} + \frac{1}{DCC_{cw-air-sub-nodecay}}}$$

Definitions of the input variables are in [Table 1](#).

CDI Equations

The composite worker ambient air land use equation, presented here, contains the following exposure routes without half-life decay:

- inhalation

$$CDI_{cw-air-inh-nodecay} \left( \frac{pCi}{year} \right) = C_{air} \left( \frac{pCi}{m^3} \right) \times EF_{cw} \left( \frac{50 \text{ weeks}}{year} \right) \times DW_{cw} \left( \frac{5 \text{ days}}{week} \right) \times ET_{cw} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{cw} \left( \frac{60 \text{ m}^3}{day} \right)$$

- external exposure to ionizing radiation

$$CDI_{cw-air-sub-nodecay} \left( \frac{pCi}{m^3} \right) = C_{air} \left( \frac{pCi}{m^3} \right) \times EF_{cw} \left( \frac{50 \text{ weeks}}{year} \right) \times DW_{cw} \left( \frac{5 \text{ days}}{week} \right) \times \left( \frac{1 \text{ yr}}{365 \text{ days}} \right) \times ET_{cw} \left( \frac{8 \text{ hr}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_a (1.0)$$

Definitions of the input variables are in [Table 1](#).

4-5-4 Construction Worker Soil Exposure to Other Construction Activities

This short-term receptor is exposed during the work day to heavy vehicles suspending dust in the air. The activities for this receptor (e.g., dozing, grading, tilling, dumping, nd excavating) typically involve on-site exposures to surface soils. The construction worker is expected to have an elevated soil ingestion rate (330 mg per day) and is assumed to be exposed to contaminants via the following pathways: incidental ingestion of soil, external radiation from contaminants in soil, and inhalation of fugitive dust. The only difference between this construction worker and the one described in section 4.5.1 is that this construction worker uses a different PEF.

The construction worker soil land use is not provided in the Generic Tables, but DCCs can be created by using the Calculator. The construction land use is described in the [supplemental soil screening guidance](#). This land use is limited to an exposure duration of

1 year and is thus subchronic. Other unique aspects of this scenario are that the PEF is based on mechanical disturbance of the soil. Two types of mechanical soil disturbance are addressed: standard vehicle traffic and other than standard vehicle traffic (e.g. wind, grading, dozing, tilling and excavating). In general, the intakes and contact rates are all greater than the outdoor worker. Exhibit 5-1 in the [supplemental soil screening guidance](#) presents the exposure parameters.

Graphical Representation



DCC Equations

The construction worker soil land use equations, presented here, contain the following exposure routes:

- incidental ingestion of soil

$$DCC_{cw-soil-ing-sa} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{oa} \left( \frac{mrem}{pCi} \right) \times EF_{cw} \left( \frac{EW_{cw}}{50 \text{ weeks}} \times \frac{5 \text{ days}}{year} \times \frac{DW_{cw}}{5 \text{ days}} \times \frac{1 \text{ week}}{week} \right) \times IRS_{cw} \left( \frac{330 \text{ mg}}{day} \right) \times \left( \frac{g}{1000 \text{ mg}} \right)}$$

- inhalation of particulates emitted from soil

$$DCC_{cw-soil-inh-sa} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{li} \left( \frac{mrem}{pCi} \right) \times EF_{cw} \left( \frac{EW_{cw}}{50 \text{ weeks}} \times \frac{5 \text{ days}}{week} \right) \times ET_{cw} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{cw} \left( \frac{60 \text{ m}^3}{day} \right) \times \frac{1}{PEF'_{sc} \left( \frac{m^3}{kg} \right) \times \left( \frac{1000 \text{ g}}{kg} \right)}$$

- external exposure to ionizing radiation

$$DCC_{cw-soil-ext-sa} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-sv} \left( \frac{mrem}{pCi/g} \right) \times EF_{cw} \left( \frac{EW_{cw}}{50 \text{ weeks}} \times \frac{5 \text{ days}}{week} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{cw} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-sv} (1.0) \times ACF_{ext-sv}}$$

- total

$$DCC_{cw-soil-tot-sa} (pCi/g) = \frac{1}{\frac{1}{DCC_{cw-soil-ing-sa}} + \frac{1}{DCC_{cw-soil-inh-sa}} + \frac{1}{DCC_{cw-soil-ext-sa}}}$$

Definitions of the input variables are in [Table 1](#).

CDI Equations

The construction worker soil land use equations, presented here, contain the following exposure routes:

- incidental ingestion of soil

$$CDI_{cw-soil-ing-sa} \left( \frac{pCi}{year} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{cw} \left( \frac{EW_{cw}}{50 \text{ weeks}} \times \frac{5 \text{ days}}{year} \times \frac{DW_{cw}}{5 \text{ days}} \times \frac{1 \text{ week}}{week} \right) \times IRS_{cw} \left( \frac{330 \text{ mg}}{day} \right) \times \left( \frac{g}{1000 \text{ mg}} \right)$$

- inhalation of particulates emitted from soil

$$CDI_{cw-soil-inh-sa} \left( \frac{pCi}{year} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{cw} \left( \frac{EW_{cw}}{50 \text{ weeks}} \times \frac{5 \text{ days}}{week} \right) \times ET_{cw} \left( \frac{8 \text{ hours}}{day} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{cw} \left( \frac{60 \text{ m}^3}{day} \right) \times \frac{1}{PEF'_{sc} \left( \frac{m^3}{kg} \right) \times \left( \frac{1000 \text{ g}}{kg} \right)}$$

- external exposure to ionizing radiation

$$CDI_{cw-soil-ext-sa} \left( \frac{pCi}{g} \right) = \left( C_{soil} \left( \frac{pCi}{g} \right) \times EF_{cw} \left( \frac{EW_{cw}}{50 \text{ weeks}} \times \frac{5 \text{ days}}{week} \right) \times \left( \frac{1 \text{ year}}{365 \text{ day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{ext-sv} (1.0) \times ACF_{ext-sv} \right)$$

Definitions of the input variables are in [Table 1](#).

4-5.5 Construction Worker Soil Exposure to Other Construction Activities 2-D External Exposure

This assessment is the same as Unpaved Roads.

4-5.6 Construction Worker Air from Exposure to Other Construction Activities

This assessment is the same as Unpaved Roads.




## 4.6 Recreator

### 4.6.1 Recreator Soil

This receptor spends time outside involved in recreational activities. There are no default DCCs for this scenario; DCCs are only site-specific.

#### Graphical Representation



#### DCC Equations

- incidental ingestion of soil

$$DCC_{\text{rec-sol-ing}} (\text{pCi/g}) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_s \left( \frac{\text{mrem}}{\text{pCi}} \right) \times IFS_{\text{rec-adj}} \left( \frac{\text{mg}}{\text{year}} \right) \times \left( \frac{\text{g}}{1000 \text{ mg}} \right)}$$

where:

$$IFS_{\text{rec-adj}} \left( \frac{\text{mg}}{\text{year}} \right) = \left( \left( EF_{\text{rec-c}} \left( \frac{\text{days}}{\text{year}} \right) \times IRS_{\text{rec-c}} \left( \frac{200 \text{ mg}}{\text{day}} \right) \times AAF_{\text{rec-c}} \right) + \left( EF_{\text{rec-a}} \left( \frac{\text{days}}{\text{year}} \right) \times IRS_{\text{rec-a}} \left( \frac{100 \text{ mg}}{\text{day}} \right) \times AAF_{\text{rec-a}} \right) \right)$$

where:

$$AAF_{\text{rec-c}} = \left( \frac{ED_{\text{rec-c}} (\text{years})}{ED_{\text{rec}} (\text{years})} \right) \text{ and } AAF_{\text{rec-a}} = \left( \frac{ED_{\text{rec-a}} (\text{years})}{ED_{\text{rec}} (\text{years})} \right)$$

- inhalation of particulates emitted from soil

$$DCC_{\text{rec-sol-inh}} (\text{pCi/g}) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_i \left( \frac{\text{mrem}}{\text{pCi}} \right) \times IFA_{\text{rec-adj}} \left( \frac{\text{m}^3}{\text{year}} \right) \times \frac{1}{PEF \left( \frac{\text{m}^3}{\text{kg}} \right)} \times \left( \frac{1000 \text{ g}}{\text{kg}} \right)}$$

where:

$$IFA_{\text{rec-adj}} \left( \frac{\text{m}^3}{\text{year}} \right) = \left( \left( EF_{\text{rec-c}} \left( \frac{\text{days}}{\text{year}} \right) \times ET_{\text{rec-c}} \left( \frac{\text{hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{rec-c}} \left( \frac{10 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{rec-c}} \right) + \left( EF_{\text{rec-a}} \left( \frac{\text{days}}{\text{year}} \right) \times ET_{\text{rec-a}} \left( \frac{\text{hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{rec-a}} \left( \frac{20 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{rec-a}} \right) \right)$$

where:

$$AAF_{\text{rec-c}} = \left( \frac{ED_{\text{rec-c}} (\text{years})}{ED_{\text{rec}} (\text{years})} \right) \text{ and } AAF_{\text{rec-a}} = \left( \frac{ED_{\text{rec-a}} (\text{years})}{ED_{\text{rec}} (\text{years})} \right)$$

- external exposure to ionizing radiation

$$DCC_{\text{rec-sol-ext}} (\text{pCi/g}) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_{\text{ext-sv}} \left( \frac{\text{mrem/year}}{\text{pCi/g}} \right) \times EF_{\text{rec}} \left( \frac{\text{days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{\text{rec}} \left( \frac{\text{hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{ext-sv}} (1.0) \times ACF_{\text{ext-sv}}}$$

- total

$$DCC_{\text{rec-sol-tot}} (\text{pCi/g}) = \frac{1}{\frac{1}{DCC_{\text{rec-sol-ing}}} + \frac{1}{DCC_{\text{rec-sol-inh}}} + \frac{1}{DCC_{\text{rec-sol-ext}}}}$$

Definitions of the input variables are in [Table 1](#).

#### CDI Equations

- incidental ingestion of soil

$$CDI_{\text{rec-soil-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{soil}} \left( \frac{\text{pCi}}{\text{g}} \right) \times FS_{\text{rec-adj}} \left( \frac{\text{mg}}{\text{year}} \right) \times \left( \frac{\text{g}}{1000 \text{ mg}} \right)$$

where:

$$IFS_{\text{rec-adj}} \left( \frac{\text{mg}}{\text{year}} \right) = \left( \left( EF_{\text{rec-c}} \left( \frac{\text{days}}{\text{year}} \right) \times IRS_{\text{rec-c}} \left( \frac{200 \text{ mg}}{\text{day}} \right) \times AAF_{\text{rec-c}} \right) + \left( EF_{\text{rec-a}} \left( \frac{\text{days}}{\text{year}} \right) \times IRS_{\text{rec-a}} \left( \frac{100 \text{ mg}}{\text{day}} \right) \times AAF_{\text{rec-a}} \right) \right)$$

where:

$$AAF_{\text{rec-c}} = \left( \frac{ED_{\text{rec-c}} (\text{years})}{ED_{\text{rec}} (\text{years})} \right) \text{ and } AAF_{\text{rec-a}} = \left( \frac{ED_{\text{rec-a}} (\text{years})}{ED_{\text{rec}} (\text{years})} \right)$$

- inhalation of particulates emitted from soil

$$CDI_{\text{rec-soil-inh}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{soil}} \left( \frac{\text{pCi}}{\text{g}} \right) \times IFA_{\text{rec-adj}} \left( \frac{\text{m}^3}{\text{year}} \right) \times \frac{1}{PEF} \left( \frac{\text{m}^3}{\text{kg}} \right) \times \left( \frac{1000 \text{ g}}{\text{kg}} \right)$$

where:

$$IFA_{\text{rec-adj}} \left( \frac{\text{m}^3}{\text{year}} \right) = \left( \left( EF_{\text{rec-c}} \left( \frac{\text{days}}{\text{year}} \right) \times ET_{\text{rec-c}} \left( \frac{\text{hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{rec-c}} \left( \frac{10 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{rec-c}} \right) + \left( EF_{\text{rec-a}} \left( \frac{\text{days}}{\text{year}} \right) \times ET_{\text{rec-a}} \left( \frac{\text{hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{rec-a}} \left( \frac{20 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{rec-a}} \right) \right)$$

where:

$$AAF_{\text{rec-c}} = \left( \frac{ED_{\text{rec-c}} (\text{years})}{ED_{\text{rec}} (\text{years})} \right) \text{ and } AAF_{\text{rec-a}} = \left( \frac{ED_{\text{rec-a}} (\text{years})}{ED_{\text{rec}} (\text{years})} \right)$$

- external exposure to ionizing radiation

$$CDI_{\text{rec-soil-ext}} \left( \frac{\text{pCi}}{\text{g}} \right) = C_{\text{soil}} \left( \frac{\text{pCi}}{\text{g}} \right) \times EF_{\text{rec}} \left( \frac{\text{days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{\text{rec}} \left( \frac{\text{hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{ext-sv}} (1.0) \times ACF_{\text{ext-sv}}$$


Definitions of the input variables are in [Table 1](#).

### 4.6.2 Recreator Soil 2-D External Exposure

This receptor spends time outside involved in recreational activities. There are no default DCCs for this scenario; DCCs are only site-specific.

This analysis is designed to look at external exposure from contamination of different area sizes. Areas considered are 1 to 1,000,000 square meters. Isotope-specific area correction factors (ACF) were developed for this analysis.

#### Graphical Representation



#### DCC Equations

- Direct External Exposure to contamination at infinite depth

$$DCC_{\text{rec-soil-sv}} \left( \frac{\text{pCi/g}}{\text{mrem}} \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_{\text{ext-sv}} \left( \frac{\text{mrem/year}}{\text{pCi/g}} \right) \times EF_{\text{rec}} \left( \frac{\text{days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{\text{rec}} \left( \frac{\text{hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{ext-sv}} (1.0) \times ACF_{\text{ext-sv}}}$$

- Direct External Exposure to contamination 1 cm thick

$$DCC_{\text{rec-soil-1cm}} \left( \frac{\text{pCi/g}}{\text{mrem}} \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_{\text{ext-1cm}} \left( \frac{\text{mrem/year}}{\text{pCi/g}} \right) \times EF_{\text{rec}} \left( \frac{\text{days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{\text{rec}} \left( \frac{\text{hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{ext-1cm}} (1.0) \times ACF_{\text{ext-1cm}}}$$

- Direct External Exposure to contamination 5 cm thick

$$DCC_{\text{rec-soil-5cm}} \left( \frac{\text{pCi/g}}{\text{mrem}} \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_{\text{ext-5cm}} \left( \frac{\text{mrem/year}}{\text{pCi/g}} \right) \times EF_{\text{rec}} \left( \frac{\text{days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{\text{rec}} \left( \frac{\text{hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{ext-5cm}} (1.0) \times ACF_{\text{ext-5cm}}}$$

- Direct External Exposure to contamination 15 cm thick

$$DCC_{rec-soil-15cm} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-15cm} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{rec} \left( \frac{days}{year} \right) \times \left( \frac{1 year}{365 days} \right) \times ET_{rec} \left( \frac{hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times GSF_{ext-15cm} (1.0) \times ACF_{ext-15cm}}$$

- Direct External Exposure to surface contamination

$$DCC_{rec-soil-gp} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_{ext-gp} \left( \frac{mrem/year}{pCi/g} \right) \times EF_{rec} \left( \frac{days}{year} \right) \times \left( \frac{1 year}{365 days} \right) \times ET_{rec} \left( \frac{hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times GSF_{ext-gp} (1.0) \times ACF_{ext-gp}}$$

Definitions of the input variables are in [Table 1](#).

CDI Equations

- Direct External Exposure to contamination at infinite depth

$$CDI_{rec-soil-sv} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{rec} \left( \frac{days}{year} \right) \times \left( \frac{1 year}{365 days} \right) \times ET_{rec} \left( \frac{hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times GSF_{ext-sv} (1.0) \times ACF_{ext-sv}$$

- Direct External Exposure to contamination 1 cm thick

$$CDI_{rec-soil-1cm} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{rec} \left( \frac{days}{year} \right) \times \left( \frac{1 year}{365 days} \right) \times ET_{rec} \left( \frac{hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times GSF_{ext-1cm} (1.0) \times ACF_{ext-1cm}$$

- Direct External Exposure to contamination 5 cm thick

$$CDI_{rec-soil-5cm} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{rec} \left( \frac{days}{year} \right) \times \left( \frac{1 year}{365 days} \right) \times ET_{rec} \left( \frac{hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times GSF_{ext-5cm} (1.0) \times ACF_{ext-5cm}$$

- Direct External Exposure to contamination 15 cm thick

$$CDI_{rec-soil-15cm} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{rec} \left( \frac{days}{year} \right) \times \left( \frac{1 year}{365 days} \right) \times ET_{rec} \left( \frac{hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times GSF_{ext-15cm} (1.0) \times ACF_{ext-15cm}$$

- Direct External Exposure to surface contamination

$$CDI_{rec-soil-gp} \left( \frac{pCi}{g} \right) = C_{soil} \left( \frac{pCi}{g} \right) \times EF_{rec} \left( \frac{days}{year} \right) \times \left( \frac{1 year}{365 days} \right) \times ET_{rec} \left( \frac{hours}{day} \right) \times \left( \frac{1 day}{24 hours} \right) \times GSF_{ext-gp} (1.0) \times ACF_{ext-gp}$$


Definitions of the input variables are in [Table 1](#).

4.6.3 Recreator Air

This receptor spends time involved in recreational activities. There are no default DCCs for this scenario; only site-specific.

Two ambient air exposure conditions are offered for this scenario. The first scenario includes a half-life decay function and the second scenario does not. In situations where the contaminant in the air is not being replenished (e.g., an accidental one-time air release from a factory), equations for the first scenario should be used. In situations where the contaminant in the air has a continual source (e.g., indoor radon from radium in the soil or an operating factory or landfill cap), equations for the second scenario should be used.

Graphical Representation



DCC Equations

The recreator ambient air land use equation, presented here, contains the following exposure routes without half-life decay:

- inhalation (without half-life decay)

$$DCC_{\text{rec-air-inh-nodecay}} \left( \text{pCi/m}^3 \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_i \left( \frac{\text{mrem}}{\text{pCi}} \right) \times ED_{\text{reca}} (\text{years}) \times IFA_{\text{rec-adj}} \left( \frac{\text{m}^3}{\text{year}} \right)}$$

where:

$$IFA_{\text{rec-adj}} \left( \frac{\text{m}^3}{\text{year}} \right) = \left( \left( EF_{\text{rec-c}} \left( \frac{\text{days}}{\text{year}} \right) \times ET_{\text{rec-c}} \left( \frac{\text{hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{rec-c}} \left( \frac{\text{m}^3}{\text{day}} \right) \times AAF_{\text{rec-c}} \right) + \left( EF_{\text{rec-a}} \left( \frac{\text{days}}{\text{year}} \right) \times ET_{\text{rec-a}} \left( \frac{\text{hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{rec-a}} \left( \frac{\text{m}^3}{\text{day}} \right) \times AAF_{\text{rec-a}} \right) \right)$$

where:

$$AAF_{\text{rec-c}} = \left( \frac{ED_{\text{rec-c}} (\text{years})}{ED_{\text{rec}} (\text{years})} \right) \text{ and } AAF_{\text{rec-a}} = \left( \frac{ED_{\text{rec-a}} (\text{years})}{ED_{\text{rec}} (\text{years})} \right)$$

- external exposure to ionizing radiation (without half-life decay)

$$DCC_{\text{rec-air-sub-nodecay}} \left( \text{pCi/m}^3 \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_{\text{sub}} \left( \frac{\text{mrem/year}}{\text{pCi/m}^3} \right) \times EF_{\text{rec}} \left( \frac{\text{days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{\text{rec}} \left( \frac{\text{hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_a (1.0)}$$

- total

$$DCC_{\text{rec-air-tot-nodecay}} \left( \text{pCi/m}^3 \right) = \frac{1}{\frac{1}{DCC_{\text{rec-air-inh-nodecay}}} + \frac{1}{DCC_{\text{rec-air-sub-nodecay}}}}$$

Definitions of the input variables are in [Table 1](#).

### CDI Equations

The recreator ambient air land use equation, presented here, contains the following exposure routes without half-life decay:

- inhalation (without half-life decay)

$$CDI_{\text{rec-air-inh-nodecay}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{air}} \left( \frac{\text{pCi}}{\text{m}^3} \right) \times IFA_{\text{rec-adj}} \left( \frac{\text{m}^3}{\text{year}} \right)$$

where:

$$IFA_{\text{rec-adj}} \left( \frac{\text{m}^3}{\text{year}} \right) = \left( \left( EF_{\text{rec-c}} \left( \frac{\text{days}}{\text{year}} \right) \times ET_{\text{rec-c}} \left( \frac{\text{hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{rec-c}} \left( \frac{\text{m}^3}{\text{day}} \right) \times AAF_{\text{rec-c}} \right) + \left( EF_{\text{rec-a}} \left( \frac{\text{days}}{\text{year}} \right) \times ET_{\text{rec-a}} \left( \frac{\text{hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{rec-a}} \left( \frac{\text{m}^3}{\text{day}} \right) \times AAF_{\text{rec-a}} \right) \right)$$

where:

$$AAF_{\text{rec-c}} = \left( \frac{ED_{\text{rec-c}} (\text{years})}{ED_{\text{rec}} (\text{years})} \right) \text{ and } AAF_{\text{rec-a}} = \left( \frac{ED_{\text{rec-a}} (\text{years})}{ED_{\text{rec}} (\text{years})} \right)$$

- external exposure to ionizing radiation (without half-life decay)

$$CDI_{\text{rec-air-sub-nodecay}} \left( \frac{\text{pCi}}{\text{m}^3} \right) = C_{\text{air}} \left( \frac{\text{pCi}}{\text{m}^3} \right) \times EF_{\text{rec}} \left( \frac{\text{days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{\text{rec}} \left( \frac{\text{hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_a (1.0)$$

Definitions of the input variables are in [Table 1](#).

### 4.6.4 Recreator Consumption of Fowl and Land Game

This receptor spends time involved in recreational hunting of waterfowl and land game. There are no default exposure assumptions this receptor.

#### Graphical Representation



#### DCC Equations

The consumption of fowl and game equations, presented here, contains the following exposure routes:

- consumption of fowl - direct

$$DCC_{rec-fowl-ing} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_o \left( \frac{mrem}{pCi} \right) \times EF_{rec} \left( \frac{days}{year} \right) \times IRGF_{rec} \left( \frac{g}{day} \right) \times CF_{rec-fowl}(1)}$$

- consumption of fowl - back-calculated to soil

$$DCC_{soil-rec-fowl-ing} (pCi/g) = \frac{DCC_{rec-fowl-ing} (pCi/g)}{TF_{fowl} \left( \frac{day}{kg} \right) \times \left[ \left( Q_{p-fowl} \left( \frac{kg}{day} \right) \times f_{p-fowl}(1) \times f_{s-fowl}(1) \times (R_{upp} + R_{es}) \right) + \left( Q_{s-fowl} \left( \frac{kg}{day} \right) \times f_{p-fowl}(1) \right) \right]}$$

where:

$$R_{upp} = Bv_{dry} \left( \frac{pCi / g-dry plant}{pCi / g-dry soil} \right); R_{es} = MLF_{pasture} \left( \frac{0.25 g-dry soil}{g-dry plant} \right)$$

The transfer factor for fowl is the same transfer factor used for poultry.

- consumption of fowl - back-calculated to water

$$DCC_{water-rec-fowl-ing} (pCi/L) = \frac{DCC_{rec-fowl-ing} (pCi/g)}{TF_{fowl} \left( \frac{day}{kg} \right) \times Q_{w-fowl} \left( \frac{L}{day} \right) \times \left( \frac{1 kg}{1000 g} \right)}$$

- consumption of land game - direct

$$DCC_{rec-game-ing} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_o \left( \frac{mrem}{pCi} \right) \times EF_{rec} \left( \frac{days}{year} \right) \times IRGL_{rec} \left( \frac{g}{day} \right) \times CF_{rec-game}(1)}$$

- consumption of land game - back-calculated to soil

$$DCC_{soil-rec-game-ing} (pCi/g) = \frac{DCC_{rec-game-ing} (pCi/g)}{TF_{game} \left( \frac{day}{kg} \right) \times \left[ \left( Q_{p-game} \left( \frac{kg}{day} \right) \times f_{p-game}(1) \times f_{s-game}(1) \times (R_{upp} + R_{es}) \right) + \left( Q_{s-game} \left( \frac{kg}{day} \right) \times f_{p-game}(1) \right) \right]}$$

where:

$$R_{upp} = Bv_{dry} \left( \frac{pCi / g-dry plant}{pCi / g-dry soil} \right); R_{es} = MLF_{pasture} \left( \frac{0.25 g-dry soil}{g-dry plant} \right)$$

The transfer factor for game is the same transfer factor used for beef.

- consumption of land game - back-calculated to water.

$$DCC_{water-rec-game-ing} (pCi/L) = \frac{DCC_{rec-game-ing} (pCi/g)}{TF_{game} \left( \frac{day}{kg} \right) \times Q_{w-game} \left( \frac{L}{day} \right) \times \left( \frac{1 kg}{1000 g} \right)}$$

Definitions of the input variables are in [Table 1](#).

### CDI Equations

The consumption of fowl and game equations, presented here, contains the following exposure routes:

- consumption of fowl - direct

$$CDI_{rec-fowl-ing} \left( \frac{pCi}{year} \right) = C_{fowl} \left( \frac{pCi}{g} \right) \times EF_{rec} \left( \frac{days}{year} \right) \times IRGF_{rec} \left( \frac{g}{day} \right) \times CF_{rec-fowl}(1)$$

- consumption of fowl - back-calculated to soil

$$CDI_{soil-rec-fowl-ing} \left( \frac{pCi}{year} \right) = \left[ \frac{C_{soil} \left( \frac{pCi}{g} \right) \times EF_{rec} \left( \frac{days}{year} \right) \times IRGF_{rec} \left( \frac{g}{day} \right) \times CF_{rec-fowl}(1) \times TF_{fowl} \left( \frac{day}{kg} \right) \times \left[ \left( Q_{p-fowl} \left( \frac{kg}{day} \right) \times f_{p-fowl}(1) \times f_{s-fowl}(1) \times (R_{upp} + R_{es}) \right) + \left( Q_{s-fowl} \left( \frac{kg}{day} \right) \times f_{p-fowl}(1) \right) \right]}{TF_{fowl} \left( \frac{day}{kg} \right) \times \left[ \left( Q_{p-fowl} \left( \frac{kg}{day} \right) \times f_{p-fowl}(1) \times f_{s-fowl}(1) \times (R_{upp} + R_{es}) \right) + \left( Q_{s-fowl} \left( \frac{kg}{day} \right) \times f_{p-fowl}(1) \right) \right]} \right]$$

where:

$$R_{upp} = Bv_{dry} \left( \frac{pCi / g-dry plant}{pCi / g-dry soil} \right); R_{es} = MLF_{pasture} \left( \frac{0.25 g-dry soil}{g-dry plant} \right)$$

The transfer factor for fowl is the same transfer factor used for poultry.

- consumption of fowl - back-calculated to water

$$CDI_{water-rec-fowl-ing} \left( \frac{pCi}{year} \right) = \left[ \frac{C_{water} \left( \frac{pCi}{L} \right) \times EF_{rec} \left( \frac{days}{year} \right) \times IRGF_{rec} \left( \frac{g}{day} \right) \times CF_{rec-fowl}(1) \times TF_{fowl} \left( \frac{day}{kg} \right) \times Q_{w-fowl} \left( \frac{L}{day} \right) \times \left( \frac{1 kg}{1000 g} \right)}{TF_{fowl} \left( \frac{day}{kg} \right) \times Q_{w-fowl} \left( \frac{L}{day} \right) \times \left( \frac{1 kg}{1000 g} \right)} \right]$$

- consumption of land game - direct

$$CDI_{rec-game-ing} \left( \frac{pCi}{year} \right) = C_{fowl} \left( \frac{pCi}{g} \right) \times EF_{rec} \left( \frac{days}{year} \right) \times IRGL_{rec} \left( \frac{g}{day} \right) \times CF_{rec-game}(1)$$

- consumption of land game - back-calculated to soil

$$CDI_{\text{soil-rec-game-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = \left[ \begin{aligned} & C_{\text{soil}} \left( \frac{\text{pCi}}{\text{g}} \right) \times EF_{\text{rec}} \left( \frac{\text{days}}{\text{year}} \right) \times IRGF_{\text{rec}} \left( \frac{\text{g}}{\text{day}} \right) \times CF_{\text{rec-game}}(1) \times TF_{\text{game}} \left( \frac{\text{day}}{\text{kg}} \right) \times \\ & \left[ \begin{aligned} & \left( Q_{\text{p-game}} \left( \frac{\text{kg}}{\text{day}} \right) \times f_{\text{p-game}}(1) \times f_{\text{s-game}}(1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \\ & \left( Q_{\text{s-game}} \left( \frac{\text{kg}}{\text{day}} \right) \times f_{\text{p-game}}(1) \right) \end{aligned} \right] \end{aligned} \right]$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{\text{pCi / g-dry plant}}{\text{pCi / g-dry soil}} \right); R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{\text{g-dry plant}} \right)$$

The transfer factor for game is the same transfer factor used for beef.

- consumption of land game - back-calculated to water.


$$CDI_{\text{water-rec-game-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = \left[ \begin{aligned} & C_{\text{water}} \left( \frac{\text{pCi}}{\text{L}} \right) \times EF_{\text{rec}} \left( \frac{\text{days}}{\text{year}} \right) \times IRGF_{\text{rec}} \left( \frac{\text{g}}{\text{day}} \right) \times CF_{\text{rec-game}}(1) \times \\ & TF_{\text{game}} \left( \frac{\text{day}}{\text{kg}} \right) \times Q_{\text{w-game}} \left( \frac{\text{L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right) \end{aligned} \right]$$

Definitions of the input variables are in [Table 1](#).

#### 4.6.5 Recreator Surface Water

This receptor is exposed to radionuclides that are present in surface water. Ingestion of water and immersion in water are appropriate pathways for all radionuclides. Inhalation is not considered due to mixing with outdoor air. There are no default DCCs for this scenario; DCCs are only site-specific.

##### Graphical Representation



##### DCC Equations

- ingestion of surface water

$$DCC_{\text{rec-water-ing}} \left( \frac{\text{pCi/L}}{\text{year}} \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_o \left( \frac{\text{mrem}}{\text{pCi}} \right) \times IFW_{\text{rec-adj}} \left( \frac{\text{L}}{\text{year}} \right)}$$

where:

$$IFW_{\text{rec-adj}} \left( \frac{\text{L}}{\text{year}} \right) = \left[ \begin{aligned} & \left( EF_{\text{rec-c}} \left( \frac{\text{days}}{\text{year}} \right) \times ET_{\text{event-rec-c}} \left( \frac{\text{hours}}{\text{event}} \right) \times EV_{\text{rec-c}} \left( \frac{\text{events}}{\text{day}} \right) \times IRW_{\text{rec-c}} \left( \frac{0.12 \text{ L}}{\text{hour}} \right) \times AAF_{\text{rec-c}} \right) + \\ & \left( EF_{\text{rec-a}} \left( \frac{\text{days}}{\text{year}} \right) \times ET_{\text{event-rec-a}} \left( \frac{\text{hours}}{\text{event}} \right) \times EV_{\text{rec-a}} \left( \frac{\text{events}}{\text{day}} \right) \times IRW_{\text{rec-a}} \left( \frac{0.071 \text{ L}}{\text{hour}} \right) \times AAF_{\text{rec-a}} \right) \end{aligned} \right]$$

where:

$$AAF_{\text{rec-c}} = \left( \frac{ED_{\text{rec-c}}(\text{years})}{ED_{\text{rec}}(\text{years})} \right) \text{ and } AAF_{\text{rec-a}} = \left( \frac{ED_{\text{rec-a}}(\text{years})}{ED_{\text{rec}}(\text{years})} \right)$$

- immersion

$$DCC_{\text{rec-water-imm}} \left( \frac{\text{pCi/L}}{\text{year}} \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_{\text{imm}} \left( \frac{\text{mrem/year}}{\text{pCi/L}} \right) \times \left( \frac{1 \text{ year}}{8760 \text{ hours}} \right) \times DFA_{\text{rec-adj}} \left( \frac{\text{hours}}{\text{year}} \right)}$$

where:

$$DFA_{\text{rec-adj}} \left( \frac{\text{hours}}{\text{year}} \right) = \left[ \begin{aligned} & \left( EF_{\text{rec-c}} \left( \frac{\text{days}}{\text{year}} \right) \times EV_{\text{rec-c}} \left( \frac{\text{event}}{\text{day}} \right) \times ET_{\text{event-rec-c}} \left( \frac{\text{hours}}{\text{event}} \right) \times AAF_{\text{rec-c}} \right) + \\ & \left( EF_{\text{rec-a}} \left( \frac{\text{days}}{\text{year}} \right) \times EV_{\text{rec-a}} \left( \frac{\text{event}}{\text{day}} \right) \times ET_{\text{event-rec-a}} \left( \frac{\text{hours}}{\text{event}} \right) \times AAF_{\text{rec-a}} \right) \end{aligned} \right]$$

where:

$$AAF_{\text{rec-c}} = \left( \frac{ED_{\text{rec-c}}(\text{years})}{ED_{\text{rec}}(\text{years})} \right) \text{ and } AAF_{\text{rec-a}} = \left( \frac{ED_{\text{rec-a}}(\text{years})}{ED_{\text{rec}}(\text{years})} \right)$$

- total



$$DCC_{rec-water-tot} (pCi/L) = \frac{1}{\frac{1}{DCC_{rec-water-ing}} + \frac{1}{DCC_{rec-water-imm}}}$$

Definitions of the input variables are in [Table 1](#).

CDI Equations

- ingestion of surface water

$$CDI_{water-rec-ing} \left( \frac{pCi}{year} \right) = C_{water} \left( \frac{pCi}{L} \right) \times IFW_{rec-adj} \left( \frac{L}{year} \right)$$

where:

$$IFW_{rec-adj} \left( \frac{L}{year} \right) = \left( \left( EF_{rec-c} \left( \frac{days}{year} \right) \times ET_{event-rec-c} \left( \frac{hours}{event} \right) \times EV_{rec-c} \left( \frac{events}{day} \right) \times IRW_{rec-c} \left( \frac{0.12 L}{hour} \right) \times AAF_{rec-c} \right) + \left( EF_{rec-a} \left( \frac{days}{year} \right) \times ET_{event-rec-a} \left( \frac{hours}{event} \right) \times EV_{rec-a} \left( \frac{events}{day} \right) \times IRW_{rec-a} \left( \frac{0.071 L}{hour} \right) \times AAF_{rec-a} \right) \right)$$

where:

$$AAF_{rec-c} = \left( \frac{ED_{rec-c} (years)}{ED_{rec} (years)} \right) \text{ and } AAF_{rec-a} = \left( \frac{ED_{rec-a} (years)}{ED_{rec} (years)} \right)$$

- immersion

$$CDI_{water-rec-imm} \left( \frac{pCi}{L} \right) = C_{water} \left( \frac{pCi}{L} \right) \times \left( \frac{1 \text{ year}}{8760 \text{ hours}} \right) \times DFA_{rec-adj} \left( \frac{hours}{year} \right)$$

where:

$$DFA_{rec-adj} \left( \frac{hours}{year} \right) = \left( \left( EF_{rec-c} \left( \frac{days}{year} \right) \times EV_{rec-c} \left( \frac{event}{day} \right) \times ET_{event-rec-c} \left( \frac{hours}{event} \right) \times AAF_{rec-c} \right) + \left( EF_{rec-a} \left( \frac{days}{year} \right) \times EV_{rec-a} \left( \frac{event}{day} \right) \times ET_{event-rec-a} \left( \frac{hours}{event} \right) \times AAF_{rec-a} \right) \right)$$

where:


$$AAF_{rec-c} = \left( \frac{ED_{rec-c} (years)}{ED_{rec} (years)} \right) \text{ and } AAF_{rec-a} = \left( \frac{ED_{rec-a} (years)}{ED_{rec} (years)} \right)$$

Definitions of the input variables are in [Table 1](#).

4.7 Consumption of Fish

The fish DCC represents the concentration in the fish that can be consumed. This is unlike the farmer scenario, where the DCC is calculated for soil levels protective of fish consumption. Further, the ingestion rate is not age-adjusted like the farmer scenario.

Graphical Representation



DCC Equation

The consumption of fish equation, presented here, contains the following exposure route:

- consumption of fish

$$DCC_{res-fish-ing} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_o \left( \frac{mrem}{pCi} \right) \times EF_{res} \left( \frac{350 \text{ days}}{year} \right) \times IRFI_{res-a} \left( \frac{54,000 \text{ mg}}{day} \right) \times \left( \frac{10^{-3} \text{ g}}{1 \text{ mg}} \right) \times CF_{res-fish} (1)}$$

**Note:** the consumption rate for fish is not age-adjusted for this land use. Also, the DCC is calculated for fish tissue rather than for soil, like for the farmer land uses.

Definitions of the input variables are in [Table 1](#).

CDI Equation

The consumption of fish equation, presented here, contains the following exposure route:

- consumption of fish

$$CDI_{res-fish-ing} \left( \frac{pCi}{year} \right) = C_{fish} \left( \frac{pCi}{g} \right) \times EF_{res} \left( \frac{350 \text{ days}}{year} \right) \times IRF_{res-a} \left( \frac{54,000 \text{ mg}}{day} \right) \times \left( \frac{g}{1000 \text{ mg}} \right) \times CF_{res-fish} (1)$$

**Note: the consumption rate for fish is not age-adjusted for this land use. Also, the DCC is calculated for fish tissue rather than for soil, like for the farmer land uses.**

Definitions of the input variables are in [Table 1](#).


## 4.8 Farmer

### 4.8.1 Farmer Direct Consumption of Agricultural Products

The farmer scenario should be considered an extension of the resident scenario and evaluate consumption of farm products for a subsistence farmer. Like the resident, the farmer assumes the receptor will be exposed via the consumption of home grown produce (100% of fruit and vegetables are from the farm). In addition to produce, 100% of consumption of the following are also considered to be from the farm: beef, milk, fish, swine, egg, and poultry. All feed (100%) for farm products is considered to have been grown on contaminated portions of the site. For these farm products, dose-based DCCs are provided for the farm product itself (vegetables, beef, milk, etc.). Also like the resident, age-adjusted intake equations were developed for all of the consumption equations to account for changes in intake as the receptor ages.

#### Graphical Representation

### Agricultural Biota, Soil and Water Graphic and Supporting Text



#### DCC Equations

- consumption of produce (fruits and vegetables). Sections 9 and 13 of the 2011 Exposure Factors Handbook were used to derive the intakes for home-grown produce.

$$DCC_{far-produce-ing-tot} (pCi/g) = \frac{1}{\left( \sum_{i=1}^n \frac{1}{DCC_{far-produce-ing} (pCi/g)} \right)}$$

where:  
n = total number of produce items included

and:

$$DCC_{far-produce-ing} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_o \left( \frac{mrem}{pCi} \right) \times IF_{far-adj} \left( \frac{g}{year} \right) \times CF_{far-produce} (1)}$$

where:

$$IF_{far-adj} \left( \frac{g}{year} \right) = \left( \left( EF_{far-c} \left( \frac{350 \text{ days}}{year} \right) \times IR_{far-c} \left( \frac{g}{day} \right) \times AAF_{far-c} (0.15) \right) + \left( EF_{far-a} \left( \frac{350 \text{ days}}{year} \right) \times IR_{far-a} \left( \frac{g}{day} \right) \times AAF_{far-a} (0.85) \right) \right)$$

and:

$$AAF_{far-c} (0.15) = \left( \frac{ED_{far-c} (6 \text{ years})}{ED_{far} (40 \text{ years})} \right) \text{ and: } AAF_{far-a} (0.85) = \left( \frac{ED_{far-a} (34 \text{ years})}{ED_{far} (40 \text{ years})} \right)$$

- consumption of poultry. Table 13-52 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced poultry.

$$DCC_{\text{far-poultry-ing}} \left( \frac{\text{pCi/g}}{\text{year}} \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_o \left( \frac{\text{mrem}}{\text{pCi}} \right) \times IFP_{\text{far-adj}} \left( \frac{34,414 \text{ g}}{\text{year}} \right) \times CF_{\text{far-poultry}} (1)}$$

where:

$$IFP_{\text{far-adj}} \left( \frac{34,414 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRP_{\text{far-c}} \left( \frac{46.9 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRP_{\text{far-a}} \left( \frac{107.4 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

• **consumption of eggs.** Table 13-40 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced eggs.

$$DCC_{\text{far-egg-ing}} \left( \frac{\text{pCi/g}}{\text{year}} \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_o \left( \frac{\text{mrem}}{\text{pCi}} \right) \times IFE_{\text{far-adj}} \left( \frac{19,395 \text{ g}}{\text{year}} \right) \times CF_{\text{far-egg}} (1)}$$

where:

$$IFE_{\text{far-adj}} \left( \frac{19,395 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRE_{\text{far-c}} \left( \frac{31.7 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRE_{\text{far-a}} \left( \frac{59.6 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

• **consumption of beef.** Table 13-33 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced beef.

$$DCC_{\text{far-beef-ing}} \left( \frac{\text{pCi/g}}{\text{year}} \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_o \left( \frac{\text{mrem}}{\text{pCi}} \right) \times IFB_{\text{far-adj}} \left( \frac{52,474 \text{ g}}{\text{year}} \right) \times CF_{\text{far-beef}} (1)}$$

where:

$$IFB_{\text{far-adj}} \left( \frac{52,474 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRB_{\text{far-c}} \left( \frac{62.8 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRB_{\text{far-a}} \left( \frac{165.3 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

• **consumption of milk.** Table 11-4 of the 2011 Exposure Factors Handbook was used to derive the intakes for total dairy.

$$DCC_{\text{far-dairy-ing}} \left( \frac{\text{pCi/g}}{\text{year}} \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_o \left( \frac{\text{risk}}{\text{pCi}} \right) \times IFD_{\text{far-adj}} \left( \frac{253,451 \text{ g}}{\text{year}} \right) \times CF_{\text{far-dairy}} (1)}$$

where:

$$IFD_{\text{far-adj}} \left( \frac{253,451 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRD_{\text{far-c}} \left( \frac{994.7 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRD_{\text{far-a}} \left( \frac{676.4 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

• **consumption of swine.** Table 13-51 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced swine.

$$DCC_{\text{far-swine-ing}} \left( \frac{\text{pCi/g}}{\text{year}} \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_o \left( \frac{\text{mrem}}{\text{pCi}} \right) \times IFSW_{\text{far-adj}} \left( \frac{29,288 \text{ g}}{\text{year}} \right) \times CF_{\text{far-swine}} (1)}$$

where:

$$IFSW_{\text{far-adj}} \left( \frac{29,288 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSW_{\text{far-c}} \left( \frac{33.7 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSW_{\text{far-a}} \left( \frac{92.5 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

• **consumption of fish.** Table 13-20 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-caught fish.

$$DCC_{\text{far-fish-ing}} \left( \frac{\text{pCi/g}}{\text{year}} \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_o \left( \frac{\text{mrem}}{\text{pCi}} \right) \times IFFI_{\text{far-adj}} \left( \frac{250,474 \text{ g}}{\text{year}} \right) \times CF_{\text{far-fish}} (1)}$$

where:

$$IFFI_{\text{far-adj}} \left( \frac{250,474 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRFI_{\text{far-c}} \left( \frac{57.4 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRFI_{\text{far-a}} \left( \frac{831.8 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

• **consumption of goat.** There are no widely accepted human intakes for home-produced goat.

$$DCC_{\text{far-goat-ing}} \left( \frac{\text{pCi/g}}{\text{year}} \right) = \frac{TR}{SF_f \left( \frac{\text{risk}}{\text{pCi}} \right) \times IFGO_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) \times CF_{\text{far-goat}} (1)}$$

where:

$$IFGO_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRGO_{\text{far-c}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRGO_{\text{far-a}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- **consumption of goat milk.** There are no widely accepted human intakes for home-produced goat dairy.

$$DCC_{\text{far-goat-milk-ing}} \left( \frac{\text{pCi/g}}{\text{year}} \right) = \frac{TR}{SF_f \left( \frac{\text{risk}}{\text{pCi}} \right) \times IFGM_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) \times CF_{\text{far-goat-milk}} (1)}$$

where:

$$IFGM_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRGM_{\text{far-c}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRGM_{\text{far-a}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- **consumption of sheep.** There are no widely accepted human intakes for home-produced sheep.

$$DCC_{\text{far-sheep-ing}} \left( \frac{\text{pCi/g}}{\text{year}} \right) = \frac{TR}{SF_f \left( \frac{\text{risk}}{\text{pCi}} \right) \times IFSH_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) \times CF_{\text{far-sheep}} (1)}$$

where:

$$IFSH_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSH_{\text{far-c}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSH_{\text{far-a}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- **consumption of sheep milk.** There are no widely accepted human intakes for home-produced sheep dairy.

$$DCC_{\text{far-sheep-milk-ing}} \left( \frac{\text{pCi/g}}{\text{year}} \right) = \frac{TR}{SF_f \left( \frac{\text{risk}}{\text{pCi}} \right) \times IFSM_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) \times CF_{\text{far-sheep-milk}} (1)}$$

where:

$$IFSM_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSM_{\text{far-c}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSM_{\text{far-a}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

Definitions of the input variables are in [Table 1](#).

### CDI Equations

- **consumption of produce (fruits and vegetables).** Sections 9 and 13 of the 2011 Exposure Factors Handbook were used to derive the intakes for home-grown produce.

$$CDI_{\text{far-produce-ing-tot}} \left( \frac{\text{pCi}}{\text{year}} \right) = \sum_{i=1}^n CDI_{\text{far-produce-ing}} \left( \frac{\text{pCi}}{\text{year}} \right)_i$$

where:

n = total number of produce items included

and:

$$CDI_{\text{far-produce-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{produce}} \left( \frac{\text{pCi}}{\text{g}} \right) \times IF_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) \times CF_{\text{far-produce}} (1)$$

where:

$$IF_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IR_{\text{far-c}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IR_{\text{far-a}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

and:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- **consumption of poultry.** Table 13-52 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced poultry.

$$CDI_{\text{far-poultry-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{poultry}} \left( \frac{\text{pCi}}{\text{g}} \right) \times IFP_{\text{far-adj}} \left( \frac{34,414 \text{ g}}{\text{year}} \right) \times CF_{\text{far-poultry}} (1)$$

where:

$$IFP_{\text{far-adj}} \left( \frac{34,414 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRP_{\text{far-c}} \left( \frac{46.9 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRP_{\text{far-a}} \left( \frac{107.4 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- **consumption of eggs.** Table 13-40 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced eggs.

$$CDI_{\text{far-egg-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{egg}} \left( \frac{\text{pCi}}{\text{g}} \right) \times IFE_{\text{far-adj}} \left( \frac{19,395 \text{ g}}{\text{year}} \right) \times CF_{\text{far-egg}} (1)$$

where:

$$IFE_{\text{far-adj}} \left( \frac{19,395 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRE_{\text{far-c}} \left( \frac{31.7 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRE_{\text{far-a}} \left( \frac{59.6 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- **consumption of beef.** Table 13-33 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced beef.

$$CDI_{\text{far-beef-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{beef}} \left( \frac{\text{pCi}}{\text{g}} \right) \times IFB_{\text{far-adj}} \left( \frac{52,474 \text{ g}}{\text{year}} \right) \times CF_{\text{far-beef}} (1)$$

where:

$$IFB_{\text{far-adj}} \left( \frac{52,474 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRB_{\text{far-c}} \left( \frac{62.8 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRB_{\text{far-a}} \left( \frac{165.3 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- **consumption of milk.** Table 11-4 of the 2011 Exposure Factors Handbook was used to derive the intakes for total dairy.

$$CDI_{\text{far-dairy-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{dairy}} \left( \frac{\text{pCi}}{\text{g}} \right) \times IFD_{\text{far-adj}} \left( \frac{253,451 \text{ g}}{\text{year}} \right) \times CF_{\text{far-dairy}} (1)$$

where:

$$IFD_{\text{far-adj}} \left( \frac{253,451 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRD_{\text{far-c}} \left( \frac{994.7 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRD_{\text{far-a}} \left( \frac{676.4 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- **consumption of swine.** Table 13-51 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced swine.

$$CDI_{\text{far-swine-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{swine}} \left( \frac{\text{pCi}}{\text{g}} \right) \times IFSW_{\text{far-adj}} \left( \frac{29,288 \text{ g}}{\text{year}} \right) \times CF_{\text{far-swine}} (1)$$

where:

$$IFSW_{\text{far-adj}} \left( \frac{29,288 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSW_{\text{far-c}} \left( \frac{33.7 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSW_{\text{far-a}} \left( \frac{92.5 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- **consumption of fish.** Table 13-20 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-caught fish.

$$CDI_{\text{far-fish-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{fish}} \left( \frac{\text{pCi}}{\text{g}} \right) \times IFFI_{\text{far-adj}} \left( \frac{250,474 \text{ g}}{\text{year}} \right) \times CF_{\text{far-fish}} (1)$$

where:

$$IFFI_{\text{far-adj}} \left( \frac{250,474 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRFI_{\text{far-c}} \left( \frac{57.4 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRFI_{\text{far-a}} \left( \frac{831.8 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- **consumption of goat.** There are no widely accepted human intakes for home-produced goat.

$$CDI_{\text{far-goat-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{goat}} \left( \frac{\text{pCi}}{\text{g}} \right) \times IFGO_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) \times CF_{\text{far-goat}} \quad (1)$$

where:

$$IFGO_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRGO_{\text{far-c}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-c}}(0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRGO_{\text{far-a}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-a}}(0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}}(0.15) = \left( \frac{ED_{\text{far-c}}(6 \text{ years})}{ED_{\text{far}}(40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}}(0.85) = \left( \frac{ED_{\text{far-a}}(34 \text{ years})}{ED_{\text{far}}(40 \text{ years})} \right)$$

- consumption of goat milk. There are no widely accepted human intakes for home-produced goat dairy.

$$CDI_{\text{far-goat-milk-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{goat-milk}} \left( \frac{\text{pCi}}{\text{g}} \right) \times IFGM_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) \times CF_{\text{far-goat-milk}} \quad (1)$$

where:

$$IFGM_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRGM_{\text{far-c}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-c}}(0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRGM_{\text{far-a}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-a}}(0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}}(0.15) = \left( \frac{ED_{\text{far-c}}(6 \text{ years})}{ED_{\text{far}}(40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}}(0.85) = \left( \frac{ED_{\text{far-a}}(34 \text{ years})}{ED_{\text{far}}(40 \text{ years})} \right)$$

- consumption of sheep. There are no widely accepted human intakes for home-produced sheep.

$$CDI_{\text{far-sheep-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{sheep}} \left( \frac{\text{pCi}}{\text{g}} \right) \times IFSH_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) \times CF_{\text{far-sheep}} \quad (1)$$

where:

$$IFSH_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSH_{\text{far-c}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-c}}(0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSH_{\text{far-a}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-a}}(0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}}(0.15) = \left( \frac{ED_{\text{far-c}}(6 \text{ years})}{ED_{\text{far}}(40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}}(0.85) = \left( \frac{ED_{\text{far-a}}(34 \text{ years})}{ED_{\text{far}}(40 \text{ years})} \right)$$

- consumption of sheep milk. There are no widely accepted human intakes for home-produced sheep dairy.

$$CDI_{\text{far-sheep-milk-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{sheep-milk}} \left( \frac{\text{pCi}}{\text{g}} \right) \times IFSM_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) \times CF_{\text{far-sheep-milk}} \quad (1)$$

where:

$$IFSM_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSM_{\text{far-c}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-c}}(0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSM_{\text{far-a}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-a}}(0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}}(0.15) = \left( \frac{ED_{\text{far-c}}(6 \text{ years})}{ED_{\text{far}}(40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}}(0.85) = \left( \frac{ED_{\text{far-a}}(34 \text{ years})}{ED_{\text{far}}(40 \text{ years})} \right)$$


Definitions of the input variables are in [Table 1](#).

#### 4.8.2 Farmer Direct Exposure and Consumption of Agricultural Products - Back-Calculated to Soil

The farmer scenario should be considered an extension of the resident scenario and evaluate consumption of farm products for a subsistence farmer. Like the resident, the farmer scenario assumes the receptor will be exposed via the following pathways: incidental ingestion of soil, external radiation from contaminants in soil, inhalation of fugitive dust, and consumption of home grown produce (100% of fruit and vegetables are from the farm). In addition to produce, 100% of consumption of the following are also considered to be from the farm: beef, milk, fish, swine, egg, and poultry. All feed (100%) for farm products is considered to have been grown on contaminated portions of the site. For these farm products, dose-based DCCs are provided for soil that may contribute contaminants to the products. Also like the resident, age-adjusted intake equations were developed for all of the ingestion/consumption equations to account for changes in intake as the receptor ages.

##### Graphical Representation





### DCC Equations

- incidental ingestion of soil

$$DCC_{\text{far-soil-ing}} (\text{pCi/g}) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_o \left( \frac{\text{mrem}}{\text{pCi}} \right) \times IFS_{\text{far-adj}} \left( \frac{40,250 \text{ mg}}{\text{year}} \right) \times \left( \frac{\text{g}}{1000 \text{ g}} \right)}$$

where:

$$IFS_{\text{far-adj}} \left( \frac{40,250 \text{ mg}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRS_{\text{far-c}} \left( \frac{200 \text{ mg}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRS_{\text{far-a}} \left( \frac{100 \text{ mg}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- inhalation of particulates emitted from soil

$$DCC_{\text{far-soil-inh}} (\text{pCi/g}) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_i \left( \frac{\text{mrem}}{\text{pCi}} \right) \times IFA_{\text{far-adj}} \left( \frac{6,475 \text{ m}^3}{\text{year}} \right) \times \frac{1}{PEF \left( \frac{\text{m}^3}{\text{kg}} \right)} \times \left( \frac{1000 \text{ g}}{\text{kg}} \right)}$$

where:

$$IFA_{\text{far-adj}} \left( \frac{6,475 \text{ m}^3}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times ET_{\text{far-c}} \left( \frac{24 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{far-c}} \left( \frac{10 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times ET_{\text{far-a}} \left( \frac{24 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{far-a}} \left( \frac{20 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- external exposure to ionizing radiation

$$DCC_{\text{far-soil-ext}} (\text{pCi/g}) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_{\text{ext-sv}} \left( \frac{\text{mrem/year}}{\text{pCi/g}} \right) \times EF_{\text{far}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ACF_{\text{ext-sv}} \times \left[ \left( ET_{\text{far-o}} \left( \frac{12,168 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{ext-sv}} (1.0) \right) + \left( ET_{\text{far-i}} \left( \frac{10,008 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{i-total}} \right) \right]}$$

- consumption of produce (fruits and vegetables). Sections 9 and 13 of the 2011 Exposure Factors Handbook were used to derive the intakes for home-grown produce.

$$DCC_{\text{soil-far-produce-ing-tot}} (\text{pCi/g}) = \frac{1}{\sum_{i=1}^n \frac{1}{DCC_{\text{soil-far-produce-ing}} (\text{pCi/g})_i}}$$

where:

n = total number of produce items included

and:

$$DCC_{\text{soil-far-produce-ing}} (\text{pCi/g}) = \frac{DCC_{\text{far-produce-ing}} (\text{pCi/g})}{(R_{\text{upv}} + R_{\text{es}})}$$

where:

$$R_{\text{upv}} = B_{\text{wet}} \left( \frac{\text{pCi / g-fresh plant}}{\text{pCi / g-dry soil}} \right); R_{\text{es}} = MLF_{\text{produce}} \left( \frac{\text{g-dry soil}}{\text{g-fresh plant}} \right)$$

The consumption of produce exposure route drives the DCCs lower than all the other routes. It is recommended that

produce-specific transfer factors ( $Bv_{wet}$ ) be used when available for a site. Further, the default transfer factors ( $Bv_{wet}$ ) from IAEA used in these DCC calculations are based on a composite of all soil groups. Transfer factors ( $Bv_{wet}$ ) for sand, loam, clay, organic, coral sand, and other soil types that may be more suited to a particular site are also provided. The site-specific option of the calculator can be used to focus on ingestion of individual produce types. When "Site-specific" is selected, if the user changes the "Select Isotope Info Type" to "User-provided", then a specific transfer factor may be changed.

where:

$$DCC_{far-produce-ing-tot} (pCi/g) = \frac{1}{\left( \sum_{i=1}^n \frac{1}{DCC_{far-produce-ing} (pCi/g)_i} \right)}$$

where:  
n = total number of produce items included

and:

$$DCC_{far-produce-ing} (pCi/g) = \frac{DL \left( \frac{mrem}{year} \right)}{DCF_o \left( \frac{mrem}{pCi} \right) \times IF_{far-adj} \left( \frac{g}{year} \right) \times CF_{far-produce} (1)}$$

where:

$$IF_{far-adj} \left( \frac{g}{year} \right) = \left( \left( EF_{far-c} \left( \frac{350 \text{ days}}{year} \right) \times IR_{far-c} \left( \frac{g}{day} \right) \times AAF_{far-c} (0.15) \right) + \left( EF_{far-a} \left( \frac{350 \text{ days}}{year} \right) \times IR_{far-a} \left( \frac{g}{day} \right) \times AAF_{far-a} (0.85) \right) \right)$$

and:

$$AAF_{far-c} (0.15) = \left( \frac{ED_{far-c} (6 \text{ years})}{ED_{far} (40 \text{ years})} \right) \text{ and: } AAF_{far-a} (0.85) = \left( \frac{ED_{far-a} (34 \text{ years})}{ED_{far} (40 \text{ years})} \right)$$

- **consumption of eggs.** Table 13-40 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced eggs.

$$DCC_{soil-far-egg-ing} (pCi/g) = \frac{DCC_{far-egg-ing} (pCi/g)}{TF_{egg} \left( \frac{day}{kg} \right) \times \left[ \left( Q_{p-poultry} \left( \frac{0.2 \text{ kg}}{day} \right) \times f_{p-poultry} (1) \times f_{s-poultry} (1) \times (R_{upp} + R_{es}) \right) + \left( Q_{s-poultry} \left( \frac{0.022 \text{ kg}}{day} \right) \times f_{p-poultry} (1) \right) \right]}$$

where:

$$R_{upp} = Bv_{dry} \left( \frac{pCi/g-dry \text{ plant}}{pCi/g-dry \text{ soil}} \right); R_{es} = MLF_{pasture} \left( \frac{0.25 \text{ g-dry soil}}{g-dry \text{ plant}} \right)$$

- **consumption of poultry.** Table 13-52 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced poultry.

$$DCC_{soil-far-poultry-ing} (pCi/g) = \frac{DCC_{far-poultry-ing} (pCi/g)}{TF_{poultry} \left( \frac{day}{kg} \right) \times \left[ \left( Q_{p-poultry} \left( \frac{0.2 \text{ kg}}{day} \right) \times f_{p-poultry} (1) \times f_{s-poultry} (1) \times (R_{upp} + R_{es}) \right) + \left( Q_{s-poultry} \left( \frac{0.022 \text{ kg}}{day} \right) \times f_{p-poultry} (1) \right) \right]}$$

where:

$$R_{upp} = Bv_{dry} \left( \frac{pCi/g-dry \text{ plant}}{pCi/g-dry \text{ soil}} \right); R_{es} = MLF_{pasture} \left( \frac{0.25 \text{ g-dry soil}}{g-dry \text{ plant}} \right)$$

- **consumption of fish.** Table 13-20 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-caught fish.

$$DCC_{soil-far-fish-ing} (pCi/g) = \frac{DCC_{far-fish-ing} (pCi/g) \times Kd \left( \frac{L}{kg} \right)}{BCF \left( \frac{L}{kg} \right)}$$

- **consumption of beef.** Table 13-33 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced beef.

$$DCC_{soil-far-beef-ing} (pCi/g) = \frac{DCC_{far-beef-ing} (pCi/g)}{TF_{beef} \left( \frac{day}{kg} \right) \times \left[ \left( Q_{p-beef} \left( \frac{11.77 \text{ kg}}{day} \right) \times f_{p-beef} (1) \times f_{s-beef} (1) \times (R_{upp} + R_{es}) \right) + \left( Q_{s-beef} \left( \frac{0.5 \text{ kg}}{day} \right) \times f_{p-beef} (1) \right) \right]}$$

where:

$$R_{upp} = Bv_{dry} \left( \frac{pCi/g-dry \text{ plant}}{pCi/g-dry \text{ soil}} \right); R_{es} = MLF_{pasture} \left( \frac{0.25 \text{ g-dry soil}}{g-dry \text{ plant}} \right)$$

- **consumption of milk.** Table 11-4 of the 2011 Exposure Factors Handbook was used to derive the intakes for total dairy.

$$DCC_{soil-far-dairy-ing} (pCi/g) = \frac{DCC_{far-dairy-ing} (pCi/g)}{TF_{dairy} \left( \frac{day}{L \text{ milk}} \right) \times \rho_m \left( \frac{1.03 \text{ kg}}{L \text{ milk}} \right)^{-1} \times \left[ \left( Q_{p-dairy} \left( \frac{20.3 \text{ kg}}{day} \right) \times f_{p-dairy} (1) \times f_{s-dairy} (1) \times (R_{upp} + R_{es}) \right) + \left( Q_{s-dairy} \left( \frac{0.4 \text{ kg}}{day} \right) \times f_{p-dairy} (1) \right) \right]}$$

where:

$$R_{upp} = Bv_{dry} \left( \frac{pCi/g-dry \text{ plant}}{pCi/g-dry \text{ soil}} \right); R_{es} = MLF_{pasture} \left( \frac{0.25 \text{ g-dry soil}}{g-dry \text{ plant}} \right)$$

- **consumption of swine.** Table 13-51 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced swine.

$$DCC_{soil-far-swine-ing} (pCi/g) = \frac{DCC_{far-swine-ing} (pCi/g)}{TF_{swine} \left( \frac{day}{kg} \right) \times \left[ \left( Q_{p-swine} \left( \frac{4.7 \text{ kg}}{day} \right) \times f_{p-swine} (1) \times f_{s-swine} (1) \times (R_{upp} + R_{es}) \right) + \left( Q_{s-swine} \left( \frac{0.37 \text{ kg}}{day} \right) \times f_{p-swine} (1) \right) \right]}$$

where:

$$R_{upp} = Bv_{dry} \left( \frac{pCi/g-dry \text{ plant}}{pCi/g-dry \text{ soil}} \right); R_{es} = MLF_{pasture} \left( \frac{0.25 \text{ g-dry soil}}{g-dry \text{ plant}} \right)$$

- **total.**

$$DCC_{soil-far-tot} (pCi/g) = \frac{\frac{1}{DCC_{soil-far-soil-ing}} + \frac{1}{DCC_{soil-far-soil-inh}} + \frac{1}{DCC_{soil-far-soil-ext}} + \frac{1}{DCC_{soil-far-produce-ing-tot}} + \frac{1}{DCC_{soil-far-egg-ing}}}{\frac{1}{DCC_{soil-far-poultry-ing}} + \frac{1}{DCC_{soil-far-fish-ing}} + \frac{1}{DCC_{soil-far-beef-ing}} + \frac{1}{DCC_{soil-far-dairy-ing}} + \frac{1}{DCC_{soil-far-swine-ing}}}$$

The following consumption routes are provided in site-specific mode only and requires the user to enter their own data, as the tool only provides a transfer factor.

consumption of goat. Please see table 1, in section 5 below, for sources of derived intakes for home-produced goat.

$$DCC_{\text{soil-far-goat-ing}} (\text{pCi/g}) = \frac{DCC_{\text{far-goat-ing}} (\text{pCi/g})}{TF_{\text{goat}} \left( \frac{\text{day}}{\text{kg}} \right) \times \left[ \left( Q_{\text{p-goat}} \left( \frac{1.27 \text{ kg}}{\text{day}} \right) \times f_{\text{p-goat}} (1) \times f_{\text{s-goat}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-goat}} \left( \frac{0.23 \text{ kg}}{\text{day}} \right) \times f_{\text{p-goat}} (1) \right) \right]}$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{\text{pCi/g-dry plant}}{\text{pCi/g-dry soil}} \right); R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{\text{g-dry plant}} \right)$$

- consumption of goat milk. Please see table 1, in section 5 below, for sources of derived intakes for home-produced goat dairy.

$$DCC_{\text{soil-far-goat-milk-ing}} (\text{pCi/g}) = \frac{DCC_{\text{far-goat-milk-ing}} (\text{pCi/g})}{TF_{\text{goat-milk}} \left( \frac{\text{day}}{\text{L milk}} \right) \times \rho_m \left( \frac{1.03 \text{ kg}}{\text{L milk}} \right)^{-1} \times \left[ \left( Q_{\text{p-goat-milk}} \left( \frac{1.59 \text{ kg}}{\text{day}} \right) \times f_{\text{p-goat-milk}} (1) \times f_{\text{s-goat-milk}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-goat-milk}} \left( \frac{0.29 \text{ kg}}{\text{day}} \right) \times f_{\text{p-goat-milk}} (1) \right) \right]}$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{\text{pCi/g-dry plant}}{\text{pCi/g-dry soil}} \right); R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{\text{g-dry plant}} \right)$$

- consumption of sheep. Please see table 1, in section 5 below, for sources of derived intakes for home-produced sheep.

$$DCC_{\text{soil-far-sheep-ing}} (\text{pCi/g}) = \frac{DCC_{\text{far-sheep-ing}} (\text{pCi/g})}{TF_{\text{sheep}} \left( \frac{\text{day}}{\text{kg}} \right) \times \left[ \left( Q_{\text{p-sheep}} \left( \frac{1.75 \text{ kg}}{\text{day}} \right) \times f_{\text{p-sheep}} (1) \times f_{\text{s-sheep}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-sheep}} \left( \frac{0.32 \text{ kg}}{\text{day}} \right) \times f_{\text{p-sheep}} (1) \right) \right]}$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{\text{pCi/g-dry plant}}{\text{pCi/g-dry soil}} \right); R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{\text{g-dry plant}} \right)$$

- consumption of sheep milk. Please see table 1, in section 5 below, for sources of derived intakes for home-produced sheep dairy.

$$DCC_{\text{soil-far-sheep-milk-ing}} (\text{pCi/g}) = \frac{DCC_{\text{far-sheep-milk-ing}} (\text{pCi/g})}{TF_{\text{sheep-milk}} \left( \frac{\text{day}}{\text{L milk}} \right) \times \rho_m \left( \frac{1.03 \text{ kg}}{\text{L milk}} \right)^{-1} \times \left[ \left( Q_{\text{p-sheep-milk}} \left( \frac{3.15 \text{ kg}}{\text{day}} \right) \times f_{\text{p-sheep-milk}} (1) \times f_{\text{s-sheep-milk}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-sheep-milk}} \left( \frac{0.57 \text{ kg}}{\text{day}} \right) \times f_{\text{p-sheep-milk}} (1) \right) \right]}$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{\text{pCi/g-dry plant}}{\text{pCi/g-dry soil}} \right); R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{\text{g-dry plant}} \right)$$

Definitions of the input variables are in [Table 1](#).

## CDI Equations

- incidental ingestion of soil

$$DCC_{\text{far-soil-ing}} (\text{pCi/g}) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_o \left( \frac{\text{mrem}}{\text{pCi}} \right) \times IFS_{\text{far-adj}} \left( \frac{40,250 \text{ mg}}{\text{year}} \right) \times \left( \frac{\text{g}}{1000 \text{ mg}} \right)}$$

where:

$$IFS_{\text{far-adj}} \left( \frac{40,250 \text{ mg}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRS_{\text{far-c}} \left( \frac{200 \text{ mg}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRS_{\text{far-a}} \left( \frac{100 \text{ mg}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- inhalation of particulates emitted from soil

$$CDI_{\text{far-soil-ext}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{soil}} \left( \frac{\text{pCi}}{\text{g}} \right) \times EF_{\text{far}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ACF_{\text{ext-sv}} \times \left[ \left( ET_{\text{far-o}} \left( \frac{12,168 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{ext-sv}} (1.0) \right) + \left( ET_{\text{far-i}} \left( \frac{10,008 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_{\text{i-total}} \right) \right]$$

- external exposure to ionizing radiation

$$CDI_{\text{far-soil-inh}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{soil}} \left( \frac{\text{pCi}}{\text{g}} \right) \times IFA_{\text{far-adj}} \left( \frac{6,475 \text{ m}^3}{\text{year}} \right) \times \frac{1}{PEF \left( \frac{\text{m}^3}{\text{kg}} \right)} \times \left( \frac{1000 \text{ g}}{\text{kg}} \right)$$

where:

$$IFA_{\text{far-adj}} \left( \frac{6,475 \text{ m}^3}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times ET_{\text{far-c}} \left( \frac{24 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{far-c}} \left( \frac{10 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times ET_{\text{far-a}} \left( \frac{24 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{far-a}} \left( \frac{20 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- consumption of produce (fruits and vegetables). Sections 9 and 13 of the 2011 Exposure Factors Handbook were used to derive the intakes for home-grown produce.

$$CDI_{\text{soil-far-produce-ing-tot}} \left( \frac{pCi}{\text{year}} \right) = \sum_{i=1}^n CDI_{\text{soil-far-produce-ing}} \left( \frac{pCi}{\text{year}} \right)_i$$

where:  
n = total number of produce items included

and:

$$CDI_{\text{soil-far-produce-ing}} \left( \frac{pCi}{\text{year}} \right) = C_{\text{soil}} \left( \frac{pCi}{g} \right) \times (R_{\text{upv}} + R_{\text{es}}) \times IF_{\text{far-adj}} \left( \frac{g}{\text{year}} \right) \times CF_{\text{far-produce}} \quad (1)$$

where:

$$R_{\text{upv}} = Bv_{\text{wet}} \left( \frac{pCi / g\text{-fresh plant}}{pCi / g\text{-dry soil}} \right); R_{\text{es}} = MLF_{\text{produce}} \left( \frac{g\text{-dry soil}}{g\text{-fresh plant}} \right)$$

and:

$$IF_{\text{far-adj}} \left( \frac{g}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IR_{\text{far-c}} \left( \frac{g}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IR_{\text{far-a}} \left( \frac{g}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

and:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

The consumption of produce exposure route drives the DCCs lower than all the other routes. It is recommended that produce-specific transfer factors ( $Bv_{\text{wet}}$ ) be used when available for a site. Further, the default transfer factors ( $Bv_{\text{wet}}$ ) from IAEA used in these DCC calculations are based on a composite of all soil groups. Transfer factors ( $Bv_{\text{wet}}$ ) for sand, loam, clay, organic, coral sand, and other soil types that may be more suited to a particular site are also provided. The site-specific option of the calculator can be used to focus on ingestion of individual produce types. When "Site-specific" is selected, if the user changes the "Select Isotope Info Type" to "User-provided", then a specific transfer factor may be changed.

where:

$$CDI_{\text{far-produce-ing-tot}} \left( \frac{pCi}{\text{year}} \right) = \sum_{i=1}^n CDI_{\text{far-produce-ing}} \left( \frac{pCi}{\text{year}} \right)_i$$

where:  
n = total number of produce items included

and:

$$CDI_{\text{far-produce-ing}} \left( \frac{pCi}{\text{year}} \right) = C_{\text{produce}} \left( \frac{pCi}{g} \right) \times IF_{\text{far-adj}} \left( \frac{g}{\text{year}} \right) \times CF_{\text{far-produce}} \quad (1)$$

where:

$$IF_{\text{far-adj}} \left( \frac{g}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IR_{\text{far-c}} \left( \frac{g}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IR_{\text{far-a}} \left( \frac{g}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

and:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- consumption of eggs. Table 13-40 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced eggs.

$$CDI_{\text{soil-far-egg-ing}} \left( \frac{pCi}{\text{year}} \right) = \left[ C_{\text{soil}} \left( \frac{pCi}{g} \right) \times IFE_{\text{far-adj}} \left( \frac{19,395 \text{ g}}{\text{year}} \right) \times CF_{\text{far-egg}} (1) \times TF_{\text{egg}} \left( \frac{\text{day}}{\text{kg}} \right) \times \left( \left( Q_{\text{p-poultry}} \left( \frac{0.2 \text{ kg}}{\text{day}} \right) \times f_{\text{p-poultry}} (1) \times f_{\text{s-poultry}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-poultry}} \left( \frac{0.022 \text{ kg}}{\text{day}} \right) \times f_{\text{p-poultry}} (1) \right) \right) \right]$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{pCi / g\text{-dry plant}}{pCi / g\text{-dry soil}} \right); R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{g\text{-dry plant}} \right)$$

and:

$$IFE_{\text{far-adj}} \left( \frac{19,395 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRE_{\text{far-c}} \left( \frac{31.7 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRE_{\text{far-a}} \left( \frac{59.6 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- consumption of poultry. Table 13-52 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced poultry.

$$CDI_{\text{soil-far-poultry-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = \left[ \begin{aligned} & C_{\text{soil}} \left( \frac{\text{pCi}}{\text{g}} \right) \times IFP_{\text{far-adj}} \left( \frac{34,414 \text{ g}}{\text{year}} \right) \times CF_{\text{far-poultry}} (1) \times TF_{\text{poultry}} \left( \frac{\text{day}}{\text{kg}} \right) \times \\ & \left[ \begin{aligned} & Q_{\text{p-poultry}} \left( \frac{0.2 \text{ kg}}{\text{day}} \right) \times f_{\text{p-poultry}} (1) \times f_{\text{s-poultry}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right] + \\ & \left[ \begin{aligned} & Q_{\text{s-poultry}} \left( \frac{0.022 \text{ kg}}{\text{day}} \right) \times f_{\text{p-poultry}} (1) \end{aligned} \right] \end{aligned} \right] \end{aligned}$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{\text{pCi / g-dry plant}}{\text{pCi / g-dry soil}} \right); R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{\text{g-dry plant}} \right)$$

and:

$$IFP_{\text{far-adj}} \left( \frac{34,414 \text{ g}}{\text{year}} \right) = \left( \begin{aligned} & \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRP_{\text{far-c}} \left( \frac{46.9 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \\ & \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRP_{\text{far-a}} \left( \frac{107.4 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \end{aligned} \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- consumption of fish. Table 13-20 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-caught fish.

$$CDI_{\text{soil-far-fish-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = \frac{C_{\text{soil}} \left( \frac{\text{pCi}}{\text{g}} \right) \times BCF \left( \frac{\text{L}}{\text{kg}} \right)}{Kd \left( \frac{\text{L}}{\text{kg}} \right)} \times IFFI_{\text{far-adj}} \left( \frac{250,474 \text{ g}}{\text{year}} \right) \times CF_{\text{far-fish}} (1)$$

where:

$$IFFI_{\text{far-adj}} \left( \frac{250,474 \text{ g}}{\text{year}} \right) = \left( \begin{aligned} & \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRFI_{\text{far-c}} \left( \frac{57.4 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \\ & \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRFI_{\text{far-a}} \left( \frac{831.8 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \end{aligned} \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- consumption of beef. Table 13-33 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced beef.

$$CDI_{\text{soil-far-beef-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = \left[ \begin{aligned} & C_{\text{soil}} \left( \frac{\text{pCi}}{\text{g}} \right) \times IFB_{\text{far-adj}} \left( \frac{52,474 \text{ g}}{\text{year}} \right) \times CF_{\text{far-beef}} (1) \times TF_{\text{beef}} \left( \frac{\text{day}}{\text{kg}} \right) \times \\ & \left[ \begin{aligned} & Q_{\text{p-beef}} \left( \frac{11.77 \text{ kg}}{\text{day}} \right) \times f_{\text{p-beef}} (1) \times f_{\text{s-beef}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right] + \\ & \left[ \begin{aligned} & Q_{\text{s-beef}} \left( \frac{0.5 \text{ kg}}{\text{day}} \right) \times f_{\text{p-beef}} (1) \end{aligned} \right] \end{aligned} \right]$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{\text{pCi / g-dry plant}}{\text{pCi / g-dry soil}} \right); R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{\text{g-dry plant}} \right)$$

and:

$$IFB_{\text{far-adj}} \left( \frac{52,474 \text{ g}}{\text{year}} \right) = \left( \begin{aligned} & \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRB_{\text{far-c}} \left( \frac{62.8 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \\ & \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRB_{\text{far-a}} \left( \frac{165.3 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \end{aligned} \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- consumption of milk. Table 11-4 of the 2011 Exposure Factors Handbook was used to derive the intakes for total dairy.

$$CDI_{\text{soil-far-dairy-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = \left[ \begin{aligned} & C_{\text{soil}} \left( \frac{\text{pCi}}{\text{g}} \right) \times IFD_{\text{far-adj}} \left( \frac{253,451 \text{ g}}{\text{year}} \right) \times CF_{\text{far-dairy}} (1) \times TF_{\text{dairy}} \left( \frac{\text{day}}{\text{kg}} \right) \times \\ & \rho_m \left( \frac{1.03 \text{ kg}}{1 \text{ L milk}} \right)^{-1} \times \left[ \begin{aligned} & Q_{\text{p-dairy}} \left( \frac{20.3 \text{ kg}}{\text{day}} \right) \times f_{\text{p-dairy}} (1) \times f_{\text{s-dairy}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right] + \\ & \left[ \begin{aligned} & Q_{\text{s-dairy}} \left( \frac{0.4 \text{ kg}}{\text{day}} \right) \times f_{\text{p-dairy}} (1) \end{aligned} \right] \end{aligned} \right]$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{\text{pCi / g-dry plant}}{\text{pCi / g-dry soil}} \right); R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{\text{g-dry plant}} \right)$$

and:

$$IFD_{\text{far-adj}} \left( \frac{253,451 \text{ g}}{\text{year}} \right) = \left( \begin{aligned} & \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRD_{\text{far-c}} \left( \frac{994.7 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \\ & \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRD_{\text{far-a}} \left( \frac{676.4 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \end{aligned} \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- consumption of swine. Table 13-51 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced swine.

$$CDI_{\text{soil-far-swine-ing}} \left( \frac{pCi}{\text{year}} \right) = \left[ \begin{aligned} & C_{\text{soil}} \left( \frac{pCi}{g} \right) \times IFSW_{\text{far-adj}} \left( \frac{29,288 g}{\text{year}} \right) \times CF_{\text{far-swine}} (1) \times TF_{\text{swine}} \left( \frac{\text{day}}{kg} \right) \times \\ & \left[ \begin{aligned} & \left( Q_{\text{p-swine}} \left( \frac{4.7 kg}{\text{day}} \right) \times f_{\text{p-swine}} (1) \times f_{\text{s-swine}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \\ & \left( Q_{\text{s-swine}} \left( \frac{0.37 kg}{\text{day}} \right) \times f_{\text{p-swine}} (1) \right) \end{aligned} \right] \end{aligned} \right]$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{pCi / g\text{-dry plant}}{pCi / g\text{-dry soil}} \right) ; R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 g\text{-dry soil}}{g\text{-dry plant}} \right)$$

and:

$$IFSW_{\text{far-adj}} \left( \frac{29,288 g}{\text{year}} \right) = \left[ \begin{aligned} & \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSW_{\text{far-c}} \left( \frac{33.7 g}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \\ & \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSW_{\text{far-a}} \left( \frac{92.5 g}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \end{aligned} \right]$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

The following consumption routes are provided in site-specific mode only and requires the user to enter their own data, as the tool only provides a transfer factor.

- **consumption of goat.** Please see table 1, in section 5 below, for sources of derived intakes for home-produced goat.

$$CDI_{\text{soil-far-goat-ing}} \left( \frac{pCi}{\text{year}} \right) = \left[ \begin{aligned} & C_{\text{soil}} \left( \frac{pCi}{g} \right) \times IFGO_{\text{far-adj}} \left( \frac{g}{\text{year}} \right) \times CF_{\text{far-goat}} (1) \times TF_{\text{goat}} \left( \frac{\text{day}}{kg} \right) \times \\ & \left[ \begin{aligned} & \left( Q_{\text{p-goat}} \left( \frac{1.27 kg}{\text{day}} \right) \times f_{\text{p-goat}} (1) \times f_{\text{s-goat}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \\ & \left( Q_{\text{s-goat}} \left( \frac{0.23 kg}{\text{day}} \right) \times f_{\text{p-goat}} (1) \right) \end{aligned} \right] \end{aligned} \right]$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{pCi / g\text{-dry plant}}{pCi / g\text{-dry soil}} \right) ; R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 g\text{-dry soil}}{g\text{-dry plant}} \right)$$

and:

$$IFGO_{\text{far-adj}} \left( \frac{g}{\text{year}} \right) = \left[ \begin{aligned} & \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRGO_{\text{far-c}} \left( \frac{g}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \\ & \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRGO_{\text{far-a}} \left( \frac{g}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \end{aligned} \right]$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- **consumption of goat milk.** Please see table 1, in section 5 below, for sources of derived intakes for home-produced goat dairy.

$$CDI_{\text{soil-far-goat-milk-ing}} \left( \frac{pCi}{\text{year}} \right) = \left[ \begin{aligned} & C_{\text{soil}} \left( \frac{pCi}{g} \right) \times IFGM_{\text{far-adj}} \left( \frac{g}{\text{year}} \right) \times CF_{\text{far-goat-milk}} (1) \times TF_{\text{goat-milk}} \left( \frac{\text{day}}{L \text{ milk}} \right) \times \\ & p_m \left( \frac{1.03 kg}{1 L \text{ milk}} \right)^{-1} \times \left[ \begin{aligned} & \left( Q_{\text{p-goat-milk}} \left( \frac{1.59 kg}{\text{day}} \right) \times f_{\text{p-goat-milk}} (1) \times f_{\text{s-goat-milk}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \\ & \left( Q_{\text{s-goat-milk}} \left( \frac{0.29 kg}{\text{day}} \right) \times f_{\text{p-goat-milk}} (1) \right) \end{aligned} \right] \end{aligned} \right]$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{pCi / g\text{-dry plant}}{pCi / g\text{-dry soil}} \right) ; R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 g\text{-dry soil}}{g\text{-dry plant}} \right)$$

and:

$$IFGM_{\text{far-adj}} \left( \frac{g}{\text{year}} \right) = \left[ \begin{aligned} & \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRGM_{\text{far-c}} \left( \frac{g}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \\ & \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRGM_{\text{far-a}} \left( \frac{g}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \end{aligned} \right]$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- **consumption of sheep.** Please see table 1, in section 5 below, for sources of derived intakes for home-produced sheep.



$$CDI_{\text{soil-far-sheep-ing}} \left( \frac{pCi}{\text{year}} \right) = \left[ \begin{aligned} & C_{\text{soil}} \left( \frac{pCi}{g} \right) \times IFSH_{\text{far-adj}} \left( \frac{g}{\text{year}} \right) \times CF_{\text{far-sheep}} (1) \times TF_{\text{sheep}} \left( \frac{\text{day}}{kg} \right) \times \\ & \left[ \begin{aligned} & Q_{\text{p-sheep}} \left( \frac{1.75 \text{ kg}}{\text{day}} \right) \times f_{\text{p-sheep}} (1) \times f_{\text{s-sheep}} (1) \times (R_{\text{upp}} + R_{\text{es}}) + \\ & Q_{\text{s-sheep}} \left( \frac{0.32 \text{ kg}}{\text{day}} \right) \times f_{\text{p-sheep}} (1) \end{aligned} \right] \end{aligned} \right]$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{pCi / g\text{-dry plant}}{pCi / g\text{-dry soil}} \right) ; R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{g\text{-dry plant}} \right)$$

and:

$$IFSH_{\text{far-adj}} \left( \frac{g}{\text{year}} \right) = \left[ \begin{aligned} & \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSH_{\text{far-c}} \left( \frac{g}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \\ & \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSH_{\text{far-a}} \left( \frac{g}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \end{aligned} \right]$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- consumption of sheep milk. Please see table 1, in section 5 below, for sources of derived intakes for home-produced sheep dairy.

$$CDI_{\text{soil-far-sheep-milk-ing}} \left( \frac{pCi}{\text{year}} \right) = \left[ \begin{aligned} & C_{\text{soil}} \left( \frac{pCi}{g} \right) \times IFSM_{\text{far-adj}} \left( \frac{g}{\text{year}} \right) \times CF_{\text{far-sheep-milk}} (1) \times TF_{\text{sheep-milk}} \left( \frac{\text{day}}{L \text{ milk}} \right) \times \\ & p_m \left( \frac{1.03 \text{ kg}}{L \text{ milk}} \right)^{-1} \times \left[ \begin{aligned} & Q_{\text{p-sheep-milk}} \left( \frac{3.15 \text{ kg}}{\text{day}} \right) \times f_{\text{p-sheep-milk}} (1) \times f_{\text{s-sheep-milk}} (1) \times (R_{\text{upp}} + R_{\text{es}}) + \\ & Q_{\text{s-sheep-milk}} \left( \frac{0.57 \text{ kg}}{\text{day}} \right) \times f_{\text{p-sheep-milk}} (1) \end{aligned} \right] \end{aligned} \right]$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{pCi / g\text{-dry plant}}{pCi / g\text{-dry soil}} \right) ; R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{g\text{-dry plant}} \right)$$

and:

$$IFSM_{\text{far-adj}} \left( \frac{g}{\text{year}} \right) = \left[ \begin{aligned} & \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSM_{\text{far-c}} \left( \frac{g}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \\ & \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSM_{\text{far-a}} \left( \frac{g}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \end{aligned} \right]$$

where:


$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

Definitions of the input variables are in [Table 1](#).

### 4.8.3 Farmer Direct Exposure and Consumption of Agricultural Products - Back-Calculated to Water

The farmer scenario should be considered an extension of the resident scenario and evaluate consumption of farm products for a subsistence farmer. Like the resident, the farmer scenario assumes the receptor will be exposed via the following pathways: ingestion of tapwater, external radiation from contaminants in tapwater, inhalation of gases in tapwater, and consumption of home grown produce (100% of fruit and vegetables are from the farm). The inhalation exposure route is only calculated for C-14, H-3, Ra-224, and Ra-226, which all volatilize. In addition to produce, 100% of consumption of the following are also considered to be from the farm: beef, milk, fish, swine, egg, and poultry. All water (100%) for farm products is considered to have been provided from contaminated portions of the site. For these farm products, dose-based DCCs are provided for the water, which may contribute contaminants to the products. Also like the resident, age-adjusted intake equations were developed for all of the ingestion/consumption equations to account for changes in intake as the receptor ages.

#### Graphical Representation



## DCC Equations

- ingestion of tapwater

$$DCC_{\text{water-far-ing}} \left( \text{pCi/L} \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_0 \left( \frac{\text{mrem}}{\text{pCi}} \right) \times \left( IFW_{\text{far-adj}} \left( \frac{785 \text{ L}}{\text{year}} \right) \right)}$$

where:

$$IFW_{\text{far-adj}} \left( \frac{785 \text{ L}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRW_{\text{far-c}} \left( \frac{0.78 \text{ L}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRW_{\text{far-a}} \left( \frac{2.5 \text{ L}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- immersion in tapwater

$$DCC_{\text{water-far-imm}} \left( \text{pCi/L} \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_{\text{imm}} \left( \frac{\text{mrem/year}}{\text{pCi/L}} \right) \times DFA_{\text{far-adj}} \left( \frac{240 \text{ hours}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{8760 \text{ hours}} \right)}$$

where:

$$DFA_{\text{far-adj}} \left( \frac{240 \text{ hours}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times EV_{\text{far-c}} \left( \frac{1 \text{ event}}{\text{day}} \right) \times ET_{\text{event-far-c}} \left( \frac{0.54 \text{ hours}}{\text{event}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times EV_{\text{far-a}} \left( \frac{1 \text{ event}}{\text{day}} \right) \times ET_{\text{event-far-a}} \left( \frac{0.71 \text{ hours}}{\text{event}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- inhalation

$$DCC_{\text{water-far-inh}} \left( \text{pCi/L} \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_i \left( \frac{\text{mrem}}{\text{pCi}} \right) \times IFA_{\text{far-adj}} \left( \frac{6,475 \text{ m}^3}{\text{year}} \right) \times K \left( \frac{0.5 \text{ L}}{\text{m}^3} \right)}$$

where:

$$IFA_{\text{far-adj}} \left( \frac{6,475 \text{ m}^3}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times ET_{\text{far-c}} \left( \frac{24 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{far-c}} \left( \frac{10 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times ET_{\text{far-a}} \left( \frac{24 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{far-a}} \left( \frac{20 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{far-c}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- consumption of produce (fruits and vegetables). Sections 9 and 13 of the 2011 Exposure Factors Handbook were used to derive the intakes for home-grown produce.

$$DCC_{\text{water-far-produce-ing-tot}} \left( \text{pCi/L} \right) = \frac{1}{\sum_{i=1}^n DCC_{\text{water-far-produce-ing}} \left( \text{pCi/L} \right)_i}$$

where:  
n = total number of produce items included

and:

$$DCC_{\text{water-far-produce}} \left( \text{pCi/L} \right) = \frac{DCC_{\text{far-produce-ing}} \left( \text{pCi/g} \right)}{\left( \frac{1 \text{ kg}}{1000 \text{ g}} \right) \times \left( Irr_{\text{rup}} \left( \frac{\text{L}}{\text{kg}} \right) + Irr_{\text{res}} \left( \frac{\text{L}}{\text{kg}} \right) + Irr_{\text{dep}} \left( \frac{\text{L}}{\text{kg}} \right) \right)}$$

where:

$$Irr_{\text{res}} \left( \frac{\text{L}}{\text{kg}} \right) = \frac{Ir \left( \frac{\text{L}}{\text{m}^2 \cdot \text{day}} \right) \times F \times MLF_{\text{produce}} \times \left[ 1 - \exp \left( - \left( \frac{\lambda_B}{\text{day}} \right) \times t_b \text{ (days)} \right) \right]}{P \left( \frac{\text{kg}}{\text{m}^2} \right) \times \left( \frac{\lambda_B}{\text{day}} \right)};$$

$$Irr_{\text{rup}} \left( \frac{\text{L}}{\text{kg}} \right) = \frac{Ir \left( \frac{\text{L}}{\text{m}^2 \cdot \text{day}} \right) \times F \times B_{\text{wet}} \times \left[ 1 - \exp \left( - \left( \frac{\lambda_B}{\text{day}} \right) \times t_b \text{ (days)} \right) \right]}{P \left( \frac{\text{kg}}{\text{m}^2} \right) \times \left( \frac{\lambda_B}{\text{day}} \right)};$$

and:

$$Irr_{\text{dep}} \left( \frac{\text{L}}{\text{kg}} \right) = \frac{Ir \left( \frac{\text{L}}{\text{m}^2 \cdot \text{day}} \right) \times F \times I_f \times T \times \left[ 1 - \exp \left( - \left( \frac{\lambda_E}{\text{day}} \right) \times t_b \text{ (days)} \right) \right]}{Y_v \left( \frac{\text{kg}}{\text{m}^2} \right) \times \left( \frac{\lambda_E}{\text{day}} \right)}$$

where:

$$DCC_{\text{far-produce-ing-tot}} (\text{pCi/g}) = \frac{1}{\left( \sum_{i=1}^n \frac{1}{DCC_{\text{far-produce-ing}} (\text{pCi/g})_i} \right)}$$

where:

n = total number of produce items included

and:

$$DCC_{\text{far-produce-ing}} (\text{pCi/g}) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_o \left( \frac{\text{mrem}}{\text{pCi}} \right) \times IF_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) \times CF_{\text{far-produce}} (1)}$$

where:

$$IF_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IR_{\text{far-c}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IR_{\text{far-a}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

and:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- **consumption of eggs.** Table 13-40 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced eggs.

$$DCC_{\text{water-far-egg-ing}} (\text{pCi/L}) = \frac{DCC_{\text{far-egg-ing}} (\text{pCi/g})}{TF_{\text{egg}} \left( \frac{\text{day}}{\text{kg}} \right) \times Q_{\text{w-poultry}} \left( \frac{0.4 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

- **consumption of poultry.** Table 13-52 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced poultry.

$$DCC_{\text{water-far-poultry-ing}} (\text{pCi/L}) = \frac{DCC_{\text{far-poultry-ing}} (\text{pCi/g})}{TF_{\text{poultry}} \left( \frac{\text{day}}{\text{kg}} \right) \times Q_{\text{w-poultry}} \left( \frac{0.4 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

- **consumption of fish.** Table 13-20 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-caught fish.

$$DCC_{\text{water-far-fish-ing}} (\text{pCi/L}) = \frac{DCC_{\text{far-fish-ing}} (\text{pCi/g})}{BCF \left( \frac{\text{L}}{\text{kg}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

- **consumption of beef.** Table 13-33 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced beef.

$$DCC_{\text{water-far-beef-ing}} (\text{pCi/L}) = \frac{DCC_{\text{far-beef-ing}} (\text{pCi/g})}{TF_{\text{beef}} \left( \frac{\text{day}}{\text{kg}} \right) \times Q_{\text{w-beef}} \left( \frac{53 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

- **consumption of milk.** Table 11-4 of the 2011 Exposure Factors Handbook was used to derive the intakes for total dairy.

$$DCC_{\text{water-far-dairy-ing}} (\text{pCi/L}) = \frac{DCC_{\text{far-dairy-ing}} (\text{pCi/g})}{TF_{\text{dairy}} \left( \frac{\text{day}}{\text{L milk}} \right) \times p_m \left( \frac{1.03 \text{ kg}}{1 \text{ L milk}} \right)^{-1} \times Q_{\text{w-dairy}} \left( \frac{92 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

- **consumption of swine.** Table 13-51 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced swine.

$$DCC_{\text{water-far-swine-ing}} (\text{pCi/L}) = \frac{DCC_{\text{far-swine-ing}} (\text{pCi/g})}{TF_{\text{swine}} \left( \frac{\text{day}}{\text{kg}} \right) \times Q_{\text{w-swine}} \left( \frac{11.4 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

- **total**

$$DCC_{\text{wat-far-tot}} (\text{pCi/L}) = \frac{1}{\frac{1}{DCC_{\text{water-far-ing}}} + \frac{1}{DCC_{\text{water-far-inh}}} + \frac{1}{DCC_{\text{water-far-imm}}} + \frac{1}{DCC_{\text{water-far-produce-ing-tot}}} + \frac{1}{DCC_{\text{water-far-egg-ing}}} + \frac{1}{DCC_{\text{water-far-poultry-ing}}} + \frac{1}{DCC_{\text{water-far-fish-ing}}} + \frac{1}{DCC_{\text{water-far-beef-ing}}} + \frac{1}{DCC_{\text{water-far-dairy-ing}}} + \frac{1}{DCC_{\text{water-far-swine-ing}}}}$$

The following consumption routes are provided in site-specific mode only and requires the user to enter their own data, as the tool only provides a transfer factor.

- **consumption of goat.** Please see table 1, in section 5 below, for sources of derived intakes for home-produced goat.

$$DCC_{\text{water-far-goat-ing}} (\text{pCi/L}) = \frac{DCC_{\text{far-goat-ing}} (\text{pCi/g})}{TF_{\text{goat}} \left( \frac{\text{day}}{\text{kg}} \right) \times Q_{\text{w-goat}} \left( \frac{3.81 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

- **consumption of goat milk.** Please see table 1, in section 5 below, for sources of derived intakes for home-produced goat dairy.

$$DCC_{\text{water-far-goat-milk-ing}} (\text{pCi/L}) = \frac{DCC_{\text{far-goat-milk-ing}} (\text{pCi/g})}{TF_{\text{goat-milk}} \left( \frac{\text{day}}{\text{L milk}} \right) \times p_m \left( \frac{1.03 \text{ kg}}{1 \text{ L milk}} \right)^{-1} \times Q_{\text{w-goat-milk}} \left( \frac{8.75 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

- **consumption of sheep.** Please see table 1, in section 5 below, for sources of derived intakes for home-produced goat.

$$DCC_{\text{water-far-sheep-ing}} (\text{pCi/L}) = \frac{DCC_{\text{far-sheep-ing}} (\text{pCi/g})}{TF_{\text{sheep}} \left( \frac{\text{day}}{\text{kg}} \right) \times Q_{\text{w-sheep}} \left( \frac{5.25 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

- **consumption of sheep milk.** Please see table 1, in section 5 below, for sources of derived intakes for home-produced sheep dairy.

$$DCC_{\text{water-far-sheep-milk-ing}} (\text{pCi/L}) = \frac{DCC_{\text{far-sheep-milk-ing}} (\text{pCi/g})}{TF_{\text{sheep-milk}} \left( \frac{\text{day}}{\text{L milk}} \right) \times p_m \left( \frac{1.03 \text{ kg}}{1 \text{ L milk}} \right)^{-1} \times Q_{\text{w-sheep-milk}} \left( \frac{10.4 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

Definitions of the input variables are in [Table 1](#).

### CDI Equations

- ingestion of tapwater

$$CDI_{\text{water-far-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{water}} \left( \frac{\text{pCi}}{\text{L}} \right) \times \left( IFW_{\text{far-adj}} \left( \frac{785 \text{ L}}{\text{year}} \right) \right)$$

where:

$$IFW_{\text{far-adj}} \left( \frac{785 \text{ L}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRW_{\text{far-c}} \left( \frac{0.78 \text{ L}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRW_{\text{far-a}} \left( \frac{2.5 \text{ L}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- immersion in tapwater

$$CDI_{\text{water-far-imm}} \left( \frac{\text{pCi}}{\text{L}} \right) = C_{\text{water}} \left( \frac{\text{pCi}}{\text{L}} \right) \times DFA_{\text{far-adj}} \left( \frac{240 \text{ hours}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{8760 \text{ hours}} \right)$$

where:

$$DFA_{\text{far-adj}} \left( \frac{240 \text{ hours}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times EV_{\text{far-c}} \left( \frac{1 \text{ event}}{\text{day}} \right) \times ET_{\text{event-far-c}} \left( \frac{0.54 \text{ hours}}{\text{event}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times EV_{\text{far-a}} \left( \frac{1 \text{ event}}{\text{day}} \right) \times ET_{\text{event-far-a}} \left( \frac{0.71 \text{ hours}}{\text{event}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- inhalation

$$CDI_{\text{water-far-inh}} \left( \frac{\text{pCi}}{\text{year}} \right) = C_{\text{water}} \left( \frac{\text{pCi}}{\text{L}} \right) \times IFA_{\text{far-adj}} \left( \frac{6,475 \text{ m}^3}{\text{year}} \right) \times K \left( \frac{0.5 \text{ L}}{\text{m}^3} \right)$$

where:

$$IFA_{\text{far-adj}} \left( \frac{6,475 \text{ m}^3}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times ET_{\text{far-c}} \left( \frac{24 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{far-c}} \left( \frac{10 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times ET_{\text{far-a}} \left( \frac{24 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{far-a}} \left( \frac{20 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{far-c}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- consumption of produce (fruits and vegetables). Sections 9 and 13 of the 2011 Exposure Factors Handbook were used to derive the intakes for home-grown produce.

$$CDI_{\text{water-far-produce-ing-tot}} \left( \frac{pCi}{\text{year}} \right) = \sum_{i=1}^n CDI_{\text{water-far-produce-ing}} \left( \frac{pCi}{\text{year}} \right)_i$$

where:

n = total number of produce items included

and:

$$CDI_{\text{water-far-produce-ing}} \left( \frac{pCi}{\text{year}} \right) = \left( C_{\text{water}} \left( \frac{pCi}{L} \right) \times IF_{\text{far-adj}} \left( \frac{g}{\text{year}} \right) \times CF_{\text{far-produce}} (1) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right) \times \left( Irr_{\text{rup}} \left( \frac{L}{\text{kg}} \right) + Irr_{\text{res}} \left( \frac{L}{\text{kg}} \right) + Irr_{\text{dep}} \left( \frac{L}{\text{kg}} \right) \right) \right)$$

where:

$$Irr_{\text{res}} \left( \frac{L}{\text{kg}} \right) = \frac{Ir \left( \frac{L}{m^2 \cdot \text{day}} \right) \times F \times MLF_{\text{produce}} \times \left[ 1 - \exp \left( - \left( \frac{\lambda_B}{\text{day}} \right) \times t_b \text{ (days)} \right) \right]}{P \left( \frac{\text{kg}}{m^2} \right) \times \left( \frac{\lambda_B}{\text{day}} \right)};$$

$$Irr_{\text{rup}} \left( \frac{L}{\text{kg}} \right) = \frac{Ir \left( \frac{L}{m^2 \cdot \text{day}} \right) \times F \times Bv_{\text{wet}} \times \left[ 1 - \exp \left( - \left( \frac{\lambda_B}{\text{day}} \right) \times t_b \text{ (days)} \right) \right]}{P \left( \frac{\text{kg}}{m^2} \right) \times \left( \frac{\lambda_B}{\text{day}} \right)};$$

$$Irr_{\text{dep}} \left( \frac{L}{\text{kg}} \right) = \frac{Ir \left( \frac{L}{m^2 \cdot \text{day}} \right) \times F \times I_f \times T \times \left[ 1 - \exp \left( - \left( \frac{\lambda_E}{\text{day}} \right) \times t_b \text{ (days)} \right) \right]}{Y_v \left( \frac{\text{kg}}{m^2} \right) \times \left( \frac{\lambda_E}{\text{day}} \right)}$$

and:

$$IF_{\text{far-adj}} \left( \frac{g}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IR_{\text{far-c}} \left( \frac{g}{\text{day}} \right) \times AAF_{\text{far-c}} (0.23) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IR_{\text{far-a}} \left( \frac{g}{\text{day}} \right) \times AAF_{\text{far-a}} (0.77) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.23) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.77) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

where:

$$CDI_{\text{far-produce-ing-tot}} \left( \frac{pCi}{\text{year}} \right) = \sum_{i=1}^n CDI_{\text{far-produce-ing}} \left( \frac{pCi}{\text{year}} \right)_i$$

where:

n = total number of produce items included

and:

$$CDI_{\text{far-produce-ing}} \left( \frac{pCi}{\text{year}} \right) = C_{\text{produce}} \left( \frac{pCi}{g} \right) \times IF_{\text{far-adj}} \left( \frac{g}{\text{year}} \right) \times CF_{\text{far-produce}} (1)$$

where:

$$IF_{\text{far-adj}} \left( \frac{g}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IR_{\text{far-c}} \left( \frac{g}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IR_{\text{far-a}} \left( \frac{g}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

and:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- consumption of eggs. Table 13-40 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced eggs.

$$CDI_{\text{water-far-egg-ing}} \left( \frac{pCi}{\text{year}} \right) = \left( C_{\text{water}} \left( \frac{pCi}{L} \right) \times IFE_{\text{far-adj}} \left( \frac{19,395 \text{ g}}{\text{year}} \right) \times CF_{\text{far-egg}} (1) \times \left( \frac{TF_{\text{egg}} \left( \frac{\text{day}}{\text{kg}} \right) \times Q_{w-\text{poultry}} \left( \frac{0.4 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)} \right) \right)$$

where:

$$IFE_{\text{far-adj}} \left( \frac{19,395 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRE_{\text{far-c}} \left( \frac{31.7 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRE_{\text{far-a}} \left( \frac{59.6 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- consumption of poultry. Table 13-52 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced poultry.

$$CDI_{\text{water-far-poultry-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = \left( C_{\text{water}} \left( \frac{\text{pCi}}{\text{L}} \right) \times IFP_{\text{far-adj}} \left( \frac{34,414 \text{ g}}{\text{year}} \right) \times CF_{\text{far-poultry}} (1) \times \right. \\ \left. TF_{\text{poultry}} \left( \frac{\text{day}}{\text{kg}} \right) \times Q_{\text{w-poultry}} \left( \frac{0.4 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right) \right)$$

where:

$$IFP_{\text{far-adj}} \left( \frac{34,414 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRP_{\text{far-c}} \left( \frac{46.9 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \right. \\ \left. \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRP_{\text{far-a}} \left( \frac{107.4 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- **consumption of fish.** Table 13-20 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-caught fish.

$$CDI_{\text{water-far-fish-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = \left( C_{\text{water}} \left( \frac{\text{pCi}}{\text{L}} \right) \times BCF \left( \frac{\text{L}}{\text{kg}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right) \times IFFI_{\text{far-adj}} \left( \frac{250,474 \text{ g}}{\text{year}} \right) \times CF_{\text{far-fish}} (1) \right)$$

where:

$$IFFI_{\text{far-adj}} \left( \frac{250,474 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRFI_{\text{far-c}} \left( \frac{57.4 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \right. \\ \left. \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRFI_{\text{far-a}} \left( \frac{831.8 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- **consumption of beef.** Table 13-33 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced beef.

$$CDI_{\text{water-far-beef-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = \left( C_{\text{water}} \left( \frac{\text{pCi}}{\text{L}} \right) \times IFB_{\text{far-adj}} \left( \frac{52,474 \text{ g}}{\text{year}} \right) \times CF_{\text{far-beef}} (1) \times \right. \\ \left. TF_{\text{beef}} \left( \frac{\text{day}}{\text{kg}} \right) \times Q_{\text{w-beef}} \left( \frac{53 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right) \right)$$

where:

$$IFB_{\text{far-adj}} \left( \frac{52,474 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRB_{\text{far-c}} \left( \frac{62.8 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \right. \\ \left. \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRB_{\text{far-a}} \left( \frac{165.3 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- **consumption of milk.** Table 11-4 of the 2011 Exposure Factors Handbook was used to derive the intakes for total dairy.

$$CDI_{\text{water-far-dairy-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = \left( C_{\text{water}} \left( \frac{\text{pCi}}{\text{L}} \right) \times IFD_{\text{far-adj}} \left( \frac{253,451 \text{ g}}{\text{year}} \right) \times CF_{\text{far-dairy}} (1) \times \right. \\ \left. TF_{\text{dairy}} \left( \frac{\text{day}}{\text{L milk}} \right) \times p_m \left( \frac{1.03 \text{ kg}}{1 \text{ L milk}} \right)^{-1} \times Q_{\text{w-dairy}} \left( \frac{92 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right) \right)$$

where:

$$IFD_{\text{far-adj}} \left( \frac{253,451 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRD_{\text{far-c}} \left( \frac{994.7 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \right. \\ \left. \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRD_{\text{far-a}} \left( \frac{676.4 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- **consumption of swine.** Table 13-51 of the 2011 Exposure Factors Handbook was used to derive the intakes for home-produced swine.

$$CDI_{\text{water-far-swine-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = \left( C_{\text{water}} \left( \frac{\text{pCi}}{\text{L}} \right) \times IFSW_{\text{far-adj}} \left( \frac{29,288 \text{ g}}{\text{year}} \right) \times CF_{\text{far-swine}} (1) \times \right. \\ \left. TF_{\text{swine}} \left( \frac{\text{day}}{\text{kg}} \right) \times Q_{\text{w-swine}} \left( \frac{11.4 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right) \right)$$

where:

$$IFE_{\text{far-adj}} \left( \frac{19,395 \text{ g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRE_{\text{far-c}} \left( \frac{31.7 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \right. \\ \left. \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRE_{\text{far-a}} \left( \frac{59.6 \text{ g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

The following consumption routes are provided in site-specific mode only and requires the user to enter their own data, as the tool only provides a transfer factor.

- **consumption of goat.** Please see table 1, in section 5 below, for sources of derived intakes for home-produced goat.



$$CDI_{\text{water-far-goat-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = \left( C_{\text{water}} \left( \frac{\text{pCi}}{\text{L}} \right) \times IFGO_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) \times CF_{\text{far-goat}} (1) \times \right. \\ \left. TF_{\text{goat}} \left( \frac{\text{day}}{\text{kg}} \right) \times Q_{\text{w-goat}} \left( \frac{3.81 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right) \right)$$

where:

$$IFGO_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRGO_{\text{far-c}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \right. \\ \left. \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRGO_{\text{far-a}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- consumption of goat milk. Please see table 1, in section 5 below, for sources of derived intakes for home-produced goat dairy.

$$CDI_{\text{water-far-goat-milk-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = \left( C_{\text{water}} \left( \frac{\text{pCi}}{\text{L}} \right) \times IFSH_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) \times CF_{\text{far-sheep}} (1) \times \right. \\ \left. TF_{\text{goat-milk}} \left( \frac{\text{day}}{\text{L milk}} \right) \times \rho_m \left( \frac{1.03 \text{ kg}}{1 \text{ L milk}} \right)^{-1} \times Q_{\text{w-goat-milk}} \left( \frac{8.75 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right) \right)$$

where:

$$IFSH_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSH_{\text{far-c}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \right. \\ \left. \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSH_{\text{far-a}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- consumption of sheep. Please see table 1, in section 5 below, for sources of derived intakes for home-produced goat.

$$CDI_{\text{water-far-sheep-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = \left( C_{\text{water}} \left( \frac{\text{pCi}}{\text{L}} \right) \times IFSM_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) \times CF_{\text{far-sheep-milk}} (1) \times \right. \\ \left. TF_{\text{sheep}} \left( \frac{\text{day}}{\text{kg}} \right) \times Q_{\text{w-sheep}} \left( \frac{5.25 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right) \right)$$

where:

$$IFSM_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSM_{\text{far-c}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \right. \\ \left. \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSM_{\text{far-a}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- consumption of sheep milk. Please see table 1, in section 5 below, for sources of derived intakes for home-produced sheep dairy.

$$CDI_{\text{water-far-sheep-milk-ing}} \left( \frac{\text{pCi}}{\text{year}} \right) = \left( C_{\text{water}} \left( \frac{\text{pCi}}{\text{L}} \right) \times IFSM_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) \times CF_{\text{far-sheep-milk}} (1) \times \right. \\ \left. TF_{\text{sheep-milk}} \left( \frac{\text{day}}{\text{L milk}} \right) \times \rho_m \left( \frac{1.03 \text{ kg}}{1 \text{ L milk}} \right)^{-1} \times Q_{\text{w-sheep-milk}} \left( \frac{10.4 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right) \right)$$

where:

$$IFSM_{\text{far-adj}} \left( \frac{\text{g}}{\text{year}} \right) = \left( \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSM_{\text{far-c}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \right. \\ \left. \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times IRSM_{\text{far-a}} \left( \frac{\text{g}}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right) \right)$$

where:


$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

Definitions of the input variables are in [Table 1](#).

#### 4.8.4 Farmer Consumption of Agricultural Products - Back-Calculated to Soil and Water

The farmer scenario should be considered an extension of the resident scenario and evaluate consumption of farm products for a subsistence farmer. Like the resident, the farmer scenario assumes the receptor will be exposed via consumption of home grown produce (100% of fruit and vegetables are from the farm). In addition to produce, 100% of consumption of the following are also considered to be from the farm: beef, milk, fish, swine, egg, and poultry. All feed and water (100%) for farm products is considered to have been grown on or provided from contaminated portions of the site. For these farm products, dose-based DCCs are provided for both soil and water, which may contribute contaminants to the products. The results are presented in an interactive graphic that shows the contribution from both sources. See section 4.8.4 for details. Also like the resident, age-adjusted intake equations were developed for all of the ingestion/consumption equations to account for changes in intake as the receptor ages.

#### Graphical Representation



**DCC Equations**

Results of back-calculating exposure to soil and water are presented in an interactive graph. See Section 4.8.4 for details.

- consumption of produce (fruits and vegetables)

y-INTERCEPT =

$$DCC_{\text{soil-far-produce-ing-tot}}(pCi/g) = \frac{1}{\sum_{i=1}^n \frac{1}{DCC_{\text{soil-far-produce-ing}}(pCi/g)_i}}$$

and:

x-INTERCEPT =

$$DCC_{\text{water-far-produce-ing-tot}}(pCi/L) = \frac{1}{\sum_{i=1}^n \frac{1}{DCC_{\text{water-far-produce-ing}}(pCi/L)_i}}$$

where:

n = total number of produce items included.

- consumption of eggs

$DCC_{\text{soil-far-egg-ing}} =$

$$y\text{-INTERCEPT} = \frac{DCC_{\text{far-egg-ing}}(pCi/g)}{TF_{\text{egg}}\left(\frac{\text{day}}{\text{kg}}\right) \times \left[ Q_{\text{p-poultry}}\left(\frac{0.2 \text{ kg}}{\text{day}}\right) \times f_{\text{p-poultry}}(1) \times f_{\text{s-poultry}}(1) \times (R_{\text{upp}} + R_{\text{es}}) \right] + \left[ Q_{\text{s-poultry}}\left(\frac{0.022 \text{ kg}}{\text{day}}\right) \times f_{\text{p-poultry}}(1) \right]}$$

and:

$DCC_{\text{water-far-egg-ing}}(pCi/L) =$

$$x\text{-INTERCEPT} = \frac{DCC_{\text{far-egg-ing}}(pCi/g)}{TF_{\text{egg}}\left(\frac{\text{day}}{\text{kg}}\right) \times Q_{\text{w-poultry}}\left(\frac{0.4 \text{ L}}{\text{day}}\right) \times \left(\frac{1 \text{ kg}}{1000 \text{ g}}\right)}$$

where:

$$SLOPE = \frac{-Q_{\text{w-poultry}}\left(\frac{0.4 \text{ L}}{\text{day}}\right)}{\left[ Q_{\text{p-poultry}}\left(\frac{0.2 \text{ kg}}{\text{day}}\right) \times f_{\text{p-poultry}}(1) \times f_{\text{s-poultry}}(1) \times (R_{\text{upp}} + R_{\text{es}}) \right] + \left[ Q_{\text{s-poultry}}\left(\frac{0.022 \text{ kg}}{\text{day}}\right) \times f_{\text{p-poultry}}(1) \right]}$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}}\left(\frac{pCi}{g\text{-dry plant}}\right); R_{\text{es}} = MLF_{\text{pasture}}\left(\frac{0.25 \text{ g-dry soil}}{g\text{-dry plant}}\right)$$

- consumption of poultry

DCC<sub>soil-far-poultry-ing</sub> =

$$y\text{-INTERCEPT} = \frac{DCC_{\text{far-poultry-ing}} (\text{pCi/g})}{TF_{\text{poultry}} \left( \frac{\text{day}}{\text{kg}} \right) \times \left[ \left( Q_{\text{p-poultry}} \left( \frac{0.2 \text{ kg}}{\text{day}} \right) \times f_{\text{p-poultry}}(t) \times f_{\text{s-poultry}}(t) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-poultry}} \left( \frac{0.022 \text{ kg}}{\text{day}} \right) \times f_{\text{p-poultry}}(t) \right) \right]}$$

and:

DCC<sub>water-far-poultry-ing</sub> (pCi/L) =

$$y\text{-INTERCEPT} = \frac{DCC_{\text{far-poultry-ing}} (\text{pCi/g})}{TF_{\text{poultry}} \left( \frac{\text{day}}{\text{kg}} \right) \times Q_{\text{w-poultry}} \left( \frac{0.4 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

where:

$$\text{SLOPE} = \frac{-Q_{\text{w-poultry}} \left( \frac{0.4 \text{ L}}{\text{day}} \right)}{\left( Q_{\text{p-poultry}} \left( \frac{0.2 \text{ kg}}{\text{day}} \right) \times f_{\text{p-poultry}}(t) \times f_{\text{s-poultry}}(t) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-poultry}} \left( \frac{0.022 \text{ kg}}{\text{day}} \right) \times f_{\text{p-poultry}}(t) \right)}$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{\text{pCi/g-dry plant}}{\text{pCi/g-dry soil}} \right); R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{\text{g-dry plant}} \right)$$

• consumption of beef

DCC<sub>soil-far-beef-ing</sub> =

$$y\text{-INTERCEPT} = \frac{DCC_{\text{far-beef-ing}} (\text{pCi/g})}{TF_{\text{beef}} \left( \frac{\text{day}}{\text{kg}} \right) \times \left[ \left( Q_{\text{p-beef}} \left( \frac{11.77 \text{ kg}}{\text{day}} \right) \times f_{\text{p-beef}}(t) \times f_{\text{s-beef}}(t) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-beef}} \left( \frac{0.5 \text{ kg}}{\text{day}} \right) \times f_{\text{p-beef}}(t) \right) \right]}$$

and:

DCC<sub>water-far-beef-ing</sub> (pCi/L) =

$$x\text{-INTERCEPT} = \frac{DCC_{\text{far-beef-ing}} (\text{pCi/g})}{TF_{\text{beef}} \left( \frac{\text{day}}{\text{kg}} \right) \times Q_{\text{w-beef}} \left( \frac{53 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

where:

$$\text{SLOPE} = \frac{-Q_{\text{w-beef}} \left( \frac{53 \text{ L}}{\text{day}} \right)}{\left( Q_{\text{p-beef}} \left( \frac{11.77 \text{ kg}}{\text{day}} \right) \times f_{\text{p-beef}}(t) \times f_{\text{s-beef}}(t) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-beef}} \left( \frac{0.5 \text{ kg}}{\text{day}} \right) \times f_{\text{p-beef}}(t) \right)}$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{\text{pCi/g-dry plant}}{\text{pCi/g-dry soil}} \right); R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{\text{g-dry plant}} \right)$$

• consumption of milk

DCC<sub>soil-far-dairy-ing</sub> =

$$y\text{-INTERCEPT} = \frac{DCC_{\text{far-dairy-ing}} (\text{pCi/g})}{TF_{\text{dairy}} \left( \frac{\text{day}}{\text{L milk}} \right) \times \rho_m \left( \frac{1.03 \text{ kg}}{1 \text{ L milk}} \right)^{-1} \times \left[ \left( Q_{\text{p-dairy}} \left( \frac{20.3 \text{ kg}}{\text{day}} \right) \times f_{\text{p-dairy}}(t) \times f_{\text{s-dairy}}(t) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-dairy}} \left( \frac{0.4 \text{ kg}}{\text{day}} \right) \times f_{\text{p-dairy}}(t) \right) \right]}$$

and:

DCC<sub>water-far-dairy-ing</sub> (pCi/L) =

$$x\text{-INTERCEPT} = \frac{DCC_{\text{far-dairy-ing}} (\text{pCi/g})}{TF_{\text{dairy}} \left( \frac{\text{day}}{\text{L milk}} \right) \times \rho_m \left( \frac{1.03 \text{ kg}}{1 \text{ L milk}} \right)^{-1} \times Q_{\text{w-dairy}} \left( \frac{92 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

where:

$$\text{SLOPE} = \frac{-Q_{\text{w-dairy}} \left( \frac{92 \text{ L}}{\text{day}} \right)}{\left( Q_{\text{p-dairy}} \left( \frac{20.3 \text{ kg}}{\text{day}} \right) \times f_{\text{p-dairy}}(t) \times f_{\text{s-dairy}}(t) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-dairy}} \left( \frac{0.4 \text{ kg}}{\text{day}} \right) \times f_{\text{p-dairy}}(t) \right)}$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{\text{pCi/g-dry plant}}{\text{pCi/g-dry soil}} \right); R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{\text{g-dry plant}} \right)$$

• consumption of swine

DCC<sub>soil-far-swine-ing</sub> =

$$y\text{-INTERCEPT} = \frac{DCC_{\text{far-swine-ing}} (pCi/g)}{TF_{\text{swine}} \left( \frac{\text{day}}{\text{kg}} \right) \times \left[ \left( Q_{\text{p-swine}} \left( \frac{4.7 \text{ kg}}{\text{day}} \right) \times f_{\text{p-swine}} (1) \times f_{\text{s-swine}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-swine}} \left( \frac{0.37 \text{ kg}}{\text{day}} \right) \times f_{\text{p-swine}} (1) \right) \right]}$$

and:

DCC<sub>water-far-swine-ing</sub> (pCi/L) =

$$x\text{-INTERCEPT} = \frac{DCC_{\text{far-swine-ing}} (pCi/g)}{TF_{\text{swine}} \left( \frac{\text{day}}{\text{kg}} \right) \times Q_{\text{w-swine}} \left( \frac{11.4 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

where:

$$\text{SLOPE} = \frac{-Q_{\text{w-swine}} \left( \frac{11.4 \text{ L}}{\text{day}} \right)}{\left( Q_{\text{p-swine}} \left( \frac{4.7 \text{ kg}}{\text{day}} \right) \times f_{\text{p-swine}} (1) \times f_{\text{s-swine}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-swine}} \left( \frac{0.37 \text{ kg}}{\text{day}} \right) \times f_{\text{p-swine}} (1) \right)}$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{pCi / \text{g-dry plant}}{pCi / \text{g-dry soil}} \right); R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{\text{g-dry plant}} \right)$$

• consumption of fish

DCC<sub>soil-far-fish-ing</sub> =

$$y\text{-INTERCEPT} = \frac{DCC_{\text{far-fish-ing}} (pCi/g) \times Kd \left( \frac{\text{L}}{\text{kg}} \right)}{BCF \left( \frac{\text{L}}{\text{kg}} \right)}$$

and:

DCC<sub>water-far-fish-ing</sub> (pCi/L) =

$$x\text{-INTERCEPT} = \frac{DCC_{\text{far-fish-ing}} (pCi/g)}{BCF \left( \frac{\text{L}}{\text{kg}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

where:

$$\text{SLOPE} = \frac{-Kd \left( \frac{\text{L}}{\text{kg}} \right)}{1000}$$

The following consumption routes are provided in site-specific mode only and require the user to enter their own data, as the tool only provides a transfer factor.

• consumption of goat

PRG<sub>soil-far-goat-ing</sub> =

$$y\text{-INTERCEPT} = \frac{DCC_{\text{far-goat-ing}} (pCi/g)}{TF_{\text{goat}} \left( \frac{\text{day}}{\text{kg}} \right) \times \left[ \left( Q_{\text{p-goat}} \left( \frac{1.27 \text{ kg}}{\text{day}} \right) \times f_{\text{p-goat}} (1) \times f_{\text{s-goat}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-goat}} \left( \frac{0.23 \text{ kg}}{\text{day}} \right) \times f_{\text{p-goat}} (1) \right) \right]}$$

and:

PRG<sub>water-far-goat-ing</sub> (pCi/L) =

$$x\text{-INTERCEPT} = \frac{DCC_{\text{far-goat-ing}} (pCi/g)}{TF_{\text{goat}} \left( \frac{\text{day}}{\text{kg}} \right) \times Q_{\text{w-goat}} \left( \frac{3.81 \text{ L}}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

where:

$$\text{SLOPE} = \frac{-Q_{\text{w-goat}} \left( \frac{3.81 \text{ L}}{\text{day}} \right)}{\left( Q_{\text{p-goat}} \left( \frac{1.27 \text{ kg}}{\text{day}} \right) \times f_{\text{p-goat}} (1) \times f_{\text{s-goat}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-goat}} \left( \frac{0.23 \text{ kg}}{\text{day}} \right) \times f_{\text{p-goat}} (1) \right)}$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{pCi / \text{g-dry plant}}{pCi / \text{g-dry soil}} \right); R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{\text{g-dry plant}} \right)$$

• consumption of goat milk

PRG<sub>soil-far-goat-milk-ing</sub> =

$$y\text{-INTERCEPT} = \frac{DCC_{\text{far-goat-milk-ing}} (pCi/g)}{TF_{\text{goat-milk}} \left( \frac{\text{day}}{L \text{ milk}} \right) \times p_m \left( \frac{1.03 \text{ kg}}{1 L \text{ milk}} \right)^{-1} \times \left[ \left( Q_{\text{p-goat-milk}} \left( \frac{1.59 \text{ kg}}{\text{day}} \right) \times f_{\text{p-goat-milk}} (1) \times f_{\text{s-goat-milk}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-goat-milk}} \left( \frac{0.29 \text{ kg}}{\text{day}} \right) \times f_{\text{p-goat-milk}} (1) \right) \right]}$$

and:

PRG<sub>water-far-goat-milk-ing</sub> (pCi/L) =

$$x\text{-INTERCEPT} = \frac{DCC_{\text{far-goat-milk-ing}} (pCi/g)}{TF_{\text{goat-milk}} \left( \frac{\text{day}}{L \text{ milk}} \right) \times p_m \left( \frac{1.03 \text{ kg}}{1 L \text{ milk}} \right)^{-1} \times Q_{\text{w-goat-milk}} \left( \frac{8.75 L}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

where:

$$\text{SLOPE} = \frac{-Q_{\text{w-goat-milk}} \left( \frac{8.75 L}{\text{day}} \right)}{\left( Q_{\text{p-goat-milk}} \left( \frac{1.59 \text{ kg}}{\text{day}} \right) \times f_{\text{p-goat-milk}} (1) \times f_{\text{s-goat-milk}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-goat-milk}} \left( \frac{0.29 \text{ kg}}{\text{day}} \right) \times f_{\text{p-goat-milk}} (1) \right)}$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{pCi / g \text{-dry plant}}{pCi / g \text{-dry soil}} \right); R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{g \text{-dry plant}} \right)$$

• consumption of sheep

PRG<sub>soil-far-sheep-ing</sub> =

$$y\text{-INTERCEPT} = \frac{DCC_{\text{far-sheep-ing}} (pCi/g)}{TF_{\text{sheep}} \left( \frac{\text{day}}{\text{kg}} \right) \times \left[ \left( Q_{\text{p-sheep}} \left( \frac{1.75 \text{ kg}}{\text{day}} \right) \times f_{\text{p-sheep}} (1) \times f_{\text{s-sheep}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-sheep}} \left( \frac{0.32 \text{ kg}}{\text{day}} \right) \times f_{\text{p-sheep}} (1) \right) \right]}$$

and:

PRG<sub>water-far-sheep-ing</sub> (pCi/L) =

$$x\text{-INTERCEPT} = \frac{DCC_{\text{far-sheep-ing}} (pCi/g)}{TF_{\text{sheep}} \left( \frac{\text{day}}{\text{kg}} \right) \times Q_{\text{w-sheep}} \left( \frac{5.25 L}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

where:

$$\text{SLOPE} = \frac{-Q_{\text{w-sheep}} \left( \frac{5.25 L}{\text{day}} \right)}{\left( Q_{\text{p-sheep}} \left( \frac{1.75 \text{ kg}}{\text{day}} \right) \times f_{\text{p-sheep}} (1) \times f_{\text{s-sheep}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-sheep}} \left( \frac{0.32 \text{ kg}}{\text{day}} \right) \times f_{\text{p-sheep}} (1) \right)}$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{pCi / g \text{-dry plant}}{pCi / g \text{-dry soil}} \right); R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{g \text{-dry plant}} \right)$$

• consumption of sheep milk

PRG<sub>soil-far-sheep-milk-ing</sub> =

$$y\text{-INTERCEPT} = \frac{DCC_{\text{far-sheep-milk-ing}} (pCi/g)}{TF_{\text{sheep-milk}} \left( \frac{\text{day}}{L \text{ milk}} \right) \times p_m \left( \frac{1.03 \text{ kg}}{1 L \text{ milk}} \right)^{-1} \times \left[ \left( Q_{\text{p-sheep-milk}} \left( \frac{3.15 \text{ kg}}{\text{day}} \right) \times f_{\text{p-sheep-milk}} (1) \times f_{\text{s-sheep-milk}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-sheep-milk}} \left( \frac{0.57 \text{ kg}}{\text{day}} \right) \times f_{\text{p-sheep-milk}} (1) \right) \right]}$$

and:

PRG<sub>water-far-sheep-milk-ing</sub> (pCi/L) =

$$x\text{-INTERCEPT} = \frac{DCC_{\text{far-sheep-milk-ing}} (pCi/g)}{TF_{\text{sheep-milk}} \left( \frac{\text{day}}{L \text{ milk}} \right) \times p_m \left( \frac{1.03 \text{ kg}}{1 L \text{ milk}} \right)^{-1} \times Q_{\text{w-sheep-milk}} \left( \frac{10.4 L}{\text{day}} \right) \times \left( \frac{1 \text{ kg}}{1000 \text{ g}} \right)}$$

where:

$$\text{SLOPE} = \frac{-Q_{\text{w-sheep-milk}} \left( \frac{10.4 L}{\text{day}} \right)}{\left( Q_{\text{p-sheep-milk}} \left( \frac{3.15 \text{ kg}}{\text{day}} \right) \times f_{\text{p-sheep-milk}} (1) \times f_{\text{s-sheep-milk}} (1) \times (R_{\text{upp}} + R_{\text{es}}) \right) + \left( Q_{\text{s-sheep-milk}} \left( \frac{0.57 \text{ kg}}{\text{day}} \right) \times f_{\text{p-sheep-milk}} (1) \right)}$$

where:

$$R_{\text{upp}} = Bv_{\text{dry}} \left( \frac{pCi / g \text{-dry plant}}{pCi / g \text{-dry soil}} \right); R_{\text{es}} = MLF_{\text{pasture}} \left( \frac{0.25 \text{ g-dry soil}}{g \text{-dry plant}} \right)$$

Results of back-calculating exposure to soil and water are presented in an interactive graph. See Section 4.8.4 for details.


Definitions of the input variables are in [Table 1](#).

### 4.8.5 Farmer Air

This receptor spends most, if not all, of the day at home except for the hours spent at work. The activities for this receptor involve typical home making chores (cooking, cleaning, and laundering) as well as gardening. This receptor is assumed to be exposed to contaminants via the following pathways: inhalation of ambient air and external radiation from contaminants in ambient air. To take into account the different inhalation rates for children and adults, age-adjusted intake equations were developed to account for changes in intake as the receptor ages.

Two ambient air exposure conditions are offered for this scenario. The first scenario includes a half-life decay function and the second scenario does not. In situations where the contaminant in the air is not being replenished (e.g., an accidental one-time air release from a factory), equations for the first scenario should be used. In situations where the contaminant in the air has a continual source (e.g., indoor radon from radium in the soil or an operating factory or landfill cap), equations for the second scenario should be used.

### Graphical Representation



### DCC Equations

The resident ambient air land use equation, presented here, contains the following exposure routes without half-life decay:

- inhalation

$$DCC_{\text{far-air-inh-no decay}} \left( \frac{\text{pCi}}{\text{m}^3} \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_i \left( \frac{\text{mrem}}{\text{pCi}} \right) \times IFA_{\text{far-adj}} \left( \frac{6,475 \text{ m}^3}{\text{year}} \right)}$$

where:

$$IFA_{\text{far-adj}} \left( \frac{6,475 \text{ m}^3}{\text{year}} \right) = \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times ET_{\text{far-c}} \left( \frac{24 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{far-c}} \left( \frac{10 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times ET_{\text{far-a}} \left( \frac{24 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{far-a}} \left( \frac{20 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- external exposure to ionizing radiation

$$DCC_{\text{far-air-sub-no decay}} \left( \frac{\text{pCi}}{\text{m}^3} \right) = \frac{DL \left( \frac{\text{mrem}}{\text{year}} \right)}{DCF_{\text{sub}} \left( \frac{\text{mrem/year}}{\text{pCi/m}^3} \right) \times EF_{\text{far}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{\text{far}} \left( \frac{24 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_a (1.0)}$$

- total

$$DCC_{\text{far-air-tot-no decay}} \left( \frac{\text{pCi}}{\text{m}^3} \right) = \frac{1}{\frac{1}{DCC_{\text{far-air-inh-no decay}}} + \frac{1}{DCC_{\text{far-air-sub-no decay}}}}$$

Definitions of the input variables are in [Table 1](#).

### CDI Equations

The resident ambient air land use equation, presented here, contains the following exposure routes without half-life decay:

- inhalation

$$CDI_{\text{far-air-inh-no decay}} \left( \frac{\text{pCi}}{\text{m}^3} \right) = C_{\text{air}} \left( \frac{\text{pCi}}{\text{m}^3} \right) \times IFA_{\text{far-adj}} \left( \frac{6,475 \text{ m}^3}{\text{year}} \right)$$

where:

$$IFA_{\text{far-adj}} \left( \frac{6,475 \text{ m}^3}{\text{year}} \right) = \left( EF_{\text{far-c}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times ET_{\text{far-c}} \left( \frac{24 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{far-c}} \left( \frac{10 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{far-c}} (0.15) \right) + \left( EF_{\text{far-a}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times ET_{\text{far-a}} \left( \frac{24 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times IRA_{\text{far-a}} \left( \frac{20 \text{ m}^3}{\text{day}} \right) \times AAF_{\text{far-a}} (0.85) \right)$$

where:

$$AAF_{\text{far-c}} (0.15) = \left( \frac{ED_{\text{far-c}} (6 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right) \text{ and: } AAF_{\text{far-a}} (0.85) = \left( \frac{ED_{\text{far-a}} (34 \text{ years})}{ED_{\text{far}} (40 \text{ years})} \right)$$

- external exposure to ionizing radiation

$$CDI_{\text{far-air-sub-no decay}} \left( \frac{\text{pCi}}{\text{m}^3} \right) = C_{\text{air}} \left( \frac{\text{pCi}}{\text{m}^3} \right) \times EF_{\text{far}} \left( \frac{350 \text{ days}}{\text{year}} \right) \times \left( \frac{1 \text{ year}}{365 \text{ days}} \right) \times ET_{\text{far}} \left( \frac{24 \text{ hours}}{\text{day}} \right) \times \left( \frac{1 \text{ day}}{24 \text{ hours}} \right) \times GSF_a (1.0)$$

Definitions of the input variables are in [Table 1](#).


## 4.9 Soil to Groundwater

The soil to groundwater scenario was developed to identify concentrations in soil that have the potential to contaminate groundwater above



dose-based DCCs or MCLs. Migration of contaminants from soil to groundwater can be envisioned as a two-stage process: (1) release of contaminant from soil to soil leachate and (2) transport of the contaminant through the underlying soil and aquifer to a receptor well. The soil to groundwater scenario considers both of these fate and transport mechanisms. First, the acceptable groundwater concentration is multiplied by a dilution factor to obtain a target leachate concentration. For example, if the dilution factor is 10 and the MCL is 0.05 mg/L, the target soil leachate concentration would be 0.5 mg/L. The partition equation (presented in the Soil Screening Guidance for Radionuclides documents) is then used to calculate the total soil concentration corresponding to this soil leachate concentration.

Graphical Representation



The SSL methodology was designed for use during the early stages of a site evaluation when information about subsurface conditions may be limited. Because of this constraint, the methodology is based on protective, simplifying assumptions about the release and transport of contaminants in the subsurface.

Equations

The user has the option to choose from two calculation methods. The first method employs the default partitioning equation for migration to groundwater. The dilution factor defaults to 20 for a 0.5-acre source. If all the parameters needed to calculate a dilution factor are available, then use Method 2 (mass-limit equation for migration to groundwater).

Method 1. Partitioning Equation for Migration to Groundwater

$$SSL\left(\frac{pCi}{g}\right) = C_{water}\left(\frac{pCi}{L}\right) \times \left(\frac{kg}{1000g}\right) \times \left(K_d\left(\frac{L}{kg}\right) + \frac{\theta_w\left(\frac{L_{water}}{L_{soil}}\right)}{\rho_b\left(\frac{kg}{L}\right)}\right)$$

where:  
 $C_{water} = MCL \times DAF$   
or  
 $C_{water} = PRG \times DAF$

Method 2. Mass-Limit Equation for Migration to Groundwater

$$SSL\left(\frac{pCi}{g}\right) = \frac{C_{water}\left(\frac{pCi}{L}\right) \times I\left(\frac{m}{year}\right) \times ED_{gw}(70\text{ years})}{\rho_b\left(\frac{kg}{L}\right) \times d_s(m) \times \left(\frac{1000g}{kg}\right)}$$

where:  
 $C_{water} = MCL \times DAF$   
or  
 $C_{water} = PRG \times DAF$

Then calculate the dilution factor using this equation.

$$DAF = 1 + \frac{\left(K\left(\frac{m}{year}\right) \times I\left(\frac{m}{m}\right) \times d(m)\right)}{\left(I\left(\frac{m}{year}\right) \times L(m)\right)}$$

Definitions of the input variables are in [Table 1](#).

4.10 Supporting Equations and Parameter Discussion

There are five parts of the above land use equations that require further explanation: two inhalation variables [the particulate emission factor (PEF) and the volatilization factor (VF)], the radionuclide decay constant (λ), the area correction factor (ACF), the outdoor soil gamma

shielding factor  $GCF_0$ ), and the groundwater transport portion of the equations involving the soil to water partition coefficient ( $K_d$ ).

4.10.1 Particulate Emission Factor (PEF) and Volatilization Factor (VF)

Inhalation of isotopes adsorbed to respirable particles (PM10) was assessed using a default PEF equal to  $1.36 \times 10^9 \text{ m}^3/\text{kg}$ . This equation relates the contaminant concentration in soil with the concentration of respirable particles in the air due to fugitive dust emissions from contaminated soils. The generic PEF was derived using default values that correspond to a receptor point concentration of approximately  $0.76 \text{ ug}/\text{m}^3$ . The relationship is derived by Cowherd (1985) for a rapid assessment procedure applicable to a typical hazardous waste site, where the surface contamination provides a relatively continuous and constant potential for emission over an extended period of time (e.g., years). This represents an annual average emission rate based on wind erosion that should be compared with chronic health criteria; it is not appropriate for evaluating the potential for more acute exposures. Definitions of the input variables are in [Table 1](#).

With the exception of a few radionuclides, the PEF does not appear to significantly affect most DCCs. The equation forms the basis for deriving a generic PEF for the inhalation pathway. For more details regarding specific parameters used in the PEF model, refer to [Soil Screening Guidance for Radionuclides: Technical Background Document](#). The use of alternate values on a specific site should be justified and presented in an Administrative Record if considered in CERCLA remedy selection.

$$PEF \left( \frac{\text{m}^3_{\text{air}}}{\text{kg}_{\text{soil}}} \right) = \frac{Q}{C_{\text{wind}}} \left( \frac{\frac{g}{\text{m}^2 \cdot s}}{\left( \frac{\text{kg}}{\text{m}^3} \right)} \right) \times \frac{3,600 \left( \frac{s}{\text{hour}} \right)}{0.036 \times (1 - V) \times \left( \frac{U_m \left( \frac{m}{s} \right)}{U_t \left( \frac{m}{s} \right)} \right)^3 \times F(x)}$$

where:

$$\frac{Q}{C_{\text{wind}}} = A \times \exp \left[ \frac{\left( \ln A_s (\text{acre}) - B \right)^2}{C} \right]$$

and:

if  $x < 2$ ,  $F(x) = 1.91207 - 0.0278085 x + 0.48113 x^2 - 1.09871 x^3 + 0.335341 x^4$

if  $x \geq 2$ ,  $F(x) = 0.18 \left( 8x^3 + 12x \right) e^{-x^2}$

where:

$$x = 0.886 \times \left( \frac{U_t}{U_m} \right)$$

**Note: the generic PEF evaluates wind-borne emissions and does not consider dust emissions from traffic or other forms of mechanical disturbance that could lead to greater emissions than assumed here.**

EPA derived a default volatilization factor (VF) value of  $17 \text{ m}^3/\text{kg}$  for tritium. The VF replaces the PEF in the DCC equations when tritium is being addressed. This VF value is based on a steady-state model that assumes, on average, tritium in soil pore water and tritium in air (as tritiated water vapor) will be distributed in the environment in proportion to the average water content in soil and air. EPA assumes a mean atmospheric humidity of 6 grams of water per cubic meter of air ( $\text{g}/\text{m}^3$ ) nationwide ([Etnier 1980](#)), and an average soil moisture content of 10% (i.e., 100 grams of water per kilogram of soil). Given these assumptions, EPA calculates the VF term for tritium as:

$$VFH-3 = 100 \text{ g H}_2\text{O}/\text{kg soil} / 6 \text{ g H}_2\text{O}/\text{m}^3 \text{ air}$$
$$= 17 \text{ m}^3 \text{ air}/\text{kg soil}$$
$$= 17 \text{ m}^3/\text{kg}$$

EPA believes that this value is appropriate for the average case, both outdoors and indoors; However, site managers can derive a site-specific VF term for tritium that may be more appropriate for a specific site, considering local atmospheric humidity and soil moisture content.

4.10.2 Standard Unpaved Road Vehicle Traffic Particulate Emission Factor (PEF<sub>sc</sub>)

The equation to calculate the subchronic particulate emission factor (PEF<sub>sc</sub>) is significantly different from the resident and non-resident PEF equations. The PEF<sub>sc</sub> focuses exclusively on emissions from truck traffic on unpaved roads, which typically contribute the majority of dust emissions during construction. This equation requires parameter estimates such as the number of days with at least 0.01 inches of rainfall, the mean vehicle weight, and the sum of fleet vehicle distance traveled during construction.

The number of days with at least 0.01 inches of rainfall can be estimated using Exhibit 5-2 in the [supplemental soil screening guidance](#). Mean vehicle weight (W) can be estimated by assuming the numbers and weights of different types of vehicles. For example, assuming that the daily unpaved road traffic consists of 20 two-ton cars and 10 twenty-ton trucks, the mean vehicle weight would be:

$$W = [(20 \text{ cars} \times 2 \text{ tons/car}) + (10 \text{ trucks} \times 20 \text{ tons/truck})]/30 \text{ vehicles} = 8 \text{ tons}$$

The sum of the fleet vehicle kilometers traveled during construction ( $\sum \text{VKT}$ ) can be estimated based on the size of the area of surface soil contamination, assuming the configuration of the unpaved road, and the amount of vehicle traffic on the road. For example, if the area of surface soil contamination is 0.5 acres (or  $2,024 \text{ m}^2$ ) and one assumes that this area is configured as a square with the unpaved road segment dividing the square evenly, the road length would be equal to the square root of  $2,024 \text{ m}^2$  ( $45 \text{ m}$  or  $0.045 \text{ km}$ ). Assuming that each vehicle travels the length of the road once per day, 5 days per week for a total of 6 months, the total fleet vehicle kilometers traveled would be:

$$\sum \text{VKT} = 30 \text{ vehicles} \times 0.045 \text{ km/day} \times (52 \text{ weeks/year} / 2) \times 5 \text{ days/week} = 175.5 \text{ km}$$

$$\begin{aligned} \text{PEF}_{\text{sc}} \left( \frac{\text{m}^3_{\text{air}}}{\text{kg}_{\text{soil}}} \right) &= \frac{Q}{C_{\text{sr}}} \left( \frac{\frac{\text{g}}{\text{m}^2 \cdot \text{s}}}{\left( \frac{\text{kg}}{\text{m}^3} \right)} \right) \times \frac{1}{F_D} \times \left[ \frac{T_t (\text{s}) \times A_R (\text{m}^2)}{2.6 \times \left( \frac{\text{s}}{12} \right)^{0.8} \times \left( \frac{W(\text{tons})}{3} \right)^{0.4} \times \left( \frac{365 \left( \frac{\text{days}}{\text{year}} \right) - p \left( \frac{\text{days}}{\text{year}} \right)}{365 \left( \frac{\text{days}}{\text{year}} \right)} \right) \times 281.9 \times \Sigma \text{VKT}(\text{km})} \right. \\ &\quad \left. \frac{\left( \frac{\text{M}_{\text{dry}}}{0.2} \right)^{0.3}}{\left( \frac{\text{kg}}{\text{m}^3} \right)} \right] \\ \frac{Q}{C_{\text{sr}}} \left( \frac{\frac{\text{g}}{\text{m}^2 \cdot \text{s}}}{\left( \frac{\text{kg}}{\text{m}^3} \right)} \right) &= A \times \exp \left[ \frac{(\ln A_s (\text{acre}) - B)^2}{C} \right] \\ A_R (\text{m}^2) &= L_R (\text{ft}) \times W_R (20 \text{ feet}) \times 0.092903 \left( \frac{\text{m}^2}{\text{feet}^2} \right) \\ W (\text{tons}) &= \frac{\left( \text{number of cars} \times \frac{\text{tons}}{\text{car}} + \text{number of trucks} \times \frac{\text{tons}}{\text{truck}} \right)}{\text{total vehicles}} \\ \Sigma \text{VKT} (\text{km}) &= \text{total vehicles} \times \text{distance} \left( \frac{\text{km}}{\text{day}} \right) \times \text{EW}_{\text{cw}} \left( \frac{\text{weeks}}{\text{year}} \right) \times \text{DW}_{\text{cw}} \left( \frac{\text{days}}{\text{week}} \right) \\ T_t (7200000 \text{ s}) &= \text{ED}_{\text{cw}} (1 \text{ years}) \times \text{EF}_{\text{cw}} \left( \frac{250 \text{ days}}{\text{year}} \right) \times \text{ET}_{\text{cw}} \left( \frac{8 \text{ hours}}{\text{day}} \right) \times \left( \frac{3600 \text{ s}}{\text{hour}} \right) \\ F_D (0.18584) &= 0.1852 + \left( 5.3537 / t_c \right) + \left( -9.6318 / t_c^2 \right) \\ t_c (8400 \text{ hours}) &= \text{ED}_{\text{cw}} (1 \text{ years}) \times \text{EW}_{\text{cw}} \left( \frac{50 \text{ weeks}}{\text{year}} \right) \times \left( \frac{7 \text{ days}}{\text{week}} \right) \times \left( \frac{24 \text{ hours}}{\text{day}} \right) \end{aligned}$$

4.10.3 Other Construction Activities Particulate Emission Factor (PEF'<sub>sc</sub>)

Other than emissions from unpaved road traffic, the construction worker may also be exposed to particulate matter emissions (PEF'<sub>sc</sub>) from wind erosion; excavation soil dumping, dozing, grading, and tilling; or similar operations. These operations may occur separately or concurrently, and the duration of each operation may be different. For these reasons, the total unit mass emitted from each operation is calculated separately, and the sum is normalized over the entire area of contamination and over the entire time during which construction activities take place. Equation E-26 in the [supplemental soil screening guidance](#) was used.

$$PEF'_{sc} \left( \frac{m^3}{kg_{soil}} \right) = \frac{Q}{C_{sa}} \left( \frac{\left( \frac{g}{m^2-s} \right)}{\left( \frac{kg}{m^3} \right)} \right) \times \frac{1}{F_D} \times \frac{1}{<I_T> \left( \frac{g}{m^2-s} \right)}$$

where:

$$\frac{Q}{C_{sa}} \left( \frac{\left( \frac{g}{m^2-s} \right)}{\left( \frac{kg}{m^3} \right)} \right) = A \times \exp \left[ \frac{(\ln A_c (acre) \cdot B)^2}{C} \right]$$

$$<I_T> \left( \frac{g}{m^2-s} \right) = \frac{M_{wind}^{PC}(g) + M_{excav}(g) + M_{doz}(g) + M_{grade}(g) + M_{till}(g)}{A_{surf}(m^2) \times T_t(s)}$$

$$M_{wind}^{PC}(g) = 0.036 \times (1-V) \times \left( \frac{U_m \left( \frac{m}{s} \right)}{U_t \left( \frac{m}{s} \right)} \right)^3 \times F(x) \times A_{surf}(m^2) \times ED(years) \times 8760 \left( \frac{hours}{year} \right)$$

$$M_{excav}(g) = 0.35 \times 0.0016 \times \frac{\left( \frac{U_m \left( \frac{m}{s} \right)}{2.2} \right)^{1.3}}{\left( \frac{M_{m-excav}(\%)}{2} \right)^{1.4}} \times \rho_{soil} \left( \frac{Mg}{m^3} \right) \times A_{excav}(m^2) \times d_{excav}(m) \times N_{A-dump} \times 1000 \left( \frac{g}{kg} \right)$$

$$M_{doz}(g) = 0.75 \times \frac{0.45 \times s_{doz}(\%)}{(M_{m-doz}(\%))^{1.4}} \times \frac{\Sigma VKT_{doz}(km)}{S_{doz} \left( \frac{km}{hour} \right)} \times 1000 \left( \frac{g}{kg} \right)$$

$$M_{grade}(g) = 0.60 \times 0.0056 \times S_{grade} \left( \frac{km}{hour} \right)^{2.0} \times \Sigma VKT_{grade}(km) \times 1000 \left( \frac{g}{kg} \right)$$

and:

$$M_{till}(g) = 1.1 \times s_{till}(\%)^{0.6} \times A_{c-till}(acres) \times 4047 \left( \frac{m^2}{acre} \right) \times 10^{-4} \left( \frac{ha}{m^2} \right) \times 1000 \left( \frac{g}{kg} \right) \times N_{A-till}$$

where:

$$\Sigma VKT_{grade}(km) = A_{c-grade}(acres) \times 4047 \left( \frac{m^2}{acre} \right) \times \frac{1}{B_{l-grade}(m)} \times \frac{1}{1000 \left( \frac{m}{km} \right)} \times N_{A-grade}$$

where:

$$\Sigma VKT_{doz}(km) = A_{c-doz}(acres) \times 4047 \left( \frac{m^2}{acre} \right) \times \frac{1}{B_{l-doz}(m)} \times \frac{1}{1000 \left( \frac{m}{km} \right)} \times N_{A-doz}$$

$$T_t(7200000 s) = ED_{cw}(1 years) \times EF_{cw} \left( \frac{250 days}{year} \right) \times ET_{cw} \left( \frac{8 hours}{day} \right) \times \left( \frac{3600 s}{hour} \right)$$

$$F_D(0.18584) = 0.1852 + (5.3537 / t_c) + (-9.6318 / t_c^2)$$

$$t_c(8400 hours) = ED_{cw}(1 years) \times EW_{cw} \left( \frac{50 weeks}{year} \right) \times \left( \frac{7 days}{week} \right) \times \left( \frac{24 hours}{day} \right)$$

and:

$$\text{if } x < 2, F(x) = 1.91207 - 0.0278085x + 0.48113x^2 - 1.09871x^3 + 0.335341x^4$$

$$\text{if } x \geq 2, F(x) = 0.18(8x^3 + 12x)e^{(-x^2)}$$

where:

$$x = 0.886 \times \left( \frac{U_t}{U_m} \right)$$

#### 4.10.4 Radionuclide Decay Constant (λ)

The decay constant term (λ), which is based on the half-life of the isotope, is used for some media in nearly all land uses. λ = 0.693/half-life in years (where 0.693=ln(2)). The term (1 - e<sup>-λt</sup>) takes into account the number of half-lives that will occur within the exposure duration to calculate an appropriate value. For the secular equilibrium DCC output option, decay is not used. In most cases, site-specific analytical data should be used to establish the actual degree of equilibrium between each parent radionuclide and its decay products in each media sampled. In the absence of empirical data, however, the secular equilibrium DCCs will provide a protective screening value. Definitions of the input variables are in [Table 1](#).

#### 4.10.5 Area Correction Factor


The RAGS/HHEM Part B model assumes that an individual is exposed to a source geometry that is effectively an infinite slab. The concept of an infinite slab means that the thickness of the contaminated zone and its aerial extent are so large that it behaves as if it were infinite in its physical dimensions. In practice, soil contaminated to a depth greater than about 15 cm and with an aerial extent greater than about 1,000 m<sup>2</sup> will create a radiation field comparable to that of an infinite slab (U.S. EPA. 2000a).

To accommodate the fact that in most residential settings the assumption of an infinite slab source will result in overly protective DCCs, an adjustment for source area is considered to be an important modification to the RAGS/HHEM Part B model. Thus, an area correction factor (ACF) has been added to the calculation of recommended DCCs. For the 2-D exposure models addressing finite areas, the ACF is made variable by isotope and area for site-specific analysis. In addition, ACFs are now available for all alternate soil analysis source depths (ground plane, 1 cm, 5 cm, and 15 cm source volumes as well as infinite source volume). This calculator allows the user to select from 19 different soil area sizes. If the default mode is selected, the ACF from the most protective area size is selected. If site-specific mode is selected, the user must select the source area. For further information on the derivation of the isotope-specific/area-specific ACF values for 2-D areas, see the [Center for Radiation Protection Knowledge ACF report and appendix](#) containing +D and +E values. For the calculation of area correction factors, a standard soil density of 1.6 g/cm<sup>3</sup> has been used. For a description of other EPA default ACF values that predate this guidance, follow the link [here](#).

4.10.6 Gamma Shielding Factor

DCCs in this guidance are calculated without any shielding between the receptor and the source (soil). In this case, a default soil gamma shielding factor for outdoor exposure to ionizing radiation (GSF<sub>o</sub>) is established at 1.0 (0% shielding). It is common to have some shielding (soil cover) over contaminated soil. For site-specific mode, the user must select a soil cover depth. Due to shielding, covering the contaminated area with soil will produce lower dose and risk coefficients than are stated in the Federal Guidance Report (FGR) 12 and 13. Therefore, gamma shielding factors are needed to apply the published EPA dose values to the buried contamination scenarios. Outdoor gamma shielding factors (GSF<sub>o</sub>) are derived by modeling various thicknesses of clean soil covering ground soil contamination. The gamma shielding factor is defined as the ratio of the contamination dose compared to the unshielded surface source in soil. The MCNP output was used to derive kerma values one meter above the soil surface for various scenarios ranging from 0 cm soil cover to 100 cm soil cover in 10 cm increments while using source thicknesses of ground plane, 1, 5, and 15 cm source volumes as well as infinite source volume. Radioisotopes published in [ICRP 107](#) were considered, along with decay chains of several radioisotopes. The [Center for Radiation Protection Knowledge](#) has provided GSF<sub>o</sub> values [here](#) and an [appendix](#) containing +D and +E values. Additional source depth-specific gamma soil shielding factors (GSF) are now given for cover depths of 2 to 10 meters. The values are presented in this [appendix](#).

- A default gamma shielding factor for indoor exposure to ionizing radiation (GSF<sub>i</sub>) is established at 0.4 (60% shielding).
- GSF<sub>o</sub> depends on soil cover depth and is shown in equation images as GSF<sub>o-ext-sv</sub>, GSF<sub>o-ext-1cm</sub>, GSF<sub>o-ext-5cm</sub>, GSF<sub>o-ext-15cm</sub>, & GSF<sub>o-ext-gp</sub>.
- In the resident, farmer, and indoor worker soil external exposure equations, GSF<sub>i-total</sub> is applied to account for the gamma shielding provided by clean soil cover and the building subfloor. GSF<sub>i-total</sub> = GSF<sub>i</sub> × GSF<sub>b</sub>; this is the product of the gamma shielding provided by the soil cover under the building (GSF<sub>b</sub>) and the subfloor of the building (GSF<sub>i</sub>). This accounts for all the gamma shielding during the exposure time of a resident while indoors.
- A default gamma shielding factor for exposure to ionizing radiation in air (GSF<sub>a</sub>) is established at 1 (0% shielding).
- For the calculation of gamma shielding factors, a standard soil density of 1.6 g/cm<sup>3</sup> has been used.



4.10.7 Using the Back-calculated to Soil and Water Interactive Graph

When multiple media are contributing to the overall site dose, it may be more practical to remediate one medium versus another. In the case of the agriculture scenario, water and soil may both contribute to the dose from ingesting produce, milk, beef, swine, poultry, eggs, and fish. The interactive graph shows the DCCs for soil and water to achieve the target dose or target hazard index for the exposure route of concern.

The back-calculated to soil and water DCC results are listed last on the results page. The interactive graph is available for ingestion of produce, ingestion of milk, and ingestion of beef DCCs and may be accessed by clicking on any highlighted blue DCC value (see the image below).

Default									
Farmer DCCs for Contaminated Soil and Water									
		Fish Soil	Fish Water		Beef Soil	Beef Water		Dairy Soil	Dairy Water
		Intercept	Intercept		Intercept	Intercept		Intercept	Intercept
		PRG	PRG		PRG	PRG		PRG	PRG
Isotope	Fish Slope	(pCi/g)	(pCi/L)	Beef Slope	(pCi/g)	(pCi/L)	Dairy Slope	(pCi/g)	(pCi/L)
Am-241	-4.00E-03	7.56E-05	1.89E-02	-1.54E+01	1.26E+01	8.17E+02	-1.68E+01	2.01E+03	1.19E+05

Clicking on a highlighted blue back-calculated to soil and water DCC will take the user to the Farmer DCC Graphical Results page,

where the interactive graph is displayed.

The x-intercept (coordinate x,0) shows where the water DCC = TR and soil concentration must equal 0. The y-intercept (coordinate 0,y) shows where the soil DCC = TR and the water concentration must equal 0. Any point between (x,0) and (0,y) shows a separate DCC for water and soil that will meet the TR. Hovering the mouse over the graph will display moving lines that follow the mouse based on the x-coordinate (water DCC). Click anywhere on the graph to stop the lines from moving and to display the soil and water DCCs associated with that specific x-coordinate.



The scale of the graph has been programmed with upper bounds of 1.0E+6 pCi/g and 1.0E+6 pCi/L and lower bounds of 1.0E-10 pCi/g and 1.0E-10 pCi/L solely for presentation purposes to avoid scaling issues.

#### 4.10.8 Half Life Decay Function

In situations where the contaminant in the soil or air is not being replenished, the equations with decay should be used. An air situation that would require decay includes an accidental one-time air release from a factory. A soil situation that would require decay is a one-time dump at a landfill. In situations where the contaminant in the soil or air has a continual source (e.g., indoor radon from radium in the soil, an operating factory or landfill cap, or uranium mining), the equations without decay should be used. The decay function, found in section 4.10.8, can be multiplied by the ambient air equations above to acquire ambient air DCCs with decay.

- The following half life should be multiplied by CDI equations:

$$CDI_{decay} = \left( \frac{(1 - e^{-\lambda t})}{t(\text{year}) \times \lambda \left( \frac{1}{\text{year}} \right)} \right)$$

- The following half life should be multiplied by DCC equations:

$$PRG_{decay} = \left( \frac{t(\text{year}) \times \lambda \left( \frac{1}{\text{year}} \right)}{(1 - e^{-\lambda t})} \right)$$

## 5. Recommended Default Exposure Parameters

Table 1 presents definitions of equation variables and their default values. The DCC default values and exposure models are consistent with the Preliminary Remediation Goals at Superfund Sites ([PRG](#)) calculator. Both the DCC and PRG calculator default values are consistent with the Regional Screening Levels for Chemical Contaminants at Superfund Sites ([RSL](#)) calculator when the same pathways are addressed (e.g., ingestion and inhalation) and are analogous when pathways are similar (e.g., dermal and external exposure). This calculator, the PRG, and the RSL all follow the recommendations in the [OSWER Directive](#) concerning use of exposure parameters from the [2011 Exposure Factors Handbook](#). Any alternative values or assumptions used in remedy evaluation or selection on a CERCLA site should be presented with supporting rationale in Administrative Records.



Table 1. Recommended Default Exposure Parameters





Dose Coefficients			
Symbol ▼	Definition (units) ↗	Default ↗	Reference ↗
DCF <sub>ext-15cm</sub>	Dose Conversion Factor - external exposure (mrem/year per pCi/g)	Isotope-specific	<a href="#">ORNL 2014c</a>
DCF <sub>ext-1cm</sub>	Dose Conversion Factor - external exposure (mrem/year per pCi/g)	Isotope-specific	<a href="#">ORNL 2014c</a>
DCF <sub>ext-5cm</sub>	Dose Conversion Factor - external exposure (mrem/year per pCi/g)	Isotope-specific	<a href="#">ORNL 2014c</a>
DCF <sub>ext-gp</sub>	Dose Conversion Factor - external exposure (mrem/year per pCi/cm²)	Isotope-specific	<a href="#">ORNL 2014c</a>
DCF <sub>ext-sv</sub>	Dose Conversion Factor - external exposure (mrem/year per pCi/g)	Isotope-specific	<a href="#">ORNL 2014c</a>
DCF <sub>i</sub>	Dose Conversion Factor - inhalation (mrem/pCi)	Isotope-specific	<a href="#">ORNL 2014c</a>
DCF <sub>o</sub>	Oral Dose Conversion Factor - population (mrem/pCi)	Isotope-specific	<a href="#">ORNL 2014c</a>
DCF <sub>oa</sub>	Oral Dose Conversion Factor - adult only (mrem/pCi)	Isotope-specific	<a href="#">ORNL 2014c</a>
DCF <sub>sub</sub>	Dose Conversion Factor - submersion (mrem/yr per pCi/cm³): For use in this tool, ORNL 2014c units were converted to (mrem/yr per pCi/m³)	Isotope-specific	<a href="#">ORNL 2014c</a>

Dose and Decay Constant Variables			
Symbol ▼	Definition (units) ↗	Default ↗	Reference ↗
DL	Dose Limit (mrem/year)	1	
t <sub>cw</sub>	Time - construction worker (year)	1	U.S. EPA 2002b Exhibit 5-1
t <sub>far</sub>	Time - farmer (year)	1	U.S. EPA 2005 (pg. C-24/C-26)
t <sub>w</sub>	Time - indoor worker (year)	1	U.S. EPA 1991a (pg. 15)
t <sub>ow</sub>	Time - outdoor worker (year)	1	U.S. EPA 1991a (pg. 15)
t <sub>rec</sub>	Time - recreator (year)	site-specific	site-specific
t <sub>res</sub>	Time - resident (year)	1	U.S. EPA 2011a, Table 16-108; 90th percentile or current residence time.
t <sub>w</sub>	Time - worker (year)	1	U.S. EPA 1991a (pg. 15)
λ	decay constant = 0.693/half-life (year <sup>1</sup> ) where 0.693 = ln(2)	Isotope-specific	Developed for Radionuclide Soil Screening calculator

Miscellaneous Variables			
Symbol ▼	Definition (units) ↗	Default ↗	Reference ↗
AAF <sub>far-a</sub>	Farmer Age Adjustment Factor - adult (unitless)	0.85	Developed for this calculator
AAF <sub>far-c</sub>	Farmer Age Adjustment Factor - child (unitless)	0.15	Developed for this calculator





AAF <sub>rec-a</sub>	Recreator Age Adjustment Factor - adult (unitless)	site-specific	Developed for this calculator
AAF <sub>rec-c</sub>	Recreator Age Adjustment Factor - child (unitless)	site-specific	Developed for this calculator
AAF <sub>res-a</sub>	Resident Age Adjustment Factor - adult (unitless)	0.77	Developed for this calculator
AAF <sub>res-c</sub>	Resident Age Adjustment Factor - child (unitless)	0.23	Developed for this calculator
ACF <sub>ext-15cm</sub>	Area Correction Factor - 15cm (unitless)	Isotope-specific	<a href="#">ORNL 2014a</a>
ACF <sub>ext-1cm</sub>	Area Correction Factor - 1cm (unitless)	Isotope-specific	<a href="#">ORNL 2014a</a>
ACF <sub>ext-5cm</sub>	Area Correction Factor - 5cm (unitless)	Isotope-specific	<a href="#">ORNL 2014a</a>
ACF <sub>ext-gp</sub>	Area Correction Factor - ground plane (unitless)	Isotope-specific	<a href="#">ORNL 2014a</a>
ACF <sub>ext-sv</sub>	Area Correction Factor - soil volume (unitless)	Isotope-specific	<a href="#">ORNL 2014a</a>
CF <sub>far-beef</sub>	Beef Contaminated Fraction - farmer (unitless)	1	Developed for Radionuclide Soil Screening calculator
CF <sub>far-dairy</sub>	Dairy Contaminated Fraction - farmer (unitless)	1	Developed for Radionuclide Soil Screening calculator
CF <sub>far-egg</sub>	Egg Contaminated Fraction - farmer (unitless)	1	Developed for Radionuclide Soil Screening calculator
CF <sub>far-fish</sub>	Fish Contaminated Fraction - farmer (unitless)	1	Developed for Radionuclide Soil Screening calculator
CF <sub>far-goat</sub>	Goat Contaminated Fraction - Farmer (unitless)	1	Developed for Radionuclide Soil Screening calculator
CF <sub>far-goat-milk</sub>	Goat Milk Contaminated Fraction - Farmer (unitless)	1	Developed for Radionuclide Soil Screening calculator
CF <sub>far-poultry</sub>	Poultry Contaminated Fraction - farmer (unitless)	1	Developed for Radionuclide Soil Screening calculator
CF <sub>far-produce</sub>	Produce Contaminated Fraction - farmer (unitless)	1	U.S. EPA 2011, U.S. EPA 2005
CF <sub>far-sheep</sub>	Sheep Contaminated Fraction - Farmer (unitless)	1	Developed for Radionuclide Soil Screening calculator
CF <sub>far-sheep-milk</sub>	Sheep Milk Contaminated Fraction - Farmer (unitless)	1	Developed for Radionuclide Soil Screening calculator
CF <sub>far-swine</sub>	Swine Contaminated Fraction - farmer (unitless)	1	Developed for Radionuclide Soil Screening calculator
CF <sub>res-produce</sub>	Produce Contaminated Fraction - resident (unitless)	1	U.S. EPA 2011, U.S. EPA 2005
GSF <sub>a</sub>	Gamma Shielding Factor - Air (unitless)	1	Developed for Radionuclide Soil Screening calculator
GSF <sub>ext-15cm</sub>	Gamma Shielding Factor - 15cm (unitless)	Isotope-specific	<a href="#">ORNL 2014b</a>
GSF <sub>ext-1cm</sub>	Gamma Shielding Factor - 1cm (unitless)	Isotope-specific	<a href="#">ORNL 2014b</a>
GSF <sub>ext-5cm</sub>	Gamma Shielding Factor - 5cm (unitless)	Isotope-specific	<a href="#">ORNL 2014b</a>
GSF <sub>ext-gp</sub>	Gamma Shielding Factor - ground plane (unitless)	Isotope-specific	<a href="#">ORNL 2014b</a>
GSF <sub>ext-sv</sub>	Gamma Shielding Factor - soil volume (unitless)	Isotope-specific	<a href="#">ORNL 2014b</a>
			U.S. EPA 2000a. (pg. 2-22). U.S. EPA 2000b. (pg. 2-

GSF <sub>i</sub>	Gamma Shielding Factor - Indoor (unitless)	0.4	18)
K	Andelman Volatilization Factor (L/m <sup>3</sup> )	0.5	U.S. EPA 1991b (pg. 20)
ρ <sub>m</sub>	Density of milk (kg/L)	1.03	<a href="#">Milk Composition &amp; Synthesis Resource Library</a>

Tissue Transfer Factors and Animal Ingestion Rates of Fodder, Water, and Soil			
Symbol 	Definition (units) 	Default 	Reference 
BCF	Fish Transfer Factor (L/kg)	radionuclide-specific	hierarchy selection in Section 2.4.2
B <sub>v dry</sub>	Soil to Plant Transfer Factor - dry (pCi/g-dry plant per pCi/g-dry soil)	radionuclide-specific	hierarchy selection in Section 2.4.2
B <sub>v wet</sub>	Soil to Plant Transfer Factor - wet (pCi/g-fresh plant per pCi/g-dry soil)	radionuclide-specific	hierarchy selection in Section 2.4.2 climate and soil selection in 2.5.1.2
F	Irrigation Period (unitless)	0.25	Personal communication
f <sub>p-beef</sub>	Animal On-site Fodder Fraction - beef (unitless)	1	Developed for this calculator
f <sub>p-dairy</sub>	Animal On-site Fodder Fraction - dairy (unitless)	1	Developed for this calculator
f <sub>p-goat</sub>	Animal On-site Fraction - goat (unitless)	1	Developed for this calculator
f <sub>p-goat-milk</sub>	Animal On-site Fraction - goat milk (unitless)	1	Developed for this calculator
f <sub>p-poultry</sub>	Animal On-site Fodder Fraction - poultry (unitless)	1	Developed for this calculator
f <sub>p-sheep</sub>	Animal On-site Fraction - sheep (unitless)	1	Developed for this calculator
f <sub>p-sheep-milk</sub>	Animal On-site Fraction - sheep milk (unitless)	1	Developed for this calculator
f <sub>p-swine</sub>	Animal On-site Fodder Fraction - swine (unitless)	1	Developed for this calculator
f <sub>s-beef</sub>	Animal On-Site Soil Fraction - beef (unitless)	1	Developed for this calculator
f <sub>s-dairy</sub>	Animal On-Site Soil Fraction - dairy (unitless)	1	Developed for this calculator
f <sub>s-poultry</sub>	Fraction of Year Animal On Site - poultry (unitless)	1	Developed for this calculator
f <sub>s-swine</sub>	Animal On-Site Soil Fraction - swine (unitless)	1	Developed for this calculator
I <sub>f</sub>	Interception Fraction (unitless)	0.42	Miller, C. W. 1980
I <sub>r</sub>	Irrigation Rate (L/m <sup>2</sup> - day)	3.62	Personal communication
Irr <sub>dep</sub>	aerial deposition from irrigation multiplier	isotope-specific	Calculated
Irr <sub>res</sub>	resuspension from irrigation multiplier	isotope-specific	Calculated
Irr <sub>rup</sub>	root uptake from irrigation multiplier	isotope-specific	Calculated
MLF <sub>pasture</sub>	Pasture Plant Mass Loading Factor (g dry soil per g dry plant)	0.25	Hinton, T. G. 1992
MLF <sub>produce</sub>	Produce Plant Mass Loading Factor (g dry soil per g fresh plant)	plant-specific	See section 2.5.3 of this guide for details
P	Area Density for Root Zone (kg/m <sup>2</sup> )	240	Hoffman, F. O., R. H. Gardner, and K. F. Eckerman.





			1982; Peterson, H. T., Jr. 1983; McKone, T. E. 1994
Q <sub>p-beef</sub>	Beef Fodder Intake Rate (kg/day)	11.77	U.S. EPA 2005 (pg. B-138)
Q <sub>p-dairy</sub>	Dairy Fodder Intake Rate (kg/day)	20.3	U.S. EPA 2005 (pg. B-145)
Q <sub>p-goat</sub>	Goat Fodder Intake Rate (kg/day)	1.27	<a href="#">Lyons et. al. 1999</a>
Q <sub>p-goat-milk</sub>	Goat Milk Fodder Intake Rate (kg/day)	1.59	<a href="#">Lyons et. al. 1999</a> and <a href="#">Tarr</a>
Q <sub>p-poultry</sub>	Chicken Fodder Intake Rate (kg/day)	0.2	U.S. EPA 2005 (pg. B-158/164)
	Duck Fodder Intake Rate (kg/day)	0.24	NRC 1994
	Turkey Fodder Intake Rate (kg/day)	0.68	NRC 1994
	Goose Fodder Intake Rate (kg/day)	0.33	NRC 1994
Q <sub>p-sheep</sub>	Sheep Fodder Intake Rate (kg/day)	1.75	<a href="#">Lyons et. al. 1999</a> and <a href="#">OMAFRA</a>
Q <sub>p-sheep-milk</sub>	Sheep Milk Fodder Intake Rate (kg/day)	3.15	<a href="#">Lyons et. al. 1999</a> and <a href="#">OMAFRA</a>
Q <sub>p-swine</sub>	Swine Fodder Intake Rate (kg/day)	4.7	U.S. EPA 2005 (pg. B-152)
Q <sub>s-beef</sub>	Beef Soil Intake Rate (kg/day)	0.5	U.S. EPA 2005 (pg. B-139)
Q <sub>s-dairy</sub>	Dairy Soil Intake Rate (kg/day)	0.4	U.S. EPA 2005 (pg. B-146)
Q <sub>s-goat</sub>	Goat Soil Intake Rate (kg/day)	0.23	<a href="#">Handbook of Ecotoxicology</a> (Q <sub>s</sub> is 18% of Q <sub>p</sub> ; sheep surrogate used)
Q <sub>s-goat-milk</sub>	Goat Milk Soil Intake Rate (kg/day)	0.29	<a href="#">Handbook of Ecotoxicology</a> (Q <sub>s</sub> is 18% of Q <sub>p</sub> ; sheep surrogate used)
Q <sub>s-poultry</sub>	Chicken Soil Intake Rate (kg/day)	0.022	U.S. EPA 2005 (pg. B-159/165)
	Duck Soil Intake Rate (kg/day)	0.024	NRC 1994 (Q <sub>s</sub> is 10% of Q <sub>p</sub> )
	Turkey Soil Intake Rate (kg/day)	0.068	NRC 1994 (Q <sub>s</sub> is 10% of Q <sub>p</sub> )
	Goose Soil Intake Rate (kg/day)	0.033	NRC 1994 (Q <sub>s</sub> is 10% of Q <sub>p</sub> )
Q <sub>s-sheep</sub>	Sheep Soil Intake Rate (kg/day)	0.32	<a href="#">Handbook of Ecotoxicology</a> (Q <sub>s</sub> is 18% of Q <sub>p</sub> )
Q <sub>s-sheep-milk</sub>	Sheep Milk Soil Intake Rate (kg/day)	0.57	<a href="#">Handbook of Ecotoxicology</a> (Q <sub>s</sub> is 18% of Q <sub>p</sub> )
Q <sub>s-swine</sub>	Swine Soil Intake Rate (kg/day)	0.37	U.S. EPA 2005 (pg. B-153)
Q <sub>w-beef</sub>	Beef Water Intake Rate (L/day)	53	U.S. EPA 1999a (pg 10-23). U.S. EPA 1997b.
Q <sub>w-dairy</sub>	Dairy Water Intake Rate (L/day)	92	U.S. EPA 1999a (pg 10-23). U.S. EPA 1997b.
Q <sub>w-goat</sub>	Goat Water Intake Rate (L/day)	3.81	<a href="#">Tarr</a>
Q <sub>w-goat-milk</sub>	Goat Milk Water Intake Rate (L/day)	8.75	<a href="#">Tarr</a>
Q <sub>w-poultry</sub>	Chicken Water Intake Rate (L/day)	0.4	U.S. EPA 2005 (pg. B-159/165), NRC 1994 pg.15 (Q <sub>w</sub> =2×Q <sub>p</sub> )
	Duck Water Intake Rate (L/day)	0.48	NRC 1994 pg. 15 (Q <sub>w</sub> = 2 × Q <sub>p</sub> )
	Turkey Water Intake Rate (L/day)	1.36	NRC 1994 pg. 15 (Q <sub>w</sub> = 2 × Q <sub>p</sub> )
	Goose Water Intake Rate (L/day)	0.66	NRC 1994 pg. 15 (Q <sub>w</sub> = 2 × Q <sub>p</sub> )
Q <sub>w-sheep</sub>	Sheep Water Intake Rate (L/day)	5.25	<a href="#">OMAFRA</a>
Q <sub>w-sheep-milk</sub>	Sheep Milk Water Intake Rate (L/day)	10.4	<a href="#">OMAFRA</a>
Q <sub>w-swine</sub>	Swine Water Intake Rate (L/day)	11.4	NEC, Swine Nutrition Guide (pg. 19). U.S. EPA 1998 (pg B-180)
R <sub>es</sub>	soil resuspension multiplier (g dry soil per g fresh plant)	=MLF (pasture or produce)	Hinton 1992
R <sub>upp</sub>	dry root uptake for pasture multiplier (pCi/g-dry plant per pCi/g-dry soil)	radionuclide-specific (=Bv <sub>dry</sub> )	hierarchy selection in Section 2.4.2

$R_{upv}$	wet root uptake for produce multiplier (pCi/g-fresh plant per pCi/g-dry soil)	radionuclide-specific (=Bv <sub>wet</sub> )	hierarchy selection in Section 2.4.2
T	Translocation Factor (unitless)	1	NCRP 1984
t <sub>b</sub>	Long Term Deposition and Buildup (day)	10950	NCRP 1985
TF <sub>beef</sub>	Beef Transfer Factor (day/kg)	radionuclide-specific	hierarchy selection in Section 2.4.2
TF <sub>dairy</sub>	Dairy Transfer Factor (day/L)	radionuclide-specific	hierarchy selection in Section 2.4.2
TF <sub>egg</sub>	Egg Transfer Factor (day/kg)	radionuclide-specific	hierarchy selection in Section 2.4.2
TF <sub>goat</sub>	Goat Transfer Factor (day/kg)	radionuclide-specific	hierarchy selection in Section 2.4.2
TF <sub>goat-milk</sub>	Goat Milk Transfer Factor (day/L)	radionuclide-specific	hierarchy selection in Section 2.4.2
TF <sub>poultry</sub>	Poultry Transfer Factor (day/kg)	radionuclide-specific	hierarchy selection in Section 2.4.2
TF <sub>sheep</sub>	Sheep Transfer Factor (day/kg)	radionuclide-specific	hierarchy selection in Section 2.4.2
TF <sub>sheep-milk</sub>	Sheep Milk Transfer Factor (day/L)	radionuclide-specific	hierarchy selection in Section 2.4.2
TF <sub>swine</sub>	Swine Transfer Factor (day/kg)	radionuclide-specific	hierarchy selection in Section 2.4.2
t <sub>v</sub>	Above Ground Exposure Time (day)	60	NCRP 1985
t <sub>w</sub>	Weathering Half-life (day)	14	NCRP 1989
Y <sub>v</sub>	Plant Yield - wet (kg/m <sup>2</sup> )	2	NCRP 1985
λ <sub>B</sub>	Effective Rate of Removal (1/day)	λ <sub>HL</sub> + λ <sub>i</sub>	NCRP 1989
λ <sub>E</sub>	Effective Rate for Removal from Produce (1/day)	λ <sub>i</sub> + (0.693/t <sub>w</sub> )	NCRP 1989
λ <sub>HL</sub>	Soil Leaching Rate (1/day)	0.000027	NCRP 1989
λ <sub>i</sub>	decay (1/day)	0.693/HL (days)	NCRP 1989

Inhalation, Ingestion, and Consumption Rates			
Symbol 	Definition (units) 	Default 	Reference 
DFA <sub>rec-adj</sub>	Recreator Surface Water Immersion Factor - age-adjusted (hour)	site-specific	U.S. EPA 1991a (pg. 15)
IFA <sub>far-adj</sub>	Farmer Inhalation Fraction - age-adjusted (m <sup>3</sup> /year)	6475	Calculated using the age-adjusted intake factors equation.
IFA <sub>rec-adj</sub>	Recreator Inhalation Fraction - age-adjusted (m <sup>3</sup> /year)	site-specific	Calculated using the age-adjusted intake factors equation.
IFA <sub>res-adj</sub>	Resident Inhalation Fraction - age-adjusted (m <sup>3</sup> /year)	6195	Calculated using the age-adjusted intake factors equation.
IFB <sub>far-adj</sub>	Farmer Beef Ingestion Fraction - age-adjusted (g/year)	52,474	Calculated using the aged adjusted intake factors equation
IFD <sub>far-adj</sub>	Farmer Dairy Ingestion Fraction - age-adjusted (g/year)	253,451	Calculated using the aged adjusted intake factors equation

IFE <sub>far-adj</sub>	Farmer Egg Ingestion Fraction - age-adjusted (g/year)	19,395	Calculated using the aged adjusted intake factors equation
IF <sub>far-adj</sub>	Farmer Produce Ingestion Fraction - age-adjusted (g)	plant-specific	Calculated using the aged adjusted intake factors equation
IFFI <sub>far-adj</sub>	Farmer Fish Ingestion Fraction - age-adjusted (g/year)	250,474	Calculated using the aged adjusted intake factors equation
IFP <sub>far-adj</sub>	Farmer Poultry Ingestion Fraction - age-adjusted (g/year)	34,414	Calculated using the aged adjusted intake factors equation
IF <sub>res-adj</sub>	Resident Produce Ingestion Fraction - age-adjusted (g)	plant-specific	Calculated using the aged adjusted intake factors equation
IFS <sub>rec-adj</sub>	Recreator Soil Ingestion Fraction - age-adjusted (mg/year)	site-specific	Calculated using the age-adjusted intake factors equation.
IFS <sub>res-adj</sub>	Resident Soil Ingestion Fraction - age-adjusted (mg/year)	43,050	Calculated using the age-adjusted intake factors equation.
IFSW <sub>far-adj</sub>	Farmer Swine Ingestion Fraction - age-adjusted (g/year)	29,288	Calculated using the aged adjusted intake factors equation
IFW <sub>rec-adj</sub>	Recreator Surface Water Ingestion Fraction - age-adjusted (L/year)	site-specific	Calculated using the age-adjusted intake factors equation.
IFW <sub>res-adj</sub>	Resident Tapwater Ingestion Fraction - age-adjusted (L/year)	737	Calculated using the age-adjusted intake factors equation.
IRA <sub>cw</sub>	Construction Worker Soil Inhalation Rate (m <sup>3</sup> /day; based on a rate of 2.5m <sup>3</sup> /hr for 24hr)	60	U.S. EPA 1997a (pg. 5-11)
IRA <sub>far-a</sub>	Farmer Inhalation Rate - adult (m <sup>3</sup> /day)	20	U.S. EPA 1991a (pg. 15)
IRA <sub>far-c</sub>	Farmer Inhalation Rate - child (m <sup>3</sup> /day)	10	U.S. EPA 1997a (pg. 5-11)
IRA <sub>rw</sub>	Indoor Worker Soil Inhalation Rate (m <sup>3</sup> /day; based on a rate of 2.5m <sup>3</sup> /hr for 24hr)	60	U.S. EPA 1997a (pg. 5-11)
IRA <sub>ow</sub>	Outdoor Worker Soil Inhalation Rate (m <sup>3</sup> /day; based on a rate of 2.5m <sup>3</sup> /hr for 24hr)	60	U.S. EPA 1997a (pg. 5-11)
IRA <sub>rec-a</sub>	Recreator Soil Inhalation Rate - adult (m <sup>3</sup> /day)	20	U.S. EPA 1991a (pg. 15)
IRA <sub>rec-c</sub>	Recreator Inhalation Rate - child (m <sup>3</sup> /day)	10	U.S. EPA 1997a (pg. 5-11)
IRA <sub>res-a</sub>	Resident Soil Inhalation Rate - adult (m <sup>3</sup> /day)	20	U.S. EPA 1991a (pg. 15)
IRA <sub>res-c</sub>	Resident Soil Inhalation Rate - child (m <sup>3</sup> /day)	10	U.S. EPA 1997a (pg. 5-11)
IRA <sub>w</sub>	Composite Worker Soil Inhalation Rate (m <sup>3</sup> /day; based on a rate of 2.5m <sup>3</sup> /hr for 24hr)	60	U.S. EPA 1997a (pg. 5-11)
IRB <sub>far-a</sub>	Farmer Beef Ingestion Rate - adult (g/day)	165.3	U.S. EPA 2011 (Table 13-33)
IRB <sub>far-c</sub>	Farmer Beef Ingestion Rate - child (g/day)	62.8	U.S. EPA 2011 (Table 13-33)
IRD <sub>far-a</sub>	Farmer Dairy Ingestion Rate - adult (g/day)	676.4	U.S. EPA 2011 (Table 11-4)
IRD <sub>far-c</sub>	Farmer Dairy Ingestion Rate - child (g/day)	994.7	U.S. EPA 2011 (Table 11-4)
IRE <sub>far-a</sub>	Farmer Egg Ingestion Rate - adult (g/day)	59.6	U.S. EPA 2011 (Table 13-40)
IRE <sub>far-c</sub>	Farmer Egg Ingestion Rate - child (g/day)	31.7	U.S. EPA 2011 (Table 13-40)
IR <sub>far-a</sub>	Farmer Produce Ingestion Rate - adult (g/day)	plant-specific	U.S. EPA 2011 (Table 13-10)
IR <sub>far-c</sub>	Farmer Produce Ingestion Rate - child (g/day)	plant-specific	U.S. EPA 2011 (Table 13-10)

$IRF_{far-a}$	Farmer Fish Ingestion Rate - adult (g/day)	831.8	U.S. EPA 2011 (Table 13-20)
$IRF_{far-c}$	Farmer Fish Ingestion Rate - child (g/day)	57.4	U.S. EPA 2011 (Table 13-20)
$IRF_{res}$	Resident Fish Ingestion Rate (g/day)	54	U.S. EPA 1991a (page 15)
$IRP_{far-a}$	Farmer Poultry Ingestion Rate - adult (g/day)	107.4	U.S. EPA 2011 (Table 13-52)
$IRP_{far-c}$	Farmer Poultry Ingestion Rate - child (g/day)	46.9	U.S. EPA 2011 (Table 13-52)
$IR_{res-a}$	Resident Produce Ingestion Rate - adult (g/day)	plant-specific	U.S. EPA 2011 (Table 13-10)
$IR_{res-c}$	Resident Produce Ingestion Rate - child (g/day)	plant-specific	U.S. EPA 2011 (Table 13-10)
$IRS_{cw}$	Construction Worker Soil Ingestion Rate (mg/day)	330	U.S. EPA 1991a (pg. 15)
$IRS_{far-a}$	Farmer Soil Ingestion Rate - adult (mg/day)	100	U.S. EPA 1991a (pg. 15)
$IRS_{far-adj}$	Farmer Soil Ingestion Rate - age-adjusted (mg/day)	115	Calculated using the age-adjusted intake factors equation.
$IRS_{far-c}$	Farmer Soil Ingestion Rate - child (mg/day)	200	U.S. EPA 1991a (pg. 15)
$IRS_{iw}$	Indoor Worker Soil Ingestion Rate (mg/day)	50	U.S. EPA 2001 (pg. 4-3)
$IRS_{ow}$	Outdoor Worker Soil Ingestion Rate (mg/day)	100	U.S. EPA 1991a (pg. 15)
$IRS_{rec-a}$	Recreator Soil Ingestion Rate - adult (mg/day)	100	U.S. EPA 1991a (pg. 15)
$IRS_{rec-c}$	Recreator Soil Ingestion Rate - child (mg/day)	200	U.S. EPA 1991a (pg. 15)
$IRS_{res-a}$	Resident Soil Ingestion Rate - adult (mg/day)	100	U.S. EPA 1991a (pg. 15)
$IRS_{res-c}$	Resident Soil Ingestion Rate - child (mg/day)	200	U.S. EPA 1991a (pg. 15)
$IRS_w$	Composite Worker Soil Ingestion Rate (mg/day)	100	U.S. EPA 1991a (pg. 15)
$IRSW_{far-a}$	Farmer Swine Ingestion Rate - adult (g/day)	92.5	U.S. EPA 2011 (Table 13-51)
$IRSW_{far-c}$	Farmer Swine Ingestion Rate - child (g/day)	33.7	U.S. EPA 2011 (Table 13-51)
$IRW_{rec-a}$	Recreator Surface Water Ingestion Rate - adult (L/hour)	0.071	Adult upper percentile from Table 3.5 of EFH 2011
$IRW_{rec-c}$	Recreator Surface Water Ingestion Rate - child (L/hour)	0.12	Child upper percentile from Table 3.5 of EFH 2011
$IRW_{res-a}$	Resident Tapwater Ingestion Rate - adult (L/day)	2.5	U.S. EPA 2011a, Tables 3-15 and 3-33; weighted average of 90th percentile consumer-only ingestion of drinking water (birth to <6 years)
$IRW_{res-c}$	Resident Tapwater Ingestion Rate - child (L/day)	0.78	U.S. EPA 2011a, Tables 3-15 and 3-33; weighted average of 90th percentile consumer-only ingestion of drinking water (birth to <6 years)

Exposure Frequency, Exposure Duration, and Exposure Time Variables			
Symbol 	Definition (units) 	Default 	Reference 
$DW_{cw}$	Construction Worker Exposure Frequency (days/week)	5	U.S. EPA 2002b Exhibit 5-1
ED	Farmer Exposure Duration (years)	40	U.S. EPA 2005 (Table 6-3)



far			
ED <sub>far-a</sub>	Farmer Exposure Duration - adult (years)	34	U.S. EPA 1994a
ED <sub>far-c</sub>	Farmer Exposure Duration - child (years)	6	U.S. EPA 2005 (Table 6-3)
ED <sub>rec</sub>	Recreator Exposure Duration (years)	site-specific	U.S. EPA 2011a, Table 16-108; 90th percentile or current residence time.
ED <sub>rec-a</sub>	Recreator Exposure Duration - adult (years)	site-specific	-
ED <sub>rec-c</sub>	Recreator Exposure Duration - child (years)	site-specific	U.S. EPA 1991a, Pages 6 and 15
ED <sub>res</sub>	Resident Exposure Duration (years)	26	U.S. EPA 2011a, Table 16-108; 90th percentile or current residence time.
ED <sub>res-a</sub>	Resident Exposure Duration - adult (years)	20	ED <sub>ress</sub> (26 years) - ED <sub>ressc</sub> (6 years)
ED <sub>res-c</sub>	Resident Exposure Duration - child (years)	6	U.S. EPA 1991a, Pages 6 and 15
EF <sub>cw</sub>	Construction Worker Exposure Frequency (days/year)	250	U.S. EPA 2002b Exhibit 5-1
EF <sub>far</sub>	Farmer Exposure Frequency (days/year)	350	U.S. EPA 1991a (pg. 15)
EF <sub>far-a</sub>	Farmer Exposure Frequency - adult (days/year)	350	U.S. EPA 1991a (pg. 15)
EF <sub>far-c</sub>	Farmer Exposure Frequency - child (days/year)	350	U.S. EPA 1991a (pg. 15)
EF <sub>iw</sub>	Indoor Worker Exposure Frequency (days/year)	250	U.S. EPA 1991a (pg. 15)
EF <sub>ow</sub>	Outdoor Worker Exposure Frequency (days/year)	225	U.S. EPA 1991a (pg. 15)
EF <sub>rec</sub>	Recreator Exposure Frequency (days/year)	site-specific	U.S. EPA 1991a (pg. 15)
EF <sub>rec-a</sub>	Recreator Exposure Frequency - adult (days/year)	site-specific	U.S. EPA 1991a (pg. 15)
EF <sub>rec-c</sub>	Recreator Exposure Frequency- child (days/year)	site-specific	U.S. EPA 1991a (pg. 15)
EF <sub>res</sub>	Resident Exposure Frequency (days/year)	350	U.S. EPA 1991a (pg. 15)
EF <sub>res-a</sub>	Resident Exposure Frequency - adult (days/year)	350	U.S. EPA 1991a (pg. 15)
EF <sub>res-c</sub>	Resident Exposure Frequency - child (days/year)	350	U.S. EPA 1991a (pg. 15)
EF <sub>w</sub>	Composite Worker Exposure Frequency (days/year)	250	U.S. EPA 1991a (pg. 15)
ET <sub>cw</sub>	Construction Worker Exposure Time (hours/day)	8	Eight Hours per 24 hour Day
ET <sub>event-far-a</sub>	Farmer Water Exposure Time - adult (hours/event)	0.71	U.S. EPA 1997a
ET <sub>event-far-c</sub>	Farmer Water Exposure Time - child (hours/event)	0.54	U.S. EPA 1997a
ET <sub>event-res-c</sub>	Resident Tapwater Exposure Time - adult (hours/event)	0.71	U.S. EPA 1997a
ET <sub>event-res-c</sub>	Resident Tapwater Exposure Time - child (hours/event)	0.54	U.S. EPA 1997a

$ET_{far-a}$	Farmer Exposure Time - adult (hours/day)	24	24 Hours per 24 hour Day
$ET_{far-away}$	Farmer Soil Exposure Time - away (hours/day)	1.83	U.S. EPA 2011 (Tables 16-20 and 16-24 total of time in vehicles, near vehicles and outdoors other than near residence 25 <sup>th</sup> %)
$ET_{far-c}$	Farmer Exposure Time - child (hours/day)	24	24 Hours per 24 hour Day
$ET_{far-i}$	Farmer Indoor Soil Exposure Time (hours/day)	10.008	24 hours/day - ( $ET_{fo}$ + $ET_{fa}$ )
$ET_{far-o}$	Farmer Outdoor Soil Exposure Time (hours/day)	12.167	U.S. EPA 2011 (Table 16-20 95 <sup>th</sup> %)
$ET_{IW}$	Indoor Worker Exposure Time (hours/day)	8	Eight Hours per 24 hour Day
$ET_{OW}$	Outdoor Worker Exposure Time (hours/day)	8	Eight Hours per 24 hour Day
$ET_{rec}$	Recreator Exposure Time (hours/day)	site-specific	site-specific
$ET_{rec-a}$	Recreator Exposure Time - adult (hours/day)	site-specific	site-specific
$ET_{rec-c}$	Recreator Exposure Time - child (hours/day)	site-specific	site-specific
$ET_{res-a}$	Resident Exposure Time - adult (hours/day)	24	24 Hours per 24 hour Day
$ET_{res-c}$	Resident Exposure Time - child (hours/day)	24	24 Hours per 24 hour Day
$ET_{res-i}$	Resident Indoor Soil Exposure Time (hours/day)	16.416	U.S. EPA 2011 (Table 16-16 50 <sup>th</sup> %)
$ET_{res-o}$	Resident Outdoor Soil Exposure Time (hours/day)	1.75	U.S. EPA 2011 (Table 16-20 50 <sup>th</sup> %)
$ET_w$	Composite Worker Exposure Time (hours/day)	8	Eight Hours per 24 hour Day
$EV_{far-a}$	Number of farmer tapwater bathing events per day - adult (events/day)	1	U.S. EPA 2004 Exhibit A-9
$EV_{far-c}$	Number of farmer tapwater bathing events per day - child (events/day)	1	U.S. EPA 2004 Exhibit A-9
$EV_{rec-a}$	Number of recreator surface water bathing events per day - adult (events/day)	site-specific	site-specific
$EV_{rec-c}$	Number of recreator surface water bathing events per day - child (events/day)	site-specific	site-specific
$EV_{res-a}$	Number of resident tapwater bathing events per day - adult (events/day)	1	U.S. EPA 2004 Exhibit A-9
$EV_{res-c}$	Number of resident tapwater bathing events per day - child (events/day)	1	U.S. EPA 2004 Exhibit A-9
$EW_{ow}$	Construction Worker Exposure Frequency (weeks/year)	50	U.S. EPA 2002b Exhibit 5-1

Soil to Groundwater SSL Factor Variables			
Symbol ▼	Definition (units) ▼	Default ▼	Reference ▼
$C_w$	Target soil leachate concentration (pCi/L)	nonzero MCL or RSL × DAF	U.S. EPA. 2002b Equation 4-14
d	mixing zone depth (m)	site-specific	U.S. EPA. 2002b Equation 4-12
$d_a$	aquifer thickness (m)	site-specific	U.S. EPA. 2002b Equation 4-10
DAF	Dilution attenuation factor (unitless)	1 (or site-specific)	U.S. EPA. 2002b Equation 4-11

d <sub>s</sub>	depth of source (m)	site-specific	U.S. EPA. 2002b Equation 4-10
ED <sub>gw</sub>	Exposure duration	70	U.S. EPA. 2002b Equation 4-14
I	Infiltration Rate (m/year)	0.18	U.S. EPA. 2002b Equation 4-11
i	hydraulic gradient (m/m)	site-specific	U.S. EPA. 2002b Equation 4-11
K	aquifer hydraulic conductivity (m/year)	site-specific	U.S. EPA. 2002b Equation 4-11
K <sub>d</sub>	soil-water partition coefficient (L/kg)	= K <sub>oc</sub> *f <sub>oc</sub> for organics	U.S. EPA. 2002b Equation 4-10
L	source length parallel to ground water flow (m)	site-specific	U.S. EPA. 2002b Equation 4-11
n	total soil porosity(L <sub>pore</sub> /L <sub>soil</sub> )	= 1-(ρ <sub>b</sub> /ρ <sub>s</sub> )	U.S. EPA. 2002b Equation 4-10
θ <sub>a</sub>	air-filled soil porosity (L <sub>air</sub> /L <sub>soil</sub> )	= n-θ <sub>w</sub>	U.S. EPA. 2002b Equation 4-10
θ <sub>w</sub>	water-filled soil porosity (L <sub>water</sub> /L <sub>soil</sub> )	0.3	U.S. EPA. 2002b Equation 4-10
ρ <sub>b</sub>	dry soil bulk density (kg/L)	1.5	U.S. EPA. 2002b Equation 4-10
ρ <sub>s</sub>	soil particle density (Kg/L)	2.65	U.S. EPA. 2002b Equation 4-10

Wind Particulate Emission Factor Variables			
Symbol ▾	Definition (units) ∞	Default ∞	Reference ∞
A	Dispersion constant (unitless)	PEF and region-specific	U.S. EPA 2002b Exhibit D-2
A <sub>s</sub>	Areal extent of the site or contamination (acres)	0.5 (range 0.5 to 500 )	U.S. EPA 2002b Exhibit D-2
B	Dispersion constant (unitless)	PEF and region-specific	U.S. EPA 2002b Exhibit D-2
C	Dispersion constant (unitless)	PEF and region-specific	U.S. EPA 2002b Exhibit D-2
F(x)	Function Dependent on 0.886 × (U <sub>t</sub> /U <sub>m</sub> ) (unitless)	0.194	U.S. EPA. 1996, Appendix D Table 2
Q/C <sub>wind</sub>	Inverse of the Mean Concentration at the Center of a 0.5-Acre-Square Source (g/m <sup>2</sup> -s per kg/m <sup>3</sup> )	93.77 (region-specific)	U.S. EPA 2002b Exhibit D-2
U <sub>m</sub>	Mean Annual Wind Speed (m/s)	4.69	U.S. EPA. 1996, Appendix D Table 2
U <sub>t</sub>	Equivalent Threshold Value of Wind Speed at 7m (m/s)	11.32	U.S. EPA. 1996, Appendix D Table 2
V	Fraction of Vegetative Cover (unitless)	0.5	U.S. EPA. 2002b Equation 4-5
PEF <sub>w</sub>	Particulate Emission Factor - Minneapolis (m <sup>3</sup> /kg)	1.36 x 10 <sup>9</sup> (region-specific)	U.S. EPA 2002b Exhibit D-2

Mechanical Particulate Emission Factor Variables from Standard Unpaved Road Vehicle Traffic			
Symbol ▾	Definition (units) ∞	Default ∞	Reference ∞
A	Dispersion constant (unitless)	12.9351	U.S. EPA 2002b Equation 5-6
A <sub>R</sub>	Surface area of contaminated road segment (m <sup>2</sup> )	(A <sub>R</sub> = L <sub>R</sub> x W <sub>R</sub> x 0.092903m <sup>2</sup> /ft <sup>2</sup> )	U.S. EPA 2002b Equation 5-5
A <sub>s</sub>	Areal extent of site surface soil contamination (acres)	0.5 (range 0.5 to 500 )	U.S. EPA 2002b Equation 5-6

B	Dispersion constant (unitless)	5.7383	U.S. EPA 2002b Equation 5-6
C	Dispersion constant (unitless)	71.7711	U.S. EPA 2002b Equation 5-6
F <sub>D</sub>	Dispersion correction factor (unitless)	0.185	U.S. EPA 2002b Equation 5-5
L <sub>R</sub>	Length of road segment (ft)	Site-specific	U.S. EPA 2002b Equation 5-5
P	Number of days with at least 0.01 inches of precipitation (days/year)	Site-specific	U.S. EPA 2002b Exhibit 5-2
Q/C <sub>sr</sub>	Inverse of the ratio of the 1-h geometric mean concentration to the emission flux along a straight road segment bisecting a square site (g/m <sup>2</sup> -s per kg/m <sup>3</sup> )	23.02 (for 0.5 acre site)	U.S. EPA 2002b Equation 5-5
T <sub>s</sub>	Total time over which construction occurs (s)	site-specific	U.S. EPA 2002b Equation 5-5
W	Mean vehicle weight (tons)	(number of cars x tons/car + number of trucks x tons/truck) / total vehicles)	U.S. EPA 2002b Equation 5-5
W <sub>R</sub>	Width of road segment (ft)	20	U.S. EPA 2002b Equation E-18
ΣVKT	Sum of fleet vehicle kilometers traveled during the exposure duration (km)	ΣVKT = total vehicles x distance (km/day) x frequency (weeks/year) x (days/year)	U.S. EPA 2002b Equation 5-5
PEF <sub>sc</sub>	Particulate Emission Factor - subchronic (m <sup>3</sup> /kg)	(site-specific)	U.S. EPA 2002b Equation 5-5

Mechanical Particulate Emission Factor Variables from Other Construction Activities			
Symbol ▼	Definition (units) ▼	Default ▼	Reference ▼
0.35	PM <sub>10</sub> particle size multiplier (unitless)	0.35	U.S. EPA 2002b Equation E-21
0.60	PM <sub>10</sub> scaling factor (unitless)	0.60	U.S. EPA 2002b Equation E-23
0.75	PM <sub>10</sub> scaling factor (unitless)	0.75	U.S. EPA 2002b Equation E-22
A	Dispersion constant (unitless)	2.4538	U.S. EPA 2002b Equation E-15
A <sub>C-doz</sub>	Areal extent of dozing (acres)	Site-specific	Necessary to solve ΣVKT <sub>doz</sub> in U.S. EPA 2002b Equation E-22
A <sub>C-till</sub>	Areal extent of tilling (acres)	Site-specific	U.S. EPA 2002b Equation E-24
A <sub>excav</sub>	Areal extent of excavation (m <sup>2</sup> )	(range 0.5 to 500)	U.S. EPA 2002b Equation E-21
A <sub>s</sub>	Areal extent of site surface soil contamination (acres)	(range 0.5 to 500)	U.S. EPA 2002b Equation E-15
A <sub>surf</sub>	Areal extent of site surface soil contamination (m <sup>2</sup> )	(range 0.5 to 500)	U.S. EPA 2002b Equation E-20
B	Dispersion constant (unitless)	17.5660	U.S. EPA 2002b Equation E-15
B <sub>d</sub>	Dozer blade length (m)	Site-specific	U.S. EPA 2002b Page E-28
B <sub>g</sub>	Grader blade length (m)	Site-specific	U.S. EPA 2002b Page E-28
C	Dispersion constant (unitless)	189.0426	U.S. EPA 2002b Equation E-15
d <sub>excav</sub>	Average depth of excavation (m)	Site-specific	U.S. EPA 2002b Equation E-21
ED	Exposure duration (years)	Site-specific	U.S. EPA 2002b Equation E-20

F(x)	Function Dependent on $0.886 \times (U_f/U_m)$ (unitless)	0.194	U.S. EPA. 1996, Appendix D Table 2
F <sub>D</sub>	Dispersion correction factor (unitless)	Site-specific	U.S. EPA 2002b Equation E-16
J <sub>T</sub> (g/m <sup>2</sup> -s)	Total time-averaged PM <sub>10</sub> unit emission flux for construction activities other than traffic on unpaved roads	Site-specific	U.S. EPA 2002b Equation E-25
M <sub>doz</sub>	Unit mass emitted from dozing operations (g)	site-specific	U.S. EPA 2002b Equation E-22
M <sub>excav</sub>	Unit mass emitted from excavation soil dumping (g)	site-specific	U.S. EPA 2002b Equation E-21
M <sub>grade</sub>	Unit mass emitted from grading operations (g)	site-specific	U.S. EPA 2002b Equation E-23
M <sub>m-doz</sub>	Gravimetric soil moisture content (%)	7.9 (mean value for overburden)	U.S. EPA 2002b Equation E-22
M <sub>m-excav</sub>	Gravimetric soil moisture content (%)	12 (mean value for municipal landfill cover)	U.S. EPA 2002b Equation E-21
M <sup>PC</sup> <sub>wind</sub>	Unit mass emitted from wind erosion (g)	site-specific	U.S. EPA 2002b Equation E-20
M <sub>till</sub>	Unit mass emitted from tilling operations (g)	site-specific	U.S. EPA 2002b Equation E-24
N <sub>A-doz</sub>	Number of times site is dozed (unitless)	Site-specific	U.S. EPA 2002b Equation E-22
N <sub>A-dump</sub>	Number of times soil is dumped (unitless)	2	U.S. EPA 2002b Equation E-21
N <sub>A-grade</sub>	Number of times site is graded (unitless)	Site-specific	U.S. EPA 2002b Equation E-23
N <sub>A-till</sub>	Number of times soil is tilled (unitless)	2	U.S. EPA 2002b Equation E-24
Q/C <sub>sa</sub>	Inverse of the ratio of the 1-h. geometric mean air concentration and the emission flux at the center of the square emission source (g/m <sup>2</sup> -s per kg/m <sup>3</sup> )	Site-specific	U.S. EPA 2002b Equation E-15
s <sub>doz</sub>	Soil silt content (%)	6.9	U.S. EPA 2002b Equation E-22
S <sub>doz</sub>	Average dozing speed (kph)	11.4 (mean value for graders)	U.S. EPA 2002b Equation E-22
S <sub>grade</sub>	Average grading speed (kph)	11.4 (mean value for graders)	U.S. EPA 2002b Equation E-23
s <sub>till</sub>	Soil silt content (%)	18	U.S. EPA 2002b Equation E-24
U <sub>m</sub>	Mean Annual Wind Speed (m/s)	4.69	U.S. EPA. 1996, Appendix D Table 2
U <sub>t</sub>	Equivalent Threshold Value of Wind Speed at 7m (m/s)	11.32	U.S. EPA. 1996, Appendix D Table 2
V	Fraction of Vegetative Cover (unitless)	0	U.S. EPA 2002b Equation E-20
ρ <sub>soil</sub>	In situ soil density (includes water) (Mg/m <sup>3</sup> )	1.68	U.S. EPA 2002b Equation E-21
ΣVKT <sub>doz</sub>	Sum of dozing kilometers traveled (km)	Site-specific	U.S. EPA 2002b Equation E-22
ΣVKT <sub>grade</sub>	Sum of grading kilometers traveled (km)		U.S. EPA 2002b Equation E-23
A <sub>c-grade</sub>	Areal extent of grading (acres)	Site-specific	Necessary to solve ΣVKT <sub>grade</sub> in U.S. EPA 2002b Equation E-23
PEF <sub>sc</sub>	Particulate Emission Factor - subchronic (m <sup>3</sup> /kg)	(site-specific)	U.S. EPA 2002b Equation E-26

## 6. References

ANL. 2001. [User's Manual for RESRAD Version 6](#). Argonne National Laboratory, Argonne, IL. ANL/EAD-4

AMEC. 2014. [Uptake of Gaseous Carbon-14 in the Biosphere](#): Development of an Assessment Model. AMEC/004041/007, Issue 2.

Calabrese E.J.; Barnes, R.; Stanek III, E.J.; Pastides, H.; Gilbert, C.E.; Veneman, P.; Wang, X.R.; Laszltity, A.; and Kostecki, P.T. 1989. How much soil do young children ingest: an epidemiologic study. Regul Toxicol Pharmacol: 10(2):123-137.

Davis, S.; Waller, P.; Buschbom, R.; Ballou, J.; and White, P. 1990. Quantitative estimates of soil ingestion in normal children between the ages of 2 and 7 years: Population-based estimates using aluminum, silicon, and titanium as soil tracer elements. Arch Environ Health 45: 112-122.

Environment Agency (E.A.). 2006. [Initial radiological assessment methodology - part 2 methods and input data](#). United Kingdom. SC030162/SR2.

E.A. 2009. [Updated technical background to the CLEA model](#). United Kingdom. SC050021/SR3. pg. 97 and pg. 100.

Etnier. 1980. Till, J. E., H. R. Meyer, E. L. Etnier, E. S. Bomar, R.D. Gentry, G. G. Killough, P. S. Rohwer, V. J. Tennery, and C. C. Travis [Tritium-An Analysis of Key Environmental and Dosimetric Questions](#). ORNL/TM-6990. pg. 15.

Hinton, T. G. 1992. Contamination of plants by resuspension: a review, with critique of measurement methods. Sci. Total Environ. 121:171-193

Hoffman, David J.; Rattner, Barnett A.; Burton, G. Allen jr.; Cairns, John jr. 2003. [Handbook of Ecotoxicology: Second Edition](#). Lewis Publishers.

IAEA. 1994. Handbook of Parameter Values for the Prediction of Radionuclide Transfer in Temperate Environments. International Atomic Energy Agency

ICRP. 2008. Nuclear Decay Data for Dosimetric Calculations. ICRP Publication 107. Ann. ICRP 38 (3).

IAEA. 2009. [Quantification of Radionuclide Transfer in Terrestrial and Freshwater Environments for Radiological Assessments](#). International Atomic Energy Agency. TECDOC1616.

IAEA. 2010. Technical Report Series No. 472: Handbook of Parameter Values for the Prediction of Radionuclide Transfer in Terrestrial and Freshwater Environments. International Atomic Energy Agency. TRS-472

Lyons, Robert K.; Machen, Rick; Forbes, T.D.A. 1999. [Understanding Forage Intake in Range Animals](#). AgriLIFE Extension. Texas A&M.

Lugo, A. E.; Brown, S.L.; Dodson, R.; Smith, T.S.; and Shugart, H.H. 1999. [The Holdridge life zones of the conterminous United States in relation to ecosystem mapping](#). Journal of Biogeography, 26: 1025-1038

Mller, Ch.W. 1980. An analysis of measured values for the fraction of a radiocative aerosol intercepted by vegetation. Health Physics, 38: 705 - 712.

NCRP. 1996. [Screening Models for Releases of Radionuclides to Atmosphere, Surface Water, and Ground. Vols. 1 and 2](#). NCRP Report No. 123. National Council on Radiation Protection and Measurements.

NEC. 1995. [Swine Nutrition Guide](#). Cooperative Extension Service / South Dakota State University and University of Nebraska / U.S. Department of Agriculture. Nebraska Cooperative Extension EC 95-273-C.

• The pig water ingestion numbers are derived from the USDA "Swine Nutrition Guide" and the EPA Human Health Risk Assessment Protocol for Hazardous Waste Combustion Facilities, [found here](#). USDA assumes a pig consumes 1/4 to 1/3 gallons of water for every pound of dry feed. The midpoint of this range (7/24 gallons of water per 1 lbs. of dry feed) was used with the default dry feed, (4.7 kg) from the U.S. EPA, to come up with 3 gallons (11.4 L) per day default water intake.

NRC. 1994. [Nutrient Requirements of Poultry: Ninth Revised Edition](#). Washington, DC: The National Academies Press, 1994.

OMAFRA. 2015. [Water Requirements of Livestock](#). Order Number 07-023. July, 2015.

ORNL. 2014a. [Area Correction Factors for Contaminated Soil for Use in Risk and Dose Assessment Models](#) and [appendix](#). Center for Radiation Protection Knowledge. September 2014.

ORNL. 2014b. [Gamma Shielding Factors for Soil Covered Contamination for Use in Risk and Dose Assessment Models](#) and [appendix](#). Center for Radiation Protection Knowledge. September 2014.

ORNL. 2014c. [Calculation of Slope Factors and Dose Coefficients](#) and [appendix](#). Center for Radiation Protection Knowledge. September 2014.

ORNL. 2016. [Biota Modeling in EPA's Preliminary Remediation Goal and Dose Compliance Concentration Calculators for Use in EPA Superfund Risk Assessment: Explanation of Intake Rate Derivation, Transfer Factor Compilation, and Mass Loading Factor Sources](#) Center for Radiation Protection Knowledge. November 2016.

Pinder, J. E. III. and K. W. McLeod. 1989. Mass loading of soil particles on plant surfaces, Health Phys. 57:935-942.

Tarr, Brian. 2011.[Guidelines to Feeding and Management of Dairy Goats](#) Nutrifax. Shur Gain.

U.S. EPA. 1988. Limiting Values of Radionuclide Intake and Air Concentration and Dose Conversion Factors for Inhalation, Submersion, and Ingestion. [Federal Guidance Report No. 11](#). Office of Radiation Programs, Washington, DC. EPA-520/1-88-020.

U.S. EPA. 1989. U.S. Environmental Protection Agency (U.S. EPA). Risk assessment guidance for Superfund. Volume I: Human health evaluation manual (Part A). Interim Final. Office of Emergency and Remedial Response. [EPA/540/1-89/002](#).

U.S. EPA. 1990. Interim Final Methodology for Assessing Health Risks Associated with Indirect Exposure to Combustor Emissions. Environmental Criteria and Assessment Office. ORD. EPA-600-90-003. January.

U.S. EPA. 1991a. U.S. Environmental Protection Agency (U.S. EPA). Human health evaluation manual, supplemental guidance: "Standard default exposure factors". OSWER Directive 9285.6-03.

U.S. EPA. 1991b. Risk Assessment Guidance for Superfund, Volume I: Human Health Evaluation Manual (Part B, Development of Risk-Based

Preliminary Remediation Goals). Office of Emergency and Remedial Response. [EPA/540/R-92/003](#). December 1991.

U.S. EPA. 1993. External Exposure to Radionuclides in Air, Water, and Soil. Federal Guidance Report No. 12. Office of Radiation and Indoor Air, Washington, DC. [EPA 402-R-93-081](#).

U.S. EPA. 1994a. Estimating Exposure to Dioxin-like Components - Volume III: Site-Specific Assessment Procedure. Review Draft. Office of Research and Development. Washington D.C. [EPA/600/6-88/005Cc](#). June. (Also see U.S. EPA. 1994c for direct link to Table).

U.S. EPA. 1994b. Radiation Site Cleanup Regulations: Technical Support Documents for the Development of Radiation Cleanup Levels for Soil (Review Draft). Office of Radiation and Indoor Air, Washington, DC. ["EPA 402-R-96-011A"](#). View Appendix C [here](#).

U.S. EPA. 1994c. Revised Draft Guidance for Performing Screening Level Risk Analyses at Combustion Facilities Burning Hazardous Wastes. [Attachment C](#), Draft Exposure Assessment Guidance for RCRA Hazardous Waste Combustion Facilities. Office of Emergency and Remedial Response. Office of Solid Waste. December 14.

U.S. EPA. 1996a. Soil Screening Guidance: User's Guide. Office of Emergency and Remedial Response. Washington, DC. [OSWER No. 9355.4-23](#).

U.S. EPA. 1996b. Soil Screening Guidance: Technical Background Document. Office of Emergency and Remedial Response. Washington, DC. [OSWER No. 9355.4-17A](#).

U.S. EPA. 1997a. Exposure Factors Handbook. Office of Research and Development, Washington, DC. [EPA/600/P-95/002Fa](#).

U.S. EPA. 1997b. Parameter Guidance Document. National Center for Environmental Assessment, NCEA-0238.

U.S. EPA. 1998. Human Health Risk Assessment Protocol for Hazardous Waste Combustion Facilities. Office of Solid Waste, Washington, DC. [EPA530-D-98-001A](#).

U.S. EPA. 1999a. [Data Collection for the Hazardous Waste Identification Rule](#). Office of Solid Waste, Washington, DC. The section on cattle fodder, soil and water intakes is [here](#).

U.S. EPA. 1999b. Understanding Variation in Partition Coefficient,  $K_d$ , Values, Volume II, "Review of Geochemistry and Available  $K_d$  Values for Cadmium, Cesium, Chromium, Lead, Plutonium, Radon, Strontium, Thorium, Tritium (3H), and Uranium". Office of Radiation and Indoor Air. Washington, DC. [EPA 402-R-99-004B](#). August 1999. .

U.S. EPA. 1999c. Cancer Risk Coefficients for Environmental Exposure to Radionuclides. Federal Guidance Report No. 13. Office of Radiation and Indoor Air. [EPA 402-R-99-001](#). September 1999.

U.S. EPA. 2000a. Soil Screening Guidance for Radionuclides: User's Guide. Office of Emergency and Remedial Response and Office of Radiation and Indoor Air. Washington, DC. [OSWER No. 9355.4-16A](#).

U.S. EPA. 2000b. Soil Screening Guidance for Radionuclides: Technical Background Document. Office of Emergency and Remedial Response and Office of Radiation and Indoor Air. Washington, DC. [OSWER No. 9355.4-16](#).

U.S. EPA. 2001. [Supplemental Guidance for Developing Soil Screening Levels for Superfund Sites](#). [OSWER 9355.4-24](#). March 2001.

U.S. EPA. 2002a. Role of Background in the CERCLA Cleanup Program. Office of Solid Waste and Emergency Response, April 26, 2002, [OSWER 9285.6-07P](#)

U.S. EPA. 2002b. Supplemental Guidance for Developing Soil Screening Levels for Superfund Sites. [OSWER 9355.4-24](#). December 2002.

U.S. EPA. 2002c. [Simulating Radionuclide Fate and Transport in the Unsaturated Zone: Evaluation and Sensitivity Analyses of Select Computer Models](#).

U.S. EPA. 2004. Risk Assessment Guidance for Superfund Volume I: Human Health Evaluation Manual (Part E, Supplemental Guidance for Dermal Risk Assessment) Final. OSWER 9285.7-02EP. July 2004. [Document](#) and [website](#).

U.S. EPA. 2005. [Human Health Risk Assessment Protocol for Hazardous Waste Combustion Facilities](#). Office of Solid Waste, Washington, DC.

U.S. EPA. 2011. [Exposure Factors Handbook 2011 Edition \(Final\)](#). National Center for Environmental Assessment, Office of Research and Development. Washington D.C.

van Wijnen J.H., Clausing P., and Brunekreef B. (1990); Estimated soil ingestion by children. Environ Res: 51(2): 147-162.

[Contact Us](#) to ask a question, provide feedback, or report a problem.